

हमारा विश्वास...
हर एक विद्यार्थी है स्वास

JEE ADVANCED
September 2020

ANSWER KEY

CHEMISTRY _ PAPER - 2

MOTION™

IIT/NIT | NEET | NTSE/IJSO/OLYMPIADS

Corporate Office : 394, Rajeev Gandhi Nagar, Kota
www.motion.ac.in | ✉: info@motion.ac.in

SECTION-1 (Maximum marks :18)


- This section contains **SIX (06)** questions.
- The answer to each question is a **SINGLE DIGIT INTEGER ranging from 0 TO 9, BOTH INCLUSIVE.**
- For Each Question, enter the correct integer corresponding to the answer using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme :
 Full marks : +3 If ONLY the correct integer is entered;
 Zero Marks : 0 If the question is unanswered.
 Negative Marks : -1 In all other cases.

1. The 1st, 2nd, and the 3rd ionization enthalpies, I_1 , I_2 and I_3 , of four atoms with atomic numbers n , $n + 1$, $n + 2$, and $n + 3$, where $n < 10$, are tabulated below. What is the value of n ?

Atomic number	Ionization Enthalpy (kJ/mol)		
	I_1	I_2	I_3
n	1681	3374	6050
$n + 1$	2081	3952	6122
$n + 2$	496	4562	6910
$n + 3$	738	1451	7733

Ans. 9

2. Consider the following compounds in the liquid form :
 O_2 , HF, H_2O , NH_3 , H_2O_2 , CCl_4 , $CHCl_3$, C_6H_6 , C_6H_5Cl .
 When a charged comb is brought near their flowing stream, how many of them show deflection as per the following figure ?



Ans. 6



ONLINE OFFLINE CLASSROOM

**REPEATER
BATCH**

JEE ADVANCED 2021
Starting from :
07th Oct. 2020


3. In the chemical reaction between stoichiometric quantities of KMnO_4 and KI in weakly basic solution, what is the number of moles of I_2 released for 4 moles of KMnO_4 consumed ?

Ans. 6

4. An acidified solution of potassium chromate was layered with an equal volume of amyl alcohol. When it was shaken after the addition of 1 mL of 3% H_2O_2 , a blue alcohol layer was obtained. The blue color is due to the formation of a chromium (VI) compound 'X'. What is the number of oxygen atoms bonded to chromium through only single bonds in a molecule of X ?

Ans. 4

5. The structure of a peptide is given below.



If the absolute values of the net charge of the peptide at $\text{pH} = 2$, $\text{pH} = 6$, and $\text{pH} = 11$ are $|z_1|$, $|z_2|$, and $|z_3|$, respectively, then what is $|z_1| + |z_2| + |z_3|$?

Ans. 5

6. An organic compound ($\text{C}_8\text{H}_{10}\text{O}_2$) rotates plane-polarized light. It produces pink color with neutral FeCl_3 solution. What is the total number of all the possible isomers for this compound ?

Ans. 6



ONLINE OFFLINE CLASSROOM

**REPEATER
BATCH**

JEE ADVANCED 2021

Starting from :
07th Oct. 2020


SECTION 2 (Maximum Marks : 24)

- Section contains **SIX (06)** questions.
- Each question has **FOUR** options. **ONE OR MORE THAN ONE** of these four option(s) is (are) correct answer(s).
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer the each question will be evaluated according to the following marking scheme:


Full Marks	: +4	If only (all the correct option(s) is (are) chosen;
Partial Marks	: +3	If all the four options are correct but ONLY three options are chosen;
Partial Marks	: +2	If three or more options are correct but ONLY two options are chosen, both of which are correct;
Partial Marks	: +1	If two or more options are correct but ONLY one option is chosen and it is a correct option;
Zero Marks	: 0	If none of the options is chosen (i.e. the question is unanswered);
Negative Marks	: -2	In all other cases.

7. In an experiment, m grams of a compound X (gas/liquid/solid) taken in a container is loaded in a balance as shown in figure I below. In the presence of a magnetic field, the pan with X is either deflected upwards (figure II), or deflected downwards (figure III), depending on the compound X. Identify the correct statement(s).


(I)
Balanced;
Magnetic field absent



(II)
Upward deflection;
Magnetic field present



(III)
Downward deflection;
Magnetic field present



- (A) If X is $\text{H}_2\text{O}(\text{l})$, deflection of the pan is upwards.
 (B) If X is $\text{K}_4[\text{Fe}(\text{CN})_6](\text{s})$, deflection of the pan is upwards.
 (C) If X is $\text{O}_2(\text{g})$, deflection of the pan is downwards.
 (D) If X is $\text{C}_6\text{H}_6(\text{l})$, deflection of the pan is downwards.

Ans. A,B,C








ONLINE OFFLINE CLASSROOM

REPEATER BATCH

JEE ADVANCED 2021
Starting from :
07th Oct. 2020

8. Which of the following plots is(are) correct for the given reaction ?
([P]₀ is the initial concentration of P)



Ans. **A**

9. Which among the following statement(s) is(are) true for the extraction of aluminium from bauxite ?
(A) Hydrated Al₂O₃ precipitates, when CO₂ is bubbled through a solution of sodium aluminate.
(B) Addition of Na₃AlF₆ lowers the melting point of alumina.
(C) CO₂ is evolved at the anode during electrolysis.
(D) The cathode is a steel vessel with a lining of carbon.

Ans. **A,B,C,D**

10. Choose the correct statement(s) among the following.
(A) SnCl₂ · 2H₂O is a reducing agent.
(B) SnO₂ reacts with KOH to form K₂[Sn(OH)₆].
(C) A solution of PbCl₂ in HCl contains Pb²⁺ and Cl⁻ ions.
(D) The reaction of Pb₃O₄ with hot dilute nitric acid to give PbO₂ is a redox reaction.

Ans. **A,B**




ONLINE OFFLINE CLASSROOM

**REPEATER
BATCH**

JEE ADVANCED 2021

Starting from :
07th Oct. 2020

11. Consider the following four compounds I, II, III, and IV.




Choose the correct statement(s).


- (A) The order of basicity is $II > I > III > IV$.
 (B) The magnitude of pK_b difference between I and II is more than that between III and IV.
 (C) Resonance effect is more in III than in IV
 (D) Steric effect makes compound IV more basic than III.

Ans. **C,D**

12. Consider the following transformations of a compound P.



Choose the correct options(s).



Ans. **B,C**

ONLINE OFFLINE CLASSROOM

**REPEATER
BATCH**

JEE ADVANCED 2021


Starting from :
07th Oct. 2020



SECTION 3 (Maximum Marks : 24)

- This section contains **SIX (06)** questions. The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value of the answer using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer. If the numerical value has more than two decimal places, **truncate/round-off** the value to **TWO** decimal places.
- Answer the each question will be evaluated according to the following marking scheme:
 Full Marks : +4 If **ONLY** the correct numerical value is entered;
 Zero Marks : 0 In all other cases.

- 13.** A solution of 0.1 M weak base (B) is titrated with 0.1 M of a strong acid (HA). The variation of pH of the solution with the volume of HA added is shown in the figure below. What is the pK_b of the base ? The neutralization reaction is given by $B + HA \rightarrow BH^+ + A^-$.



Ans. 3.3

- 14.** Liquids A and B form ideal solution for all compositions of A and B at 25 °C. Two such solutions with 0.25 and 0.50 mole fractions of A have the total vapor pressures of 0.3 and 0.4 bar, respectively. What is the vapor pressure of pure liquid B in bar ?

Ans. 0.2




ONLINE OFFLINE CLASSROOM

**REPEATER
BATCH**

JEE ADVANCED 2021


Starting from :
07th Oct. 2020

15. The figure below is the plot of potential energy versus internuclear distance (d) of H_2 molecule in the electronic ground state. What is the value of the net potential energy E_0 (as indicated in the figure) in kJ mol^{-1} , for $d = d_0$ at which the electron-electron repulsion and the nucleus-nucleus repulsion energies are absent? As reference, the potential energy of H atom is taken as zero when its electron and the nucleus are infinitely far apart. Use Avogadro constant as $6.023 \times 10^{23} \text{ mol}^{-1}$.



Ans. **-5242.42**

16. Consider the reaction sequence from P to Q shown below. The overall yield of the major product Q from P is 75%. What is the amount in grams of Q obtained from 9.3 mL of P? (Use density of P = 1.00 g mL^{-1} ; Molar mass of C = 12.0, H = 1.0, O = 16.0 and N = 14.0 g mol^{-1})



Ans. **18.6**

17. Tin is obtained from cassiterite by reduction with coke. Use the data given below to determine the minimum temperature (in K) at which the reduction of cassiterite by coke would take place. At 298 K: $\Delta_f H^\circ(\text{SnO}_2(\text{s})) = -581.0 \text{ kJ mol}^{-1}$, $\Delta_f H^\circ(\text{CO}_2(\text{g})) = -394.0 \text{ kJ mol}^{-1}$. $S^\circ(\text{SnO}_2(\text{s})) = 56.0 \text{ J K}^{-1} \text{ mol}^{-1}$, $S^\circ(\text{Sn}(\text{s})) = 52.0 \text{ J K}^{-1} \text{ mol}^{-1}$, $S^\circ(\text{C}(\text{s})) = 6.0 \text{ J K}^{-1} \text{ mol}^{-1}$, $S^\circ(\text{CO}_2(\text{g})) = 210.0 \text{ J K}^{-1} \text{ mol}^{-1}$. Assume that the enthalpies and the entropies are temperature independent.

Ans. **935**

18. An acidified solution of 0.05 M Zn^{2+} is saturated with $0.1 \text{ M H}_2\text{S}$. What is the minimum molar concentration (M) of H^+ required to prevent the precipitation of ZnS ? use $K_{\text{sp}}(\text{ZnS}) = 1.25 \times 10^{-22}$ and overall dissociation constant of H_2S , $K_{\text{NET}} = K_1 K_2 = 1 \times 10^{-21}$.

Ans. **0.2**



ONLINE OFFLINE CLASSROOM

REPEATER BATCH

JEE ADVANCED 2021
Starting from :
07th Oct. 2020

जब इन्होंने पूरा किया अपना सपना
तो आप भी पा सकते हैं लक्ष्य अपना

Admission
OPEN

JEE ADVANCED RESULT 2019



Aniket Agrawal

AIR-34



Dilendra Uikey

AIR-117



Karan

AIR-140



Shubham Kumar

AIR-174

KOTA'S PIONEER IN DIGITAL EDUCATION

1,95,00,000+ viewers | 72,67,900+ viewing hours | 2,11,000+ Subscribers

SERVICES	SILVER	GOLD	PLATINUM
Classroom Lectures (VOD)			
Live interaction	NA		
Doubt Support	NA		
Academic & Technical Support	NA		
Complete access to all content	NA		
Classroom Study Material	NA		
Exercise Sheets	NA		
Recorded Video Solutions	NA		
Online Test Series	NA		
Revision Material	NA		
Upgrade to Regular Classroom program	Chargeable	Chargeable	Free
Physical Classroom	NA	NA	
Computer Based Test	NA	NA	
Student Performance Report	NA	NA	
Workshop & Camp	NA	NA	
Motion Solution Lab- Supervised learning and instant doubt clearance	NA	NA	
Personalised guidance and mentoring	NA	NA	

FEE STRUCTURE

CLASS	SILVER	GOLD	PLATINUM
7th/8th	FREE	₹ 12,000	₹ 35,000
9th/10th	FREE	₹ 15,000	₹ 40,000
11th	FREE	₹ 29,999	₹ 49,999
12th	FREE	₹ 39,999	₹ 54,999
12th Pass	FREE	₹ 39,999	₹ 59,999

+ Student Kit will be provided at extra cost to Platinum Student.

- * **SILVER (Trial)** Only valid 7 DAYS or First 10 Hour's Lectures.
- ** **GOLD (Online)** can be converted to regular classroom (Any MOTION Center) by paying difference amount after lockdown.
- *** **PLATINUM (Online + Regular)** can be converted to regular classroom (Any MOTION Center) without any cost after lockdown.

New Batch Starting from :
07th Oct. 2020

Zero Cost EMI Available

MOTION™

H.O. : 394, Rajeev Gandhi Nagar, Kota
www.motion.ac.in | ✉ : info@motion.ac.in