

JEE ADVANCED ANSWER KEY

2021

MATHEMATICS
Paper-1

QUESTION WITH SOLUTION



32700+ SELECTIONS
SINCE 2007

Motion®

हो चुकी है ऑफलाइन क्लासरूम की शुरुआत
अपने सपने को करो साकार, कोटा कोविंग के साथ

Directors of Nucleus Education & Wizard of Mathematics

Now Offline associated with Motion Kota Classroom



Nitin Vijay (NV Sir)
Managing Director
Exp. : 18 yrs



Akhilesh Kanther (AKK Sir)
Exp. : 17 yrs

Vishal Joshi (VJ Sir)
Exp. : 18 yrs

Surendra K. Mishra (SKM Sir)
Exp. : 16 yrs



Gavesh Bhardwaj (GB Sir)
Exp. : 17 yrs

Academic Pillars of JEE Motion Kota



Ram Ratan Dwivedi (RRD Sir)
Joint Director
Exp. : 20 yrs



Amit Verma (AV Sir)
Joint Director
Exp. : 16 yrs



Vijay Pratap Singh (VPS Sir)
Vice President
Exp. : 20 yrs



Nikhil Srivastava (NS Sir)
Head JEE Academics
Exp. : 17 yrs



Aatish Agarwal (AA Sir)
Sr. Faculty
Exp. : 17 yrs



Jayant Chittora (JC Sir)
Sr. Faculty
Exp. : 16 yrs



Anurag Garg (AG Sir)
Sr. Faculty
Exp. : 17 yrs



Arjun Gupta (Arjun Sir)
Sr. Faculty
Exp. : 14 yrs



Devki Nandan Pathak (DN Sir)
Sr. Faculty
Exp. : 13 yrs



Avinash Kishore (AVN Sir)
Sr. Faculty
Exp. : 9 yrs



Vipin Sharma (VS Sir)
Sr. Faculty
Exp. : 12 yrs



Durgesh Pandey (Pandey Sir)
Sr. Faculty
Exp. : 8 yrs

Join

English & Hindi Medium

JEE DROPPER BATCH

Online + Offline Mode

Batch Starting from :
6th October 2021

SECTION – 1

- This section contains **FOUR (04)** questions.
 - Each question has **FOUR** options (A), (B), (C) and (D). **ONLY ONE** of these four options is the correct answer.
 - For each question, choose the option corresponding to the correct answer.
 - Answer to each question will be evaluated according to the following marking scheme:
- | | |
|----------------|---|
| Full Marks | : +3 If ONLY the correct option is chosen; |
| Zero Marks | : 0 If none of the options is chosen (i.e. the question is unanswered); |
| Negative Marks | : -1 In all other cases. |

Q.id 521118

Circle


1. Consider a triangle Δ whose two sides lie on the x-axis and the line $x+y+1=0$. If the orthocenter of Δ is $(1,1)$, then the equation of the circle passing through the vertices of the triangle Δ is

$(A) x^2 + y^2 - 3x + y = 0$	$(B) x^2 + y^2 + x + 3y = 0$
$(C) x^2 + y^2 + 2y - 1 = 0$	$(D) x^2 + y^2 + x + y = 0$

माना एक त्रिभुज की दो भुजाएँ x-अक्ष तथा रेखा $x+y+1=0$ पर स्थित हैं। यदि Δ का लम्बकेन्द्र $(1,1)$ है, तब त्रिभुज के शीर्ष से गुजरने वाले वृत्त का समीकरण है।

$(A) x^2 + y^2 - 3x + y = 0$	$(B) x^2 + y^2 + x + 3y = 0$
$(C) x^2 + y^2 + 2y - 1 = 0$	$(D) x^2 + y^2 + x + y = 0$

Ans. B



$$L_{BC} : x + y = 1 \Rightarrow L_{AM} \perp L_{BC}$$


$$L_{AM} : y - x = 0$$

$$\Rightarrow A(0,0) \text{ & } B : (-1,0)$$

$$L_{CN} \perp L_{AB}$$

$$L_{CN} : x = 1 \Rightarrow C : (1,-2)$$

$$\text{Now centroid of } \Delta \text{ is } C_1 : \left(0, -\frac{2}{3}\right)$$



$$\text{Equation of circle is } \left(x + \frac{1}{2}\right)^2 + \left(y + \frac{3}{2}\right)^2 = \left(\frac{\sqrt{10}}{2}\right)^2$$

$$S: x^2 + y^2 + x + 3y = 0$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



521139

Area under curve

2. The area of the region $\left\{ (x, y) : 0 \leq x \leq \frac{9}{4}, 0 \leq y \leq 1, x \geq 3y, x + y \geq 2 \right\}$ is

(A) $\frac{11}{32}$

(B) $\frac{35}{96}$

(C) $\frac{37}{96}$

(D) $\frac{13}{32}$

क्षेत्र $\left\{ (x, y) : 0 \leq x \leq \frac{9}{4}, 0 \leq y \leq 1, x \geq 3y, x + y \geq 2 \right\}$ का क्षेत्रफल है।


(A) $\frac{11}{32}$

(B) $\frac{35}{96}$

(C) $\frac{37}{96}$

(D) $\frac{13}{32}$

Ans. A



$$\text{Area} = \Delta_1 + \Delta_2$$

$$\Delta = \frac{1}{2} \begin{vmatrix} 3 & 1 & 1 \\ \frac{3}{2} & \frac{1}{2} & 1 \\ 2 & 0 & 1 \end{vmatrix} + \frac{1}{2} \begin{vmatrix} 2 & 0 & 1 \\ \frac{9}{4} & 0 & 1 \\ \frac{9}{4} & \frac{3}{4} & 1 \end{vmatrix}$$

$$\Delta = \frac{1}{2} \left[-2 \left(\frac{1}{2} - \frac{3}{4} \right) - 1 \left(\frac{3}{2} - \frac{3}{4} - \frac{9}{4} \cdot \frac{1}{2} \right) \right] + \frac{1}{2} \left| \frac{3}{4} \left(2 - \frac{9}{4} \right) \right|$$

$$\Delta = \frac{1}{2} \left| -2 \left(-\frac{1}{4} \right) - \left(\frac{9}{8} - \frac{9}{8} \right) \right| + \frac{1}{2} \left| \frac{3}{4} \left(-\frac{1}{4} \right) \right|$$

$$\Delta = \frac{1}{4} + \frac{3}{32} = \frac{8+3}{32} = \frac{11}{32}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



521121**Probability**

3. Consider three sets $E_1 = \{1, 2, 3\}$, $F_1 = \{1, 3, 4\}$ and $G_1 = \{2, 3, 4, 5\}$. Two elements are chosen at random, without replacement, from the set E_1 , and let S_1 denote the set of these chosen elements. Let $E_2 = E_1 - S_1$ and $F_2 = F_1 \cup S_1$. Now two elements are chosen at random, without replacement, from the set F_2 and let S_2 denote the set of these chosen elements.

Let $G_2 = G_1 \cup S_2$. Finally, two elements are chosen at random, without replacement, from the set G_2 and let S_3 denote the set of these chosen elements.

Let $E_3 = E_2 \cup S_3$. Given that $E_1 = E_3$, let p be the conditional probability of the event $S_1 = \{1, 2\}$.

Then the value of p is

- (A) $\frac{1}{5}$ (B) $\frac{3}{5}$ (C) $\frac{1}{2}$ (D) $\frac{2}{5}$

माना तीन समुच्चय $E_1 = \{1, 2, 3\}$, $F_1 = \{1, 3, 4\}$ तथा $G_1 = \{2, 3, 4, 5\}$ हैं। दो अवयवों समुच्चय E_1 से बिना प्रतिस्थापन के यादृच्छया चयनित किए जाते हैं तथा माना S_1 इन चयनित अवयवों के समुच्चय को निरूपित करता है। माना $E_2 = E_1 - S_1$ तथा $F_2 =$ अब दो अवयव समुच्चय F_2 से बिना प्रतिस्थापन के यादृच्छया चयनित किए जाते हैं तथा माना S_2 इन चयनित अवयवों के समुच्चय को निरूपित करता है। माना $G_2 =$ अन्ततः, दो अवयव समुच्चय G_2 से बिना प्रतिस्थापन के यादृच्छया चयनित किए जाते हैं तथा माना S_3 इन चयनित अवयवों के समुच्चय को निरूपित करता है। माना $E_3 =$ दिया है कि $E_1 = E_3$, माना p घटना $S_1 = \{1, 2\}$ की प्रतिबंधात्मक प्रायिकता है। तब p का मान है

- (A) $\frac{1}{5}$ (B) $\frac{3}{5}$ (C) $\frac{1}{2}$ (D) $\frac{2}{5}$

Ans. A

$$P = \frac{P(S_1 \cap (E_1 = E_2))}{P(E_2 = E_3)} = \frac{P(B_{1/2})}{P(B)}$$

$$P(B) = P(B_{1,2}) + P(B_{1,3}) + P(B_{2,3})$$

↑ ↑ ↑
 If 1.2 If 1.3 If 2.3
 chosen chosen chosen
 at start at start at start

$$P(B_{1,2}) = \frac{1}{3} \times \underbrace{\frac{1 \times {}^2C_1}{{}^3C_2}}_{\substack{= \\ =}} \times \underbrace{\frac{1}{{}^5C_2}}_{\substack{=}}$$

$$P(B_{2,3}) = \frac{1}{3} \times \left[\underbrace{\frac{{}^3C_2 \times 1}{{}^4C_2}}_{\substack{= \\ =}} \times \frac{1}{{}^4C_2} + \underbrace{\frac{1 \times {}^3C_3}{{}^4C_2}}_{\substack{= \\ =}} \times \frac{1}{{}^5C_2} \right]$$

$$\frac{P(B_{1,2})}{P(B)} = \frac{1}{5}$$

521126 Complex No.

4. Let $\theta_1, \theta_2, \dots, \theta_{10}$ be positive valued angles (in radian) such that

$\theta_1 + \theta_2 + \dots + \theta_{10} = 2\pi$. Define the complex numbers $z_1 = e^{i\theta_1}, z_k = z_{k-1} e^{i\theta_k}$

for $k=2, 3, \dots, 10$, where $i = \sqrt{-1}$. Consider the statements P and Q given below:



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



$$P : |z_2 - z_1| + |z_3 - z_2| + \dots + |z_{10} - z_9| + |z_1 - z_{10}| \leq 2\pi$$

$$Q : |z_2^2 - z_1^2| + |z_3^2 - z_2^2| + \dots + |z_{10}^2 - z_9^2| + |z_1^2 - z_{10}^2| \leq 4\pi$$

Then,

(A) P is **TRUE** and Q is **FALSE**

(B) Q is **TRUE** and P IS **FALSE**

(C) Both P and Q are **TRUE**

(D) Both P and Q are **FALSE**

माना $\theta_1, \theta_2, \dots, \theta_{10}$ धनात्मक मान वाले कोण (रेडियन में) हैं जबकि $\theta_1 + \theta_2 + \dots + \theta_{10} = 2\pi$. $k=2, 3, \dots, 10$ के लिए सम्मिश्र संख्याओं को परिभासित कीजिए, जहाँ नीचे दिये गये कथनों P तथा Q पर विचार कीजिए:

$$P : |z_2 - z_1| + |z_3 - z_2| + \dots + |z_{10} - z_9| + |z_1 - z_{10}| \leq 2\pi$$

$$Q : |z_2^2 - z_1^2| + |z_3^2 - z_2^2| + \dots + |z_{10}^2 - z_9^2| + |z_1^2 - z_{10}^2| \leq 4\pi$$

Then,

(A) P सही है तथा Q गलत है

(B) Q सही है तथा P गलत है

(C) P तथा Q दोनों सही हैं

(D) P तथा Q दोनों गलत हैं

Ans. C

$$z_k = z_{k-1}, e^{i\theta_k}$$


$$|z_{k+1} - z_k| = \text{side of polygon's}$$

$$P = |z_2 - z_1| + |z_3 - z_2| + \dots + |z_{10} - z_9| + |z_1 - z_{10}|$$

P = Sum of sides of polygon

P \leq perimeter of circumference

$$\Rightarrow [P \leq 2\pi]$$



$$\theta : |z_2^2 - z_1^2| + |z_3^2 - z_2^2| + \dots + |z_{10}^2 - z_9^2|$$

$$\theta : |z_2 - z_1| \frac{|z_2 + z_1|}{\leq 2} + |z_3 - z_2| \frac{|z_3 + z_1|}{\leq 2} + \dots + |z_{10} - z_9| \frac{|z_{10} + z_1|}{\leq 2}$$

$$\theta \leq 2(|z_2 - z_1| + |z_3 - z_2| + \dots + |z_{10} - z_9|)$$

$$\theta \leq 2.2\pi$$

$$\boxed{\theta \leq 4\pi}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



Section – 2

- This section contains **THREE (03)** question stems.
- There are **TWO (02)** questions corresponding to each question stem.
- The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value corresponding to the answer in the designated place using the mouse and the on-screen virtual numeric keypad.
- If the numerical value has more than two decimal places, **truncate/round-off** the value to **TWO** decimal places.
- Answer to each question will be evaluated according to the following marking scheme:
Full Marks : +2 If ONLY the correct numerical value is entered at the designated place;
Zero Marks : 0 In all other cases.

Question Stem for Question Nos. 5 and 6

Question Stem

Three numbers are chosen at random, one after another with replacement, from the set $S = \{1, 2, 3, \dots, 100\}$. Let p_1 be the probability that the maximum of chosen numbers is at least 81 and p_2 be the probability that the minimum of chosen numbers is at most 40.

प्रश्न संख्या 5 तथा 6 के लिए अनुच्छेद

समुच्चय $S = \{1, 2, 3, \dots, 100\}$ से तीन संख्याएँ यादृच्छया एक के बाद एक प्रतिस्थापन सहित चयनित की जाती है। माना p_1 अधिकतम चयनित संख्या कम से कम 81 होने की प्रायिकता है तथा p_2 न्यूनतम चयनित संख्या अधिक से अधिक 40 होने की प्रायिकता है। तब

521141

Probability

5. The value of $\frac{625}{4} p_1$ is _____.

$\frac{625}{4} p_1$ का मान _____ है।

Ans. 76.25

521142

Probability

6. The value of $\frac{125}{4} p_2$ is _____.

$\frac{125}{4} p_2$ का मान _____ है।

Ans. 24.5

$P_1 = \dots$ at least 81

$$P_1 = 1 - (< 81)$$

$$= 1 - \left(\left(\frac{80}{100} \right)^3 \right) = 1 - \left(\frac{4}{5} \right)^3 = \frac{125 - 64}{125} = \frac{61}{125}$$

$P_2 = \dots$ minimum chosen number

$$P_2 = 1 - \left(\frac{60}{100} \right)^3 = 1 - \left(\frac{3}{5} \right)^3 = \frac{125 - 27}{125} = \frac{98}{125}$$

$$\frac{625}{4} \cdot P_1 = \frac{625}{4} \cdot \frac{61}{125} = \frac{61}{4} = 76.25$$

$$\frac{125}{4} \cdot P_2 = \frac{125}{4} \cdot \frac{98}{125} = 24.5$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



Question Stem for Question Nos. 7 and 8

Question Stem

Let α , β and γ be real numbers such that the system of linear equations

$$\begin{aligned}x + 2y + 3z &= \alpha \\4x + 5y + 6z &= \beta \\7x + 8y + 9z &= \gamma - 1\end{aligned}$$

is consistent. Let $|M|$ represent the determinant of the matrix

$$M = \begin{bmatrix} \alpha & 2 & \gamma \\ \beta & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}$$

Let P be the plane containing all those (α, β) for which the above system of linear equations is consistent, and D be the square of the distance of the point $(0, 1, 0)$ from the plane P .

प्रश्न संख्या 7 व 8 के लिए अनुच्छेद

माना α , β व γ वास्तविक संख्याएँ जबकि ϵ एखिक समीकरण निकाय

$$\begin{aligned}x + 2y + 3z &= \alpha \\4x + 5y + 6z &= \beta \\7x + 8y + 9z &= \gamma - 1\end{aligned}$$

संसार के सभी विषयों को निरूपित करता है। माना $|M|$ निम्न आव्यह के सारणिक को निरूपित करता है

$$M = \begin{bmatrix} \alpha & 2 & \gamma \\ \beta & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}$$

माना P उन सभी (α, β, γ) का तल है जिनके लिए उपरोक्त रेखिक समीकरण निकाय सुसंगत है तथा D समतल P से बिन्दु $(0,1,0)$ की दूरी का वर्ग है।

521386

DETERMINANT

7. The value of $|M|$ is

ग्रन्थालय |

Ans. 1

$$\Delta = \begin{vmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{vmatrix} = 0$$

\Rightarrow System of equation has ' ∞ ' solution.

$$\Delta_1 = \begin{vmatrix} \alpha & 2 & 3 \\ \beta & 5 & 6 \\ \gamma - 1 & 8 & 9 \end{vmatrix} = 0$$

$$-3\alpha + 6\beta - 3(\gamma - 1) = 0$$

$$-\alpha + 2\beta - \gamma + 1 = 0$$

$$\alpha - 2\beta + \gamma = 1 \quad \dots \dots \dots \quad (1)$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



$$\Delta_2 = \begin{vmatrix} 1 & \alpha & 3 \\ 4 & \beta & 6 \\ 7 & \gamma - 1 & 9 \end{vmatrix} = 0$$

$$-\alpha(36 - 42) + \beta(9 - 21) - (\gamma - 1)(6 - 12) = 0$$

$$6\alpha - 12\beta - 6(\gamma - 1) = 0$$

$$\alpha - 2\beta + \gamma = 1 \quad \dots \dots \dots (2)$$

$$\Delta_3 = \begin{vmatrix} 1 & 2 & \alpha \\ 4 & 5 & \beta \\ 7 & 8 & \gamma - 1 \end{vmatrix} = 0$$

$$\alpha(-3) - \beta(-6) + (\gamma - 1)(-3) = 0$$

$$\alpha - 2\beta + \gamma = 1 \quad \dots \dots \dots (3)$$

$|M|$

$$= \alpha - 2(\beta) + \gamma(1)$$

$$= \alpha - 2\beta + \gamma$$

$$= 1$$

521392

8. D का मान _____ है।

Ans. 1.5

(α, β, γ) lie on plane

$x - 2y + z = 1$ hence its distance
from $(0, 1, 0)$ is

$$D = \left| \frac{-2 - 1}{\sqrt{1 + 4 + 1}} \right| = \frac{3}{\sqrt{6}}$$

$$D^2 = \frac{9}{6} = \frac{3}{2} = 1.5$$

Question Stem for Question Nos. 9 and 10

Question Stem

Consider the lines L_1 and L_2 defined by

$$L_1 : x\sqrt{2} + y - 1 = 0 \text{ and } L_2 : x\sqrt{2} - y + 1 = 0$$

For a fixed constant λ , let C be the locus of a point P such that the product of the distance of P from L_1 and the distance of P from L_2 is λ^2 . The line $y = 2x + 1$ meets C at two points R and S, where the distance between R and S is $\sqrt{270}$.

Let the perpendicular bisector of RS meet C at two distinct points R' and S' . Let D be the square of the distance between R' and S' .

प्रश्न संख्या 9 व 10 के लिए अनुच्छेद

माना रेखाएँ L_1 व L_2 , $L_1 : x\sqrt{2} + y - 1 = 0$ व $L_2 : x\sqrt{2} - y + 1 = 0$ द्वारा परिभाषित है।

एक निश्चित अचर λ के लिए, माना C एक बिन्दुपथ है जबकि P की L_1 से दूरी तथा P की L_2 से दूरी का गुणनफल λ^2 है। रेखा $y = 2x + 1$, C से दो बिंदुओं R तथा S पर मिलती है, जहाँ R व S के मध्य दूरी $\sqrt{270}$ है।

माना RS का लम्ब समद्विभाजक C से दो भिन्न बिंदुओं R' व S' पर मिलता है। माना R' व S' के मध्य दूरी का वर्ग D है।

[JEE 2021]



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



522138**Straight Line**

9. The value of λ^2 is _____.
 λ^2 का मान _____ है।

Ans. 9

$$d_1 \cdot d_2 = \lambda^2 \quad \text{Let } P : (h, k)$$

$$\left| \frac{h\sqrt{2} + k - 1}{\sqrt{3}} \right| \left| \frac{h\sqrt{2} - k + 1}{\sqrt{3}} \right| = \lambda^2$$

$$\left| \sqrt{2}x + y - 1 \right| \left| \sqrt{2}x - y + 1 \right| = 3\lambda^2 \dots\dots\dots(1)$$

Now solve this with $y = 2x + 1$ to

Let R & S

$$\left| x\sqrt{2} + 2x + 1 - 1 \right| \left| x\sqrt{2} - 2x - 1 + 1 \right| = 3\lambda^2$$

$$2|x|^2 = 3\lambda^2$$

$$\lambda^2 = \frac{2x^2}{3}$$

$x_1 = \sqrt{\frac{3}{2}}\lambda \Rightarrow y_1 = \sqrt{6}\lambda + 1$

$x_2 = -\sqrt{\frac{3}{2}}\lambda \Rightarrow y_2 = 1 - \sqrt{6}\lambda$

$$R: \left(\frac{\sqrt{3}}{2}\lambda, \sqrt{6}\lambda + 1 \right), S: \left(-\frac{\sqrt{3}}{2}\lambda, 1 - \sqrt{6}\lambda \right)$$

$$(x_1 - x_2)^2 + (y_1 - y_2)^2 = 270$$

$$\Rightarrow (\sqrt{6}\lambda)^2 + (2\sqrt{6}\lambda)^2 = 270$$

$$6\lambda^2 + 24\lambda^2 = 270 > 0$$

$$\lambda^2 = \frac{270}{30}$$

$$\Rightarrow \boxed{\lambda^2 = 9}$$

522146**Straight Line**

10. The value of D is _____.
D का मान _____ है।

Ans. 77.14

\perp bisector of RS

$$T = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

Here $x_1 + x_2 = 0$

$T = (0, 1)$.

Equation of



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



$$R'S''(y-1) = -\frac{1}{2}(x-0) \Rightarrow x+2y=2$$

$R'(a_1, b)$ $S(a_2, b_2)$

$$D = (a_1 - a_2)^2 + (b_1 - b_2)^2 = 5(b_1 - b_2)^2$$

Solve $x + 2y = 2$ and $|2x^2 - (y-1)^2| = 3\lambda^2$

$$\left| 8(y-1)^2 - (y-1)^2 \right| = 3\lambda^2 \Rightarrow (y-1)^2 = \left(\frac{\sqrt{3}\lambda}{\sqrt{7}} \right)^2$$

$$y-1 = \pm \frac{\sqrt{3}\lambda}{\sqrt{7}} \Rightarrow b_1 = 1 + \frac{\sqrt{3}\lambda}{\sqrt{7}}, b_2 = 1 - \frac{\sqrt{3}\lambda}{\sqrt{17}}$$

$$D = 5 \left(\frac{2\sqrt{3}\lambda}{\sqrt{7}} \right)^2 = \frac{5 \times 4 \times 3\lambda^2}{7} = \frac{5 \times 4 \times 27}{7} = 77.14$$

Section – 3

- This section contains SIX (06) questions.
- Each question has FOUR options (A), (B), (C) and (D). ONE OR MORE THAN ONE of these four option(s) is (are) correct answer(s).
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme:

Full Marks	: +4 If only (all) the correct option(s) is(are) chosen;
Partial Marks	: +3 If all the four options are correct but ONLY three options are chosen;
Partial Marks	: +2 If three or more options are correct but ONLY two options are chosen, both of which are correct;
Partial Marks	: +1 If two or more options are correct but ONLY one option is chosen and it is a correct option;
Zero Marks	: 0 If unanswered;
Negative Marks	: -2 In all other cases.
- For example, in a question, if (A), (B) and (D) are the ONLY three options corresponding to correct answers, then
 - choosing ONLY (A), (B) and (D) will get +4 marks;
 - choosing ONLY (A) and (B) will get +2 marks;
 - choosing ONLY (A) and (D) will get +2 marks;
 - choosing ONLY (B) and (D) will get +2 marks;
 - choosing ONLY (A) will get +1 mark;
 - choosing ONLY (B) will get +1 mark;
 - choosing ONLY (D) will get +1 mark;
 - choosing no option(s) (i.e. the question is unanswered) will get 0 marks and choosing any other option(s) will get -2 marks.

522171 Matrix

11. For any 3×3 matrix M, let $|M|$ denote the determinant of M. Let

$$E = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 8 & 13 & 18 \end{bmatrix}, P = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \text{ and } F = \begin{bmatrix} 1 & 3 & 2 \\ 8 & 18 & 13 \\ 2 & 4 & 3 \end{bmatrix}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



If Q is a nonsingular matrix of order 3×3 , then which of the following statements is (are) **TRUE?**

- (A) $F = PEP$ and $P^2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$
- (B) $|EQ + PFQ^{-1}| = |EQ| + |PFQ^{-1}|$
- (C) $|(EF)^3| > |EF|^2$
- (D) Sum of the diagonal entries of $P^{-1}EP + F$ is equal to the sum of diagonal entries of $E = P^{-1}FP$

किसी 3×3 आव्यूह M के लिए, माना $|M|, M$ के सारणिक को निरूपित करता है। माना

$$E = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 8 & 13 & 18 \end{bmatrix}, P = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \text{ and } F = \begin{bmatrix} 1 & 3 & 2 \\ 8 & 18 & 13 \\ 2 & 4 & 3 \end{bmatrix}$$

यदि Q कोटि 3×3 का एक व्युत्क्रमणीय है, तब निम्न में से कौनसा / कौनसे सही है / हैं?

- (A) $F = PEP$ तथा $P^2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$
- (B) $|EQ + PFQ^{-1}| = |EQ| + |PFQ^{-1}|$
- (C) $|(EF)^3| > |EF|^2$
- (D) $P^{-1}EP + F$ के विकर्ण अवयवों का योगफल $E = P^{-1}FP$ के विकर्ण अवयवों के योगफल के बराबर है

Ans. A,B,D

$$PEP = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 8 & 13 & 18 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

$$= \begin{pmatrix} 1 & 2 & 3 \\ 8 & 13 & 18 \\ 2 & 3 & 4 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 3 & 2 \\ 8 & 18 & 13 \\ 2 & 4 & 3 \end{pmatrix}$$

$$P^2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 10 \\ 0 & 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

(B) $|EQ + PFQ^{-1}| = |EQ| + |PFQ^{-1}|$

$|E| = 0$ and $|F| = 0$ and $|Q| \neq 0$

$$|EQ| = |E||Q| = 0, |PFQ^{-1}| = \frac{|P||F|}{|Q|} = 0$$

$T = EQ + PFQ^{-1}$

$TQ = EQ^2 + PF = EQ^2 + P^2EP = EQ^2 + EP = E(Q^2 + P)$

$|TQ| = |E(Q^2 + P)| \Rightarrow |T||Q| = |E||Q^2 + P| = 0 \Rightarrow |T| = 0$ (as $|Q| \neq 0$)

(C) $|(EF)^3| > |EF|^2$

Here $0 > 0$ (false)



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



(D) as $P^2 = I \Rightarrow P^{-1} = P$ so $P^{-1}FP = PFP = PPEPP = E$

So $E + P^{-1}FP = E + E = 2E$

$P^{-1}EP + F \Rightarrow PEP + F = 2PEP$

$\text{Tr}(2PEP) = 2\text{Tr}(PEP) = 2\text{Tr}(EPP) = 2\text{Tr}(E)$

522175

Monotonocity

12. Let $f : R \rightarrow R$ be defined by

$$f(x) = \frac{x^2 - 3x - 6}{x^2 + 2x + 4}$$

Then which of the following statements is (are) **TRUE**?

(A) f is decreasing in the interval $(-2, -1)$

(B) f is increasing in the interval $(1, 2)$

(C) f is onto

(D) Range of f is $\left[-\frac{3}{2}, 2\right]$

माना $f : R \rightarrow R$ निम्न प्रकार परिभाषित है:

$$f(x) = \frac{x^2 - 3x - 6}{x^2 + 2x + 4}$$

[JEE 2021]

तब निम्न में से कौनसा/कौनसे कथन सही है/हैं?

(A) f , अन्तराल $(-2, -1)$ से हासमान है


(B) f , अन्तराल $(1, 2)$ में वर्धमान है

(C) f आच्छादक है

(D) f का परिसर $\left[-\frac{3}{2}, 2\right]$ है

Ans. A,B

$$y = \frac{x^2 - 3x - 6}{x^2 + 2x + 4}$$



$$\frac{dy}{dx} = \frac{(x^2 + 2x + 4)(2x - 3) - (x^2 - 3x - 6)(2x + 2)}{(x^2 + 2x + 4)^2}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



$$\begin{aligned}
 &= \frac{[2x^3 + 4x^2 + 8x - 3x^2 - 6x - 12] - [2x^3 - 6x^2 - 12x + 2x^2 - 6x - 12]}{[x^2 + 2x + 4]^2} \\
 &= \frac{5x^2 + 20x}{(x^2 + 2x + 4)^2} \\
 &\quad \begin{array}{c|c|c}
 + & - & + \\
 \hline
 -4 & 0
 \end{array}
 \end{aligned}$$

Increasing in $(-\infty, -4] \cup [0, \infty)$

Decreasing in $[-4, 0]$

$$\text{Range} \Rightarrow x^2y + 2xy + 4y = x^2 - 3x - 6$$

$$\Rightarrow x^2(y - 1) + x(2y + 3) + 4y + 6 = 0$$

Case I $\rightarrow y \neq 1, D \geq 0$

$$(2y + 3)^2 - 4(y - 1)(4y + 6) \geq 0$$

$$\Rightarrow (4y^2 + 9 + 12) - 4[4y^2 + 2y - 6] \geq 0$$

$$\Rightarrow -12y^2 + 4y + 33 \geq 0$$

$$\Rightarrow 12y^2 - 4y - 33 \leq 0$$

$$\Rightarrow (6y - 11)(2y + 3) \leq 0$$

$$y \in \left[-\frac{3}{2}, \frac{11}{6}\right] - \{1\}$$

Case-II $\rightarrow y = 1$

$$x^2 + 2x + 4 = x^2 - 3x - 6$$

$$\Rightarrow 5x = -10$$

$x = -2$ from case -I and case - II

Ans. $y \in \left[-\frac{3}{2}, \frac{11}{6}\right]$

522252

Probability

13. Let E, F and G be three events having probabilities

$$P(E) = \frac{1}{8}, P(F) = \frac{1}{6} \text{ and } P(G) = \frac{1}{4}, \text{ and let } P(E \cap F \cap G) = \frac{1}{10}.$$

For any event H, if H^c denotes its complement, then which of the following statements is (are) TRUE?

(A) $P(E \cap F \cap G^c) \leq \frac{1}{40}$

(B) $P(E^c \cap F \cap G) \leq \frac{1}{15}$

(C) $P(E \cap F \cap G) \leq \frac{13}{24}$

(D) $P(E^c \cap F^c \cap G^c) \leq \frac{5}{12}$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



माना E, F व G तीन घटनाएँ हैं जिनकी प्रायिकताएँ $P(E) = \frac{1}{8}$, $P(F) = \frac{1}{6}$ तथा $P(G) = \frac{1}{4}$, तथा $P(E \cap F \cap G) = \frac{1}{10}$.

किसी घटना H के लिए, यदि H^c इसके पूरक को निरूपित करता है, तब निम्न में से कौनसा/कौनसे कथन सही है/हैं?

- | | |
|--|--|
| (A) $P(E \cap F \cap G^c) \leq \frac{1}{40}$ | (B) $P(E^c \cap F \cap G) \leq \frac{1}{15}$ |
| (C) $P(E \cap F \cap G) \leq \frac{13}{24}$ | (D) $P(E^c \cap F^c \cap G^c) \leq \frac{5}{12}$ |

Ans. A,B,C

$$P(6) = \frac{1}{B}, P(F) = \frac{1}{6}, P(C_1) = \frac{1}{4}, P(E \cap F \cap G) = \frac{1}{10}$$

$$(A) P(E \cap F \cap G^c) = \therefore \{P(E \cap F) \leq P(E)\}$$

$$\begin{aligned} &\leq P(E) - P(G) \\ &\leq \frac{1}{8} - \frac{1}{10} \\ &\leq \frac{1}{40} \end{aligned}$$

$$(B) P(E^c \cap F \cap G) \therefore \{P(E \cap F) \leq P(E)\}$$

$$\begin{aligned} &\leq P(F) - P(G) \\ &\leq \frac{1}{6} - \frac{1}{10} \\ &\leq \frac{10 - 6}{60} = \frac{4}{60} \\ &\leq \frac{1}{15} \end{aligned}$$

$$(C) P(E \cup F \cup G) \leq P(E) + P(F) + P(G)$$

$$\begin{aligned} &\leq \frac{1}{8} + \frac{1}{6} + \frac{1}{4} \\ &\leq \frac{15 + 20 + 30}{120} = \frac{65}{120} = \frac{13}{24} \end{aligned}$$

$$(D) P(E^c \cap F^c \cap G^c) = 1 - P(E \cup F \cup G)$$

$$\begin{aligned} &\geq 1 - \frac{13}{24} \\ &\geq \frac{11}{24} \end{aligned}$$

522263 Matrix

14. For any 3×3 matrix M , let $|M|$ denote the determinant of M . Let I be the 3×3 identity matrix. Let E and F be two 3×3 matrices such that $(I - EF)$ is invertible. If $G = (I - EF)^{-1}$, then which of the following statements is (are) **TRUE**?

- | | |
|----------------------------|-----------------------------|
| (A) $ FE = I - FE FGE $ | (B) $(I - FE)(I + FGE) = I$ |
| (C) $EFG = GEF$ | (D) $(I - FE)(I - FGE) = I$ |

किसी 3×3 आव्यूह M के लिए $|M|, M$ के सारणिक को निरूपित करता है। माना $I, 3 \times 3$ तत्समक आव्यूह है। माना E व F दो 3×3 आव्यूह हैं जबकि $(I - EF)$ प्रतिलोमीय है। यदि $G = (I - EF)^{-1}$, तब निम्न में से कौनसा/कौनसे कथन सही है/हैं?

- | | |
|----------------------------|-----------------------------|
| (A) $ FE = I - FE FGE $ | (B) $(I - FE)(I + FGE) = I$ |
| (C) $EFG = GEF$ | (D) $(I - FE)(I - FGE) = I$ |



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



Ans. A,B,C

$$G = (I - EF)^{-1}$$

$$G^{-1} = (I - EF)$$

$$GG^{-1} = G - GEF$$

$$I = G - GEF$$

$$\text{Also } I = G - EFG$$

$$\Rightarrow GEF = EFG \rightarrow \boxed{C}$$

$$(B) (I - FE)(I + FGE) = I - FE + FGE - FEFGE$$

$$= I - FE + FGE - F(G - I)E$$

$$= I - FE + FGE - FGE + FE$$

$$(D) |FE| = |I - FE| |FGE|$$

$$\text{Now } (I - FE)(FGE)$$

$$= FGE - FEFGE$$

$$= FGE - FGEEFE$$

$$= FGE - F(G - I)E$$

$$= FGE - FGE + FE$$

$$= FE$$

$$\Rightarrow |I - FE| |FGE| = |FE|$$

If B is correct then D is not correct.

522274

Inverse Trigonometric Functions

15. For any positive integer n , let $S_n: (0, \infty) \rightarrow \mathbb{R}$ be defined by

$$S_n(x) = \sum_{k=1}^n \cot^{-1} \left(\frac{1 + k(k+1)x^2}{x} \right),$$

where for any $x \in \mathbb{R}$, $\cot^{-1}(x) \in (0, \pi)$ and $\tan^{-1}(x) \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$. Then which of the following statements is (are) **TRUE**?

$$(A) S_{10}(x) = \frac{\pi}{2} - \tan^{-1} \left(\frac{1 + 11x^2}{10x} \right), \text{ for all } x > 0$$

$$(B) \lim_{n \rightarrow \infty} \cot(S_n(x)) = x, \text{ for all } x > 0$$

$$(C) \text{The equation } S_3(x) = \frac{\pi}{4} \text{ has a root in } (0, \infty)$$

$$(D) \tan(S_n(x)) \leq \frac{1}{2}, \text{ for all } n \geq 1 \text{ and } x > 0$$

किसी धनात्मक पूर्णांक n के लिए, माना

$$S_n: (0, \infty) \rightarrow \mathbb{R}$$

$$S_n(x) = \sum_{k=1}^n \cot^{-1} \left(\frac{1 + k(k+1)x^2}{x} \right),$$

द्वारा परिभाषित है जहाँ किसी R के लिए $\cot^{-1}(x) \in (0, \pi)$ तथा

$$\tan^{-1}(x) \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right). \text{ तब निम्न में से कौनसा / कौनसे कथन सही है / हैं?}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



(A) सभी $x > 0$ के लिए $S_{10}(x) = \frac{\pi}{2} - \tan^{-1}\left(\frac{1+11x^2}{10x}\right)$

(B) सभी $x > 0$ के लिए, $\lim_{n \rightarrow \infty} \cot(S_n(x)) = x$

(C) समीकरण $S_3(x) = \frac{\pi}{4}$ का एक मूल $(0, \infty)$ में है

(D) $\tan(S_n(x)) \leq \frac{1}{2}$, सभी $n \geq 1$ तथा $x > 0$ के लिए

Ans. A,B

$$\begin{aligned} S_n(x) &= \sum_{k=1}^n \tan^{-1} \left[\frac{x}{1 + (kx)(k+1)x} \right] \\ &= \sum_{k=1}^n \tan^{-1} \left[\frac{(k+1)x - kx}{1 + (kx)(k+1)x} \right] \\ &= \sum_{k=1}^n [\tan^{-1}((k+1)x) - \tan^{-1}(kx)] \\ &= [\tan^{-1} 2x - \tan^{-1} x] + [\tan^{-1} 3x - \tan^{-1} 2x] + [\tan^{-1} 4x - \tan^{-1} 3x] + \dots + [\tan^{-1}(n+1)x - \tan^{-1} nx] \end{aligned}$$

$$S_n(x) = \tan^{-1}(n+1)x - \tan^{-1}x$$

$$\text{Now, } S_{10}(x) = \tan^{-1}x(11x) - \tan^{-1}x$$

$$= \tan^{-1} \left[\frac{10x}{1 + (11x)(x)} \right]$$

$$= \tan^{-1} \left[\frac{10x}{1 + 11x^2} \right]$$

$$= \frac{\pi}{2} - \cot^{-1} \left(\frac{10x}{1 + 11x^2} \right)$$

$$= \frac{\pi}{2} - \tan^{-1} \left(\frac{1 + 11x^2}{10x} \right) \quad (x > 0)$$

(b) $\lim_{n \rightarrow \infty} \cot[\sin(x)] = \lim_{n \rightarrow \infty} \cot \left(\tan \left[\frac{(n+1)x - x}{1 + (n+1)x \cdot x} \right] \right)$

$$\lim_{n \rightarrow \infty} \cot \left[\cot^{-1} \left(\frac{1 + (n+1)x \cdot x}{nx} \right) \right]$$

$$\lim_{n \rightarrow \infty} \frac{1 + (n+1)x^2}{(nx)} \Rightarrow x$$

$$S_n(x) = \tan^{-1}(n+1)x - \tan^{-1}x$$

$$S_3(x) = \tan^{-1}(4x) - \tan^{-1}x$$

$$\Rightarrow \tan^{-1} \left[\frac{3x}{1 + 4x^2} \right] = \frac{\pi}{4} \quad (\text{given})$$

$$\Rightarrow \frac{3x}{1 + 4x^2} = 1$$

$$\Rightarrow 1 + 4x^2 = 3x$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



$$\Rightarrow 4x^2 - 3x + 1 = 0$$

$$D = 9 - 16 < 0$$

No real roots

$$(d) \tan(S_n(x)) \leq \frac{1}{2}$$

$$\tan\left[\tan^{-1}\left(\frac{nx}{1+(n+1)x^2}\right)\right] = \frac{nx}{1+(n+1)x^2}$$

$$\lim_{n \rightarrow \infty} \frac{nx}{1+(n+1)x^2} = \frac{1}{x}$$

522286

Complex Number

16. For any complex number $w = c + id$, let $\arg(w) \in (-\pi, \pi]$, where $i = \sqrt{-1}$. Let α and β be real numbers such that for all complex numbers $z = x + iy$ satisfying $\arg\left(\frac{z+\alpha}{z+\beta}\right) = \frac{\pi}{4}$, the ordered pair (x, y) lies on the circle

$$x^2 + y^2 + 5x - 3y + 4 = 0$$

Then which of the following statements is (are) **TRUE** ?

- (A) $\alpha = -1$ (B) $\alpha\beta = 4$ (C) $\alpha\beta = -4$ (D) $\beta = 4$

किसी सम्मिश्र संख्या $w = c + id$ के लिए माना

$\arg(w) \in (-\pi, \pi]$, जहाँ $i = \sqrt{-1}$. माना α तथा β वास्तविक संख्याएँ हैं जबकि सभी सम्मिश्र संख्याओं $z = x + iy$ के लिए $\arg\left(\frac{z+\alpha}{z+\beta}\right) = \frac{\pi}{4}$ को संतुष्ट करती है, क्रमित युग्म (x, y) वृत्त $x^2 + y^2 + 5x - 3y + 4 = 0$ पर स्थित है। तब निम्न में


से कौनसा/कौनसे कथन सही है/है?

- (A) $\alpha = -1$ (B) $\alpha\beta = 4$ (C) $\alpha\beta = -4$ (D) $\beta = 4$

Ans. B,D

$$S : x^2 + y^2 + 5x - 3y + 4 = 0$$

$$C : \left(\frac{-5}{2}, \frac{3}{2}\right) \quad R = \sqrt{\frac{25}{4} + \frac{9}{4} - 4} = \frac{3}{\sqrt{12}}$$



$$\arg\left(\frac{z - (-\alpha)}{z - (-\beta)}\right) = \frac{\pi}{4}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



$\Rightarrow A : (-1, 0) \text{ & } B : (-4, 0)$

$\alpha = 1, \text{ & } \beta = 4$

Alternate:

Let $Z = x + iy$

$$\tan^{-1} \left[\frac{y}{x+\alpha} \right] - \tan^{-1} \left[\frac{y}{x+\beta} \right] = \frac{\pi}{4}$$

$$\tan^{-1} \left[\frac{\frac{y}{x+\alpha} - \frac{y}{x+\beta}}{1 + \frac{y^2}{(x+\alpha)(x+\beta)}} \right] = \frac{\pi}{4}$$

$$\frac{\frac{y(\beta-\alpha)}{(x+\alpha)(x+\beta)}}{\frac{(x+\alpha)(x+\beta)+y^2}{(x+\alpha)(x+\beta)}} = 1$$

$$y(\beta-\alpha) = x^2 + (\alpha+\beta)x + \alpha\beta + y^2$$

$$\Rightarrow x^2 + y^2 + (\alpha+\beta)x - (\beta-\alpha)y + \alpha\beta = 0$$

$$\begin{array}{l} \alpha+\beta=5 \\ \alpha-\beta=-3 \end{array} \quad \alpha\beta=4$$

$$\alpha=1$$

$$\beta=4$$

SECTION 4

- This section contains **THREE (03)** questions.
- The answer to each question is a **NON-NEGATIVE INTEGER**.
- For each question, enter the correct integer corresponding to the answer using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme:
Full Marks : +4 If ONLY the correct integer is entered;
Zero Marks : 0 In all other cases.

522289

Quadratic Equation

- 17.** For $x \in \mathbb{R}$, the number of real roots of the equation $3x^2 - 4|x^2 - 1| + x - 1 = 0$ is _____.

$x \in \mathbb{R}$ के लिए, समीकरण $3x^2 - 4|x^2 - 1| + x - 1 = 0$ के वास्तविक मूलों की संख्या है _____.

Ans. 4

$$3x^2 - 4|x^2 - 1| + x - 1 = 0$$

$$3x^2 + x - 1 = 4|x^2 - 1|$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



$$x^2 \geq 1 \Rightarrow 3x^2 + x - 1 = 4 \times 2 - 4$$

$$x^2 - x - 3 = 0$$

$$x = \frac{1 + \sqrt{13}}{2}$$

$$x = \frac{1 - \sqrt{13}}{2}$$

$$\Rightarrow 3x^2 + x - 1 = 4 - 4x^2$$


$$x = \frac{-1 + \sqrt{141}}{14}$$

$$x = \frac{-1 - \sqrt{141}}{14}$$

$$7x^2 + x - 5 = 0$$

$$3x^2 + x - 1 = 4 |x^2 - 1|$$

Alternate method



522293

Solution of Triangle

18. In a triangle ABC, let AB = $\sqrt{23}$, BC = 3 and CA = 4. Then the value of $\frac{\cot A + \cot C}{\cot B}$ is ____.

एक त्रिभुज ABC में AB = $\sqrt{23}$, BC = 3 तथा CA = 4 तब $\frac{\cot A + \cot C}{\cot B}$ का मान ____ है।

Ans. 2

$$AB = c = \sqrt{23}$$

$$BC = a = 3$$

$$CA = b = 4$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



$$\cot A = \frac{b^2 + c^2 - a^2}{2bcKa}$$

$$\cot B = \frac{c^2 + a^2 - b^2}{2cakb}$$

$$\cot C = \frac{a^2 + b^2 - c^2}{2abkc}$$

$$\frac{\cot A + \cot C}{\cot B}$$

$$\frac{b^2 + c^2 - a^2 + a^2 + b^2 - c^2}{c^2 + a^2 - b^2}$$

$$= \frac{2.16}{23 + 9 - 16}$$

$$= \frac{32}{16} = [2]$$

522298

Vector

19. Let \vec{u}, \vec{v} and \vec{w} , be vectors in three-dimensional space, where \vec{u} and \vec{v} are unit vectors which are not perpendicular to each other and $\vec{u} \cdot \vec{w} = 1$, $\vec{v} \cdot \vec{w} = 1$, $\vec{w} \cdot \vec{w} = 4$

If the volume of the parallelopiped, whose adjacent sides are represented by the vectors \vec{u}, \vec{v} and \vec{w} , is $\sqrt{2}$ then the value of $|3\vec{u} + 5\vec{v}|$ is _____.

माना \vec{u}, \vec{v} व \vec{w} त्रिविम में सदिश हैं जहाँ \vec{u} व \vec{v} इकाई सदिश हैं जो एक दूसरे के लम्बवत् नहीं हैं तथा $\vec{u} \cdot \vec{w} = 1$, $\vec{v} \cdot \vec{w} = 1$, $\vec{w} \cdot \vec{w} = 4$ यदि समान्तर षटफलक जिसकी आसन्न भुजाएँ सदिश \vec{u}, \vec{v} व \vec{w} , द्वारा निरूपित की गयी हैं, का आयतन $\sqrt{2}$ है, तब $|3\vec{u} + 5\vec{v}|$ का मान _____ है।

Ans. 7

$$\vec{u} \cdot \vec{u} = 1, \vec{u} \cdot \vec{w} = 1, |\vec{w}| = 2$$

$$\vec{v} \cdot \vec{v} = 1, \vec{v} \cdot \vec{w} = 1$$

$$\vec{u} \cdot \vec{v} \neq 0$$

$$\text{Volume of parallelopiped} = [\vec{u} \cdot \vec{v} \cdot \vec{w}] = \sqrt{2}$$

$$(\sqrt{2})^2 = \begin{vmatrix} \vec{u} \cdot \vec{u} & \vec{u} \cdot \vec{v} & \vec{u} \cdot \vec{w} \\ \vec{v} \cdot \vec{u} & \vec{v} \cdot \vec{v} & \vec{v} \cdot \vec{w} \\ \vec{w} \cdot \vec{u} & \vec{w} \cdot \vec{v} & \vec{w} \cdot \vec{w} \end{vmatrix}$$

$$\text{Let } \vec{u} \cdot \vec{v} = \lambda \neq 0$$

$$2 = \begin{vmatrix} 1 & \lambda & 1 \\ \lambda & 1 & 1 \\ 1 & 1 & 4 \end{vmatrix}$$

$$2 = (4 - 1) - \lambda(4\lambda - 1) + 1(\lambda - 1)$$

$$2 = 3 - 4\lambda^2 + \lambda + \lambda - 1$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



$$4\lambda^2 - 2\lambda = 0$$

$\lambda=0$ (not possible)

$\lambda=\frac{1}{2}$

$$\begin{aligned} \text{Now } |3\bar{u} + 5\bar{v}| &= \sqrt{9 + 25 + 30 \cdot \frac{1}{2}} \\ &= \sqrt{9 + 25 + 15} \\ &= \sqrt{9 + 40} \\ &= \sqrt{49} = 7 \end{aligned}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**



हो चुकी है ऑफलाइन क्लासरूम की शुरुआत
अपने सपने को करो साकार, कोटा कोचिंग के साथ

Directors of Nucleus Education & Wizard of Mathematics

Now Offline associated with Motion Kota Classroom



Nitin Vijay (NV Sir)
Managing Director
Exp. : 18 yrs



Akhilesh Kanther
(AKK Sir)
Exp. : 17 yrs

Vishal Joshi
(VJ Sir)
Exp. : 18 yrs

Surendra K. Mishra
(SKM Sir)
Exp. : 16 yrs



Gavesh Bhardwaj
(GB Sir)
Exp. : 17 yrs

Academic Pillars of JEE Motion Kota



Ram Ratan Dwivedi
(RRD Sir)
Joint Director
Exp. : 20 yrs



Amit Verma
(AV Sir)
Joint Director
Exp. : 16 yrs



Vijay Pratap Singh
(VPS Sir)
Vice President
Exp. : 20 yrs



Nikhil Srivastava
(NS Sir)
Head JEE Academics
Exp. : 17 yrs



Aatish Agarwal
(AA Sir)
Sr. Faculty
Exp. : 17 yrs



Jayant Chittora
(JC Sir)
Sr. Faculty
Exp. : 16 yrs



Anurag Garg
(AG Sir)
Sr. Faculty
Exp. : 17 yrs



Arjun Gupta
(Arjun Sir)
Sr. Faculty
Exp. : 14 yrs



Devki Nandan Pathak
(DN Sir)
Sr. Faculty
Exp. : 13 yrs



Avinash Kishore
(AVN Sir)
Sr. Faculty
Exp. : 9 yrs



Vipin Sharma
(VS Sir)
Sr. Faculty
Exp. : 12 yrs



Durgesh Pandey
(Pandey Sir)
Sr. Faculty
Exp. : 8 yrs

Join

English & Hindi Medium

JEE DROPPER BATCH

Online + Offline Mode

Batch Starting from :
6th October 2021



KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**





An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **6th October 2021**

