


हमारा विश्वास... हर एक विद्यार्थी है सवास

**JEE  
MAIN  
JAN  
2020**

**PAPER WITH SOLUTION**

**7<sup>th</sup> January 2020 \_ SHIFT - 1**

**MATHEMATICS**



**24000+**  
SELECTIONS SINCE 2007

JEE (Advanced)

**5392**

(Under 50000 Rank)

JEE (Main)

**16241**

NEET / AIIMS

**1305**

(since 2016)

NTSE / OLYMPIADS

**1158**

(5<sup>th</sup> to 10<sup>th</sup> class)

**MOTION™**

Nurturing potential through education

H.O. : 394, Rajeev Gandhi Nagar, Kota

www.motion.ac.in | ✉: info@motion.ac.in

1. Let  $x^k + y^k = a^k$ , ( $a, k > 0$ ) and  $\frac{dy}{dx} + \left(\frac{y}{x}\right)^{\frac{1}{3}} = 0$ , then  $k$  is:

यदि  $x^k + y^k = a^k$ , ( $a, k > 0$ ) तथा  $\frac{dy}{dx} + \left(\frac{y}{x}\right)^{\frac{1}{3}} = 0$ , तो  $k$  बराबर है :

- (1)  $\frac{2}{3}$                       (2)  $\frac{4}{3}$                       (3)  $\frac{3}{2}$                       (4)  $\frac{1}{3}$

**Sol. 1**

$$k \cdot x^{k-1} \cdot k \cdot y^{k-1} \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = -\left(\frac{x}{y}\right)^{k-1}$$

$$\frac{dy}{dx} + \left(\frac{x}{y}\right)^{k-1} = 0$$

$$k - 1 = -\frac{1}{3}$$

$$k = 1 - \frac{1}{3} = \frac{2}{3}$$

2. Let  $\alpha$  be a root of the equation  $x^2 + x + 1 = 0$  and the matrix  $A = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 & 1 & 1 \\ 1 & \alpha & \alpha^2 \\ 1 & \alpha^2 & \alpha^4 \end{bmatrix}$ , then the matrix  $A^{31}$  is equal to

यदि समीकरण  $x^2 + x + 1 = 0$  का एक मूल  $\alpha$  है तथा आव्यूह  $A = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 & 1 & 1 \\ 1 & \alpha & \alpha^2 \\ 1 & \alpha^2 & \alpha^4 \end{bmatrix}$  है, तो आव्यूह  $A^{31}$  बराबर है

- (1)  $A$                       (2)  $A^3$                       (3)  $A^2$                       (4)  $I_3$

**Sol. 2**

$$A^2 = \frac{1}{3} \begin{bmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

$$\Rightarrow A^4 = 1$$

$$\Rightarrow A^{31} = A^{28} \times A^3 = A^3$$

3. Let  $\alpha$  and  $\beta$  be two real roots of the equation  $(k + 1)\tan^2 x - \sqrt{2} \cdot \lambda \tan x = (1 - k)$ , where  $k(\neq -1)$  and  $\lambda$  are real numbers. if  $\tan^2(\alpha + \beta) = 50$ , then a value of  $\lambda$  is:

माना समीकरण  $(k + 1)\tan^2 x - \sqrt{2} \cdot \lambda \tan x = (1 - k)$ ,  $k(\neq -1)$ ,  $\lambda \in \mathbb{R}$  के  $\alpha$  तथा  $\beta$  दो वास्तविक मूल हैं। यदि  $\tan^2(\alpha + \beta) = 50$  है, तो  $\lambda$  का एक मान है -

- (1) 5                      (2) 10                      (3)  $10\sqrt{2}$                       (4)  $5\sqrt{2}$

**Sol. 2**

$$(k + 1) \tan^2 x - \sqrt{2}\lambda \tan x + (k - 1) = 0$$

$$\tan \alpha + \tan \beta = \frac{\sqrt{2}\lambda}{k+1}$$

$$\tan \alpha \times \tan \beta = \frac{k-1}{k+1}$$

$$\tan(\alpha + \beta) = \frac{\frac{\sqrt{2}\lambda}{k+1}}{1 - \frac{k-1}{k+1}} = \frac{\sqrt{2}\lambda}{2} = \frac{\lambda}{\sqrt{2}}$$

$$\tan^2(\alpha + \beta) = \frac{\lambda^2}{2} = 50$$

$$\lambda = 10$$

4. A vector  $\vec{a} = \alpha\hat{i} + 2\hat{j} + \beta\hat{k}$  ( $\alpha, \beta \in \mathbb{R}$ ) lies in the plane of the vectors,  $\vec{b} = \hat{i} + \hat{j}$  and  $\vec{c} = \hat{i} - \hat{j} + 4\hat{k}$ . If  $\vec{a}$  bisects the angle between  $\vec{b}$  and  $\vec{c}$ , then:

एक सदिश  $\vec{a} = \alpha\hat{i} + 2\hat{j} + \beta\hat{k}$  ( $\alpha, \beta \in \mathbb{R}$ ) उस समतल में, जिसमें दोनों सदिश  $\vec{b} = \hat{i} + \hat{j}$  तथा  $\vec{c} = \hat{i} - \hat{j} + 4\hat{k}$  स्थित है, स्थित है। यदि  $\vec{a}$  सदिशों  $\vec{b}$  और  $\vec{c}$  के बीच के कोण को समद्विभाजित करता है तो :

$$(1) \vec{a} \cdot \hat{i} + 1 = 0$$

$$(2) \vec{a} \cdot \hat{k} + 2 = 0$$

$$(3) \vec{a} \cdot \hat{j} + 3 = 0$$

$$(4) \vec{a} \cdot \hat{k} + 4 = 0$$

**Sol. 2**

angle bisector can be  $\vec{a} = \lambda(\vec{b} + \vec{c})$  or  $\vec{a} = \mu(\vec{b} - \vec{c})$

$$\vec{a} = \lambda \left( \frac{\hat{i} + \hat{j}}{\sqrt{2}} + \frac{\hat{i} + \hat{j} + 4\hat{k}}{3\sqrt{2}} \right) = \frac{\lambda}{3\sqrt{2}} [3\hat{i} + 3\hat{j} + \hat{i} - \hat{j} + 4\hat{k}] = \frac{\lambda}{3\sqrt{2}} [4\hat{i} + 2\hat{j} + 4\hat{k}]$$

compare with  $\vec{a} = \alpha\hat{i} + 2\hat{j} + \beta\hat{k}$

$$\frac{2\lambda}{3\sqrt{2}} = 2 \Rightarrow \lambda = 3\sqrt{2}$$

$$\vec{a} = 4\hat{i} + 2\hat{j} + 4\hat{k}$$

Not in option so now consider  $\vec{a} = \mu \left( \frac{\hat{i} + \hat{j}}{\sqrt{2}} - \frac{\hat{i} - \hat{j} + 4\hat{k}}{3\sqrt{2}} \right)$

$$\vec{a} = \frac{\mu}{3\sqrt{2}} (3\hat{i} + 3\hat{j} - \hat{i} + \hat{j} - 4\hat{k})$$

$$= \frac{\mu}{3\sqrt{2}} (2\hat{i} + 4\hat{j} - 4\hat{k})$$

**Increase Your Score  
for JEE Main April'2020**

**उत्कर्ष**  
15 JAN 2020

percentile between 97.0 to 98.99  
in JEE Main (Jan-2020)

Fees - ₹ 22000 Including GST

**उन्नति**  
17 JAN 2020

Below 97 percentile in JEE Main (Jan-2020)

Tenure: 62 Days | Schedule: 5 Classes Per Day

Fees - ₹ 27500 Including GST

**उत्थान**  
17 JAN 2020

99 percentile and above  
in JEE Main (Jan-2020)

Fees - ₹ 11000  
score 160-200

Fees - ₹ 5500  
score 200-240

Fees - ₹ 0  
score above 240

compare with  $\vec{a} = \alpha \hat{i} + 2\hat{j} + \beta \hat{k}$

$$\frac{4\mu}{3\sqrt{2}} = 2 \Rightarrow \mu = \frac{3\sqrt{2}}{2} \Rightarrow \vec{a} = \hat{i} + 2\hat{j} - 2\hat{k}$$

$$\vec{a} \cdot \hat{k} + 2 = 0$$

5. If  $\operatorname{Re}\left(\frac{z-1}{2z+i}\right) = 1$ , where  $z = x + iy$ , then the point  $(x, y)$  lies on a:

यदि  $\operatorname{Re}\left(\frac{z-1}{2z+i}\right) = 1$ , जहाँ  $z = x + iy$ , तो बिन्दु  $(x, y)$  स्थित है :

(1) straight line whose slope is  $\frac{3}{2}$ .

(2) circle whose center is  $\left(-\frac{1}{2}, -\frac{3}{2}\right)$ .

(3) circle whose diameter is  $\frac{\sqrt{5}}{2}$

(4) straight line whose slope is  $-\frac{2}{3}$ .

(1) एक सरल रेखा पर, जिसका ढाल  $\frac{3}{2}$  है।

(2) एक वृत्त पर, जिसका केन्द्र बिन्दु  $\left(-\frac{1}{2}, -\frac{3}{2}\right)$  है।

(3) एक वृत्त पर, जिसका व्यास  $\frac{\sqrt{5}}{2}$  है।

(4) एक सरल रेखा पर, जिसका ढाल  $-\frac{2}{3}$  है।

**Sol. 3**

$$z = x + iy$$

$$\left(\frac{z-1}{2z+i}\right) = \frac{(x-1)+iy}{2(x+iy)+i} = \frac{(x-1)+iy}{2x+(2y+1)i} \times \frac{2x-(2y+1)i}{2x-(2y+1)i}$$

$$\operatorname{Re}\left(\frac{z+1}{2z+i}\right) = \frac{2x(x-1)+y(2y+1)}{(2x)^2+(2y+1)^2} = 1$$

$$\Rightarrow 2x^2 + 2y^2 - 2x + y = 4x^2 + 4y^2 + 4y + 1$$

$$\Rightarrow 2x^2 + 2y^2 + 2x + 3y + 1 = 0$$

$$\Rightarrow x^2 + y^2 - x + \frac{3}{2}y + \frac{1}{2} = 0$$

Circle with center  $\left(-\frac{1}{2}, -\frac{3}{4}\right)$

$$r = \sqrt{\frac{1}{4} + \frac{9}{16} - \frac{1}{2}} = \sqrt{\frac{4+9-8}{16}} = \frac{\sqrt{5}}{4}$$

6. Total number of 6-digit numbers in which only and all the five digits 1, 3, 5, 7 and 9 appear, is:  
छः अंकों वाली सभी संख्याओं की कुल संख्या जिनमें केवल तथा सभी पाँच अंक 1, 3, 5, 7 और 9 ही हों, है:

(1)  $\frac{1}{2}(6!)$

(2)  $6!$

(3)  $\frac{5}{2}(6!)$

(4)  $5^6$

**24000+**  
SELECTIONS SINCE 2007

JEE (Advanced)

**5392**

JEE (Main)

**16241**

(Under 50000 Rank)

NEET / AIIMS

**1305**

(since 2016)

NTSE / OLYMPIADS

**1158**

(5th to 10th class)

H.O. : 394, Rajeev Gandhi Nagar, Kota

Toll Free : 1800-212-1799

www.motion.ac.in | ✉ info@motion.ac.in



**Sol. 3**

1, 3, 5, 7, 9

For digit to repeat we have  ${}^5C_1$  choices

and six digits can be arranged in  $\frac{6!}{2}$  ways.

Hence total such numbers =  $\frac{5!}{2} = \frac{5 \cdot 6!}{2}$

**7.** If  $y = mx + 4$  is a tangent to both the parabolas,  $y^2 = 4x$  and  $x^2 = 2by$ , then  $b$  is equal to:

यदि  $y = mx + 4$  दोनो परवलयों,  $y^2 = 4x$  तथा  $x^2 = 2by$  को स्पर्श करती है, तो  $b$  बराबर है :

(1) 128                      (2) -32                      (3) -128                      (4) -64

**Sol. 3**

$y = mx + 4$

.....(i)

$y^2 = 4x$  tangent  $y = mx + \frac{a}{m} \Rightarrow y = mx + \frac{1}{m}$

.....(ii)

from (i) and (ii)

$4 = \frac{1}{m} \Rightarrow m = \frac{1}{4}$

So line  $y = \frac{1}{4}x + 4$  is also tangent to parabola  $x^2 = 2by$ , so solve

$x^2 = 2b \left( \frac{x+16}{4} \right) \Rightarrow 2x^2 - bx - 16b = 0 \Rightarrow d = 0$

$\Rightarrow b^2 - 4 \times 2 \times (-16b) = 0 \Rightarrow b^2 + 32 \times 4b = 0$

$b = -128, b = 0$  (not possible)

**8.** If the distance between the foci of an ellipse is 6 and the distance between its directrices is 12, then the length of its latus rectum is:

यदि एक दीर्घवृत्त की नाभियों के बीच की दूरी 6 है तथा इसकी नियताओं के बीच की दूरी 12 है, तो इसकी नाभिलम्ब जीवा की लम्बाई है :

(1)  $3\sqrt{2}$

(2)  $2\sqrt{3}$

(3)  $\frac{3}{\sqrt{2}}$

(4)  $\sqrt{3}$

**Sol. 1**

$2ae = 6$                       and                       $\frac{2a}{e} = 12$

$\Rightarrow ae = 3$                       and                       $\frac{a}{e} = 6$

$\Rightarrow a^2 = 18$

$\Rightarrow b^2 = a^2 - a^2e^2 = 18 - 9 = 9$

$\Rightarrow \text{L.R.} = \frac{2b^2}{a} = \frac{2 \times 9}{3\sqrt{2}} = 3\sqrt{2}$

**Increase Your Score  
for JEE Main April'2020**

**उन्नति**  
17 JAN 2020

Below 97 percentile in JEE Main (Jan-2020)

Tenure: 62 Days | Schedule: 5 Classes Per Day

Fees - ₹ 27500 Including GST

**उत्कर्ष**  
15 JAN 2020

percentile between 97.0 to 98.99  
in JEE Main (Jan-2020)

Fees - ₹ 22000 Including GST

**उत्थान**  
17 JAN 2020

99 percentile and above  
in JEE Main (Jan-2020)

Fees - ₹ 11000 | Fees - ₹ 5500 | Fees - ₹ 0  
score 160-200 | score 200-240 | score above 240


9. The area of the region, enclosed by the circle  $x^2 + y^2 = 2$  which is not common to the region bounded by the parabola  $y^2 = x$  and the straight line  $y = x$ , is:

वक्त  $x^2 + y^2 = 2$  द्वारा परिबद्ध क्षेत्र का वह क्षेत्रफल जो परवलय  $y^2 = x$  तथा सरल रेखा  $y = x$  द्वारा परिबद्ध क्षेत्र में नहीं है, है

- (1)  $\frac{1}{3}(6\pi - 1)$       (2)  $\frac{1}{6}(12\pi - 1)$       (3)  $\frac{1}{3}(12\pi - 1)$       (4)  $\frac{1}{6}(24\pi - 1)$

Sol. 2

Total area - enclosed area



$$2\pi - \int_0^1 [\sqrt{x} - x] dx$$

$$2\pi - \left( \frac{2x^{3/2}}{3} - \frac{x^2}{2} \right)_0^1$$

$$2\pi - \left( \frac{2}{3} - \frac{1}{2} \right) \Rightarrow 2\pi - \left( \frac{1}{6} \right) \Rightarrow \frac{12\pi - 1}{6}$$

10. An unbiased coin is tossed 5 times. Suppose that a variable X is assigned the value of k when k consecutive heads are obtained for k = 3, 4, 5, otherwise X takes the value -1. Then the expected value of X, is:

एक अनभिन्न सिक्के को पाँच बार उछाला जाता है। माना, एक चर X को, k = 3, 4, 5 के लिए, मान k दिया जाता है जब सिक्के पर क्रमागत k चित आएँ तथा अन्य सभी स्थितियों में X का मान -1 है, तो X का अपेक्षित मान है।

- (1)  $\frac{1}{8}$       (2)  $-\frac{3}{16}$       (3)  $\frac{3}{16}$       (4)  $-\frac{1}{8}$

Sol. 1

K	0	1	2	3	4	5
P(k)	$\frac{1}{32}$	$\frac{12}{32}$	$\frac{11}{32}$	$\frac{5}{32}$	$\frac{2}{32}$	$\frac{1}{32}$

k = no. of times head occur consecutively  
now expected value is

$$= \sum xP(k) = (-1) \times \frac{1}{32} + (-1) \times \frac{12}{32} + (-1) \times \frac{11}{32} + 3 \times \frac{5}{32} + 4 \times \frac{2}{32} + 5 \times \frac{1}{32} = \frac{1}{8}$$

- 11.** Let the function,  $f: [-7, 0] \rightarrow \mathbb{R}$  be continuous on  $[-7, 0]$  and differentiable on  $(-7, 0)$ . If  $f(-7) = -3$  and  $f'(x) \leq 2$ , for all  $x \in (-7, 0)$ , then for all such functions  $f$ ,  $f(-1) + f(0)$  lies in the interval:  
माना फलन  $f: [-7, 0] \rightarrow \mathbb{R}$ ,  $[-7, 0]$  पर संतत है तथा  $(-7, 0)$  पर अवकलनीय है। यदि  $f(-7) = -3$  और सभी  $x \in (-7, 0)$  के लिए,  $f'(x) \leq 2$  है, तो ऐसे सभी फलनों  $f$  के लिए  $f(-1) + f(0)$  जिस अन्तराल में है, वह है :
- (1)  $(-\infty, 20]$                       (2)  $[-3, 11]$                       (3)  $(-\infty, 11]$                       (4)  $[-6, 20]$

**Sol. 1**

Lets use LMVT for  $x \in [-7, -1]$

$$\frac{f(-1) - f(-7)}{(-1 + 7)} \leq 2$$

$$\frac{f(-1) + 3}{6} \leq 2 \Rightarrow f(-1) \leq 9$$

Also use LMVT for  $x \in [-7, 0]$

$$\frac{f(0) - f(-7)}{(0 + 7)} \leq 2$$

$$\frac{f(0) + 3}{7} \leq 2 \Rightarrow f(0) \leq 11$$

$$\therefore f(0) + f(-1) \leq 20$$

- 12.** The greatest positive integer  $k$ , for which  $49^k + 1$  is a factor of the sum  $49^{125} + 49^{124} + \dots + 49^2 + 49 + 1$ , is:  
सबसे बड़ी धन पूर्णांक संख्या  $k$ , जिसके लिए  $49^k + 1$  योगफल  $49^{125} + 49^{124} + \dots + 49^2 + 49 + 1$ , का एक गुणखंड है, है :

- (1) 32                      (2) 63                      (3) 60                      (4) 65

**Sol. 2**

$$\frac{(49)^{126} - 1}{48} = \frac{((49)^{63} + 1)(49^{63} - 1)}{48}$$

- 13.** If  $g(x) = x^2 + x - 1$  and  $(g \circ f)(x) = 4x^2 - 10x + 5$ , then  $f\left(\frac{5}{4}\right)$  is equal to:

यदि  $g(x) = x^2 + x - 1$  तथा  $(g \circ f)(x) = 4x^2 - 10x + 5$ , तो  $f\left(\frac{5}{4}\right)$  बराबर है :

- (1)  $-\frac{1}{2}$                       (2)  $\frac{3}{2}$                       (3)  $-\frac{3}{2}$                       (4)  $\frac{1}{2}$

**Sol. 1**

$$g(f(x)) = f^2(x) + f(x) - 1$$

$$g\left(f\left(\frac{5}{4}\right)\right) = 4\left(\frac{5}{4}\right)^2 - 10 \cdot \frac{5}{4} + 5 = \frac{-5}{4}$$

$$g\left(f\left(\frac{5}{4}\right)\right) = f^2\left(\frac{5}{4}\right) + f\left(\frac{5}{4}\right) - 1$$

**Increase Your Score  
for JEE Main April'2020**

**उन्नति**  
17 JAN 2020

Below 97 percentile in JEE Main (Jan-2020)

Tenure: 62 Days | Schedule: 5 Classes Per Day

Fees - ₹ 27500 Including GST

**उत्कर्ष**  
15 JAN 2020

percentile between 97.0 to 98.99  
in JEE Main (Jan-2020)

Fees - ₹ 22000 Including GST

**उत्थान**  
17 JAN 2020

99 percentile and above  
in JEE Main (Jan-2020)

Fees - ₹ 11000 score 160-200 | Fees - ₹ 5500 score 200-240 | Fees - ₹ 0 score above 240

$$-\frac{5}{4} = f^2\left(\frac{5}{4}\right) + f\left(\frac{5}{4}\right) - 1$$

$$f^2\left(\frac{5}{4}\right) + f\left(\frac{5}{4}\right) + \frac{1}{4} = 0$$

$$\left(f\left(\frac{5}{4}\right) + \frac{1}{2}\right)^2 = 0$$

$$f\left(\frac{5}{4}\right) = -\frac{1}{2}$$

- 14.** Let P be a plane passing through the points (2, 1, 0), (4, 1, 1) and (5, 0, 1) and R be any point (2, 1, 6). Then the image of R in the plane P is:

यदि एक समतल P तीन बिन्दुओं (2, 1, 0), (4, 1, 1) और (5, 0, 1) से होकर जाता है, तथा कोई और बिन्दु R (2, 1, 6) है, तो समतल P में R का प्रतिबिम्ब (image) है :

- (1) (6, 5, -2)      (2) (4, 3, 2)      (3) (6, 5, 2)      (4) (3, 4, -2)

**Sol. 1**

$$\text{Plane is } x + y - 2z = 3 \Rightarrow \frac{x-2}{1} = \frac{y-1}{1} = \frac{z-6}{-2} = \frac{-2(2+1-12-3)}{6}$$

$$\Rightarrow (x, y, z) = (6, 5, -2)$$

- 15.** If  $f(a + b + 1 - x) = f(x)$ , for all x, where a and b are fixed positive real numbers, then

$$\frac{1}{a+b} \int_a^b x(f(x) + f(x+1))dx \text{ is equal to:}$$

यदि सभी x के लिए,  $f(a + b + 1 - x) = f(x)$  है, जबकि a तथा b स्थिर (fixed) धन वास्तविक संख्याएँ हैं, तो

$$\frac{1}{a+b} \int_a^b x(f(x) + f(x+1))dx \text{ बराबर है :}$$

- (1)  $\int_{a-1}^{b-1} f(x)dx$       (2)  $\int_{a+1}^{b+1} f(x)dx$       (3)  $\int_{a-1}^{b-1} f(x+1)dx$       (4)  $\int_{a+1}^{b+1} f(x+1)dx$

**Sol. 3**

$$I = \frac{1}{(a+b)} \int_a^b x[f(x) + f(x+1)]dx \quad \dots(i)$$

$$x \rightarrow a + b - x$$

$$I = \frac{1}{(a+b)} \int_a^b (a+b-x)[f(a+b-x) + f(a+b+1-x)]dx$$

$$I = \frac{1}{(a+b)} \int_a^b (a+b-x)[f(x+1) + f(x)]dx \quad \dots(ii)$$

[∴ put  $x \rightarrow x + 1$  in given equation]

(i) + (ii)

$$2I = \int_a^b [f(x+1) + f(x)] dx$$

$$2I = \int_a^b f(x+1) dx + \int_a^b f(x) dx$$

$$\int_a^b f(a+b+1-x) dx + \int_a^b f(x) dx$$

$$2I = 2 \int_a^b f(x) dx$$

16. If the system of linear equations

$$2x + 2ay + az = 0$$

$$2x + 3by + bz = 0$$

$$2x + 4cy + cz = 0,$$

where  $a, b, c \in \mathbb{R}$  are non-zero distinct; has a non-zero solution, then:

(1)  $\frac{1}{a}, \frac{1}{b}, \frac{1}{c}$  are in A.P.

(2)  $a + b + c = 0$

(3)  $a, b, c$  are in A.P.

(4)  $a, b, c$  are in G.P.

यदि निम्न रैखिक समीकरण निकाय

$$2x + 2ay + az = 0$$

$$2x + 3by + bz = 0$$

$$2x + 4cy + cz = 0,$$

जहाँ  $a, b$ , तथा  $c$  विभिन्न शून्येतर वास्तविक संख्याएँ हैं का एक शून्येतर हल है, तो :-

(1)  $\frac{1}{a}, \frac{1}{b}, \frac{1}{c}$  समान्तर श्रेणी में है।

(2)  $a + b + c = 0$

(3)  $a, b, c$  समान्तर श्रेणी में है।

(4)  $a, b, c$  गुणोत्तर श्रेणी में है।

**Sol. 1**

For non - trivial solution

$$\begin{vmatrix} 2 & 2a & a \\ 2 & 3b & b \\ 2 & 4c & c \end{vmatrix} = 0$$

$$\begin{vmatrix} 1 & 2a & a \\ 1 & 3b & b \\ 1 & 4c & c \end{vmatrix} = 0$$

$$(3bc - 4bc) - (2ac - 4ac) + (2ab - 3ab) = 0$$

$$-bc + 2ac - ab = 0$$

$a, b, c$  in H.P.

$$\Rightarrow \frac{1}{a}, \frac{1}{b}, \frac{1}{c} \text{ are in A.P.}$$

**Increase Your Score  
for JEE Main April'2020**

**उत्कर्ष**  
15 JAN 2020

percentile between 97.0 to 98.99  
in JEE Main (Jan-2020)

Fees - ₹ 22000 Including GST

**उन्नति**  
17 JAN 2020

Below 97 percentile in JEE Main (Jan-2020)

Tenure: 62 Days | Schedule: 5 Classes Per Day

Fees - ₹ 27500 Including GST

**उत्थान**  
17 JAN 2020

99 percentile and above  
in JEE Main (Jan-2020)

Fees - ₹ 11000  
score 160-200

Fees - ₹ 5500  
score 200-240

Fees - ₹ 0  
score above 240

17. If  $y(\alpha) = \sqrt{2\left(\frac{\tan \alpha + \cot \alpha}{1 + \tan^2 \alpha}\right) + \frac{1}{\sin^2 \alpha}}$ ,  $\alpha \in \left(\frac{3\pi}{4}, \pi\right)$ , then  $\frac{dy}{d\alpha}$  at  $\alpha = \frac{5\pi}{6}$  is:

यदि  $y(\alpha) = \sqrt{2\left(\frac{\tan \alpha + \cot \alpha}{1 + \tan^2 \alpha}\right) + \frac{1}{\sin^2 \alpha}}$ ,  $\alpha \in \left(\frac{3\pi}{4}, \pi\right)$ , है, तो  $\alpha = \frac{5\pi}{6}$  पर  $\frac{dy}{d\alpha}$  का मान है :

- (1)  $\frac{4}{3}$                       (2) -4                      (3) 4                      (4)  $-\frac{1}{4}$

Sol. 3

$$y = \sqrt{\frac{2 \cos^2 \alpha}{\sin \alpha \cos \alpha} + \frac{1}{\sin^2 \alpha}} = \sqrt{2 \cot \alpha + \operatorname{cosec}^2 \alpha} = |1 + \cot \alpha| = -1 - \cot \alpha$$

$$\frac{dy}{d\alpha} = \operatorname{cosec}^2 \alpha \Rightarrow \left(\frac{dy}{d\alpha}\right)_{\alpha = \frac{5\pi}{6}} \text{ will be } = 4$$

18. If  $y = y(x)$  is the solution of the differential equation,  $e^y \left(\frac{dy}{dx} - 1\right) = e^x$  such that  $y(0) = 0$ , then  $y(1)$  is equal to:

यदि अवकलन समीकरण,  $e^y \left(\frac{dy}{dx} - 1\right) = e^x$ , जबकि  $y(0) = 0$ , का हल  $y=y(x)$  है, तो  $y(1)$  बराबर है:

- (1)  $1 + \log_e 2$                       (2)  $\log_e 2$                       (3)  $2 + \log_e 2$                       (4)  $2e$

Sol. 1

$$e^y = t$$

$$e^y \frac{dy}{dx} = \frac{dt}{dx}$$

$$\frac{dt}{dx} - t = e^x$$

$$\text{IF} = e^{\int -1 dx} = e^{-x}$$

$$t(e^{-x}) = \int e^x \cdot e^{-x} dx$$

$$e^{y-x} = x + C$$

$$\text{Put } x = 0, y = 0 \text{ then } C = 1$$

$$e^{y-x} = x + 1$$

$$y = x - \log(x + 1)$$

$$\text{at } x = 1, y = 1 + \log_e(2)$$

19. The logical statement  $(p \Rightarrow q) \wedge (q \Rightarrow \sim p)$  is equivalent to:

तर्कसंगत कथन  $(p \Rightarrow q) \wedge (q \Rightarrow \sim p)$  निम्न कथनों में से किसके तुल्य है ?

- (1)  $p$                       (2)  $q$                       (3)  $\sim p$                       (4)  $\sim q$

Sol. 3

**24000+**  
SELECTIONS SINCE 2007

JEE (Advanced)	JEE (Main)	NEET / AIIMS	NTSE / OLYMPIADS
<b>5392</b>	<b>16241</b>	<b>1305</b>	<b>1158</b>
(Under 50000 Rank)	(since 2016)	(5th to 10th class)	

H.O. : 394, Rajeev Gandhi Nagar, Kota  
Toll Free : 1800-212-1799  
www.motion.ac.in | info@motion.ac.in



p	q	$p \rightarrow q$	$\sim p$	$q \rightarrow \sim p$	$(p \rightarrow q) \wedge (p \rightarrow \sim q)$
T	T	T	F	F	F
T	F	F	F	T	F
F	T	T	T	T	T
F	F	T	T	T	T

Clearly  $(p \rightarrow q) \wedge (q \rightarrow \sim p)$  is equivalent to  $\sim p$

20. Five numbers are in A.P. whose sum is 25 and product is 2520. If one of these five numbers is  $-\frac{1}{2}$ , then the greatest number amongst them is:

पाँच संख्याएँ समान्तर श्रेणी में हैं, जिनका योगफल 25 तथा गुणनफल 2520 है। यदि इन पाँच संख्याओं में से एक  $-\frac{1}{2}$  है, तो इनमें सबसे बड़ी संख्या है

- (1) 16                      (2) 7                      (3)  $\frac{21}{2}$                       (4) 27

Sol. 1

Let terms be  $a - 2d, a, a - d, a + d, a + 2d$   
 sum = 25  $\Rightarrow 5a = 25 \Rightarrow a = 5$   
 product = 2520  
 $(5 - 2d)(5 - d) + 5(5 + d)(5 + 2d) = 2520$   
 $\Rightarrow (25 - 4d^2)(25 - d^2) = 504$   
 $\Rightarrow 625 - 100d^2 - 25d^2 + 4d^4 = 504$   
 $\Rightarrow 4d^4 - 125d^2 + 121 = 0$   
 $\Rightarrow 4d^4 - 121d^2 - 4d^2 + 121 = 0$   
 $\Rightarrow (d^2 - 1)(4d^2 - 121) = 0$

$$\Rightarrow d = \pm 1, \quad d = \pm \frac{11}{2}$$

$d = \pm 1,$  does not give  $-\frac{1}{2}$  as a term

$$\therefore d = \frac{11}{2}$$

$\therefore$  largest term =  $5 + 2d = 5 + 11 = 16$

21. If the sum of the coefficients of all even powers of  $x$  in the product  $(1 + x + x^2 + \dots + x^{2n})(1 - x + x^2 - x^3 + \dots + x^{2n})$  is 61, then  $n$  is equal to \_\_\_\_\_.

यदि गुणनफल  $(1 + x + x^2 + \dots + x^{2n})(1 - x + x^2 - x^3 + \dots + x^{2n})$  में,  $x$  के सभी समघातों वाले गुणांकों का योगफल 61 है, तो  $n$  बराबर है \_\_\_\_\_.

Sol. 30

Let  $(1 - x + x^2 + \dots)(1 + x + x^2 + \dots) = a_0 + a_1x + a_2x^2 + \dots$

Put  $x = 1$

$$1(2n + 1) = a_0 + a_1 + a_2 + \dots + a_{2n} \quad \dots(i)$$

put  $x = -1$

$$(2n + 1) \times 1 = a_0 - a_1 + a_2 + \dots + a_{2n} \quad \dots(ii)$$

Form (i) and (ii)

$$4n + 2 = 2(a_0 + a_2 + \dots)$$

$$= 2 \times 61$$

$$\Rightarrow 2n + 1 = 61 \Rightarrow n = 30$$

**Increase Your Score  
for JEE Main April'2020**

**उन्नति**  
17 JAN 2020

Below 97 percentile in JEE Main (Jan-2020)

Tenure: 62 Days | Schedule: 5 Classes Per Day

Fees - ₹ 27500 Including GST

**उत्कर्ष**  
15 JAN 2020

percentile between 97.0 to 98.99  
in JEE Main (Jan-2020)

Fees - ₹ 22000 Including GST

**उत्थान**  
17 JAN 2020

99 percentile and above  
in JEE Main (Jan-2020)

Fees - ₹ 11000 | Fees - ₹ 5500 | Fees - ₹ 0  
score 160-200 | score 200-240 | score above 240

**22.** Let S be the set of points where the function,  $f(x) = |2-|x-3||$ ,  $x \in \mathbb{R}$  is not differentiable. Then  $\sum_{x \in S} f(f(x))$  is equal to\_\_\_\_\_.

यदि S उन सभी बिन्दुओं का समुच्चय है, जिनके लिए फलन,  $f(x) = |2-|x-3||$ ,  $x \in \mathbb{R}$  अवकलनीय नहीं है, तो  $\sum_{x \in S} f(f(x))$  बराबर है\_\_\_\_\_.

**Sol. 3**  
 $\therefore f(x)$  is non differentiable at  $x = 1, 3, 5$   
 $\sum f(f(x)) = f(f(1)) + f(f(3)) + f(f(5))$   
 $= 1 + 1 + 1$   
 $= 3$

**23.** Let A(1, 0), B(6, 2) and  $C\left(\frac{3}{2}, 6\right)$  be the vertices of a triangle ABC. If P is a Point inside the triangle ABC such that the triangles APC, APB and BPC have equal areas, then the length of the line segment PQ, where Q is the point  $\left(-\frac{7}{6}, -\frac{1}{3}\right)$ , is\_\_\_\_\_.

माना A(1, 0), B(6, 2) तथा  $C\left(\frac{3}{2}, 6\right)$ , एक त्रिभुज ABC के शीर्ष बिन्दु है। यदि एक बिन्दु P,  $\Delta ABC$  के अन्दर इस प्रकार

है, कि त्रिभुजों APC, APB और BPC के क्षेत्रफल बराबर हैं, तो रेखाखण्ड PQ, जबकि बिन्दु  $Q\left(-\frac{7}{6}, -\frac{1}{3}\right)$  है, की लम्बाई है\_\_\_\_\_.

**Sol. 5**  
P will be centroid of  $\Delta ABC$   
 $P = \left(\frac{17}{6}, \frac{8}{3}\right) \Rightarrow PQ = \sqrt{\left(\frac{24}{6}\right)^2 + 3^2}$   
 $= 5$

**24.** If the variance of the first n natural numbers is 10 and the variance of the first m even natural numbers is 16, then m + n is equal to\_\_\_\_\_.

यदि प्रथम n प्राकृत संख्याओं का प्रसरण 10 है और प्रथम m सम-प्राकृत संख्याओं का प्रसरण 16 है, तो m+n बराबर है\_\_\_\_\_.

**Sol. 18**  
 $\text{var}(1, 2, \dots, n) = 10 \Rightarrow \frac{1^2+2^2+\dots+n^2}{n} - \left(\frac{1+2+\dots+n}{n}\right)^2 = 10$   
 $\Rightarrow \frac{(n+1)(2n+1)}{6} - \left(\frac{n+1}{2}\right)^2 = 10$   
 $\Rightarrow n^2 - 1 = 120 \Rightarrow n = 11$   
 $\text{var}(2, 4, 6, \dots, 2m) = 16 \Rightarrow \text{var}(1, 2, \dots, m) = 4$   
 $\Rightarrow m^2 - 1 = 48 \Rightarrow m = 7 \Rightarrow m + n = 18$

25.  $\lim_{x \rightarrow 2} \frac{3^x + 3^{3-x} - 12}{3^{-x/2} - 3^{1-x}}$  is equal to \_\_\_\_\_.

$\lim_{x \rightarrow 2} \frac{3^x + 3^{3-x} - 12}{3^{-x/2} - 3^{1-x}}$  बराबर है \_\_\_\_\_.

Sol. **72**

Put  $3^{\frac{x}{2}} = t$

$$\Rightarrow \lim_{x \rightarrow 3} \frac{\frac{4t^2}{3} - 12}{-\frac{3}{t^2} + \frac{1}{t}} = \lim_{x \rightarrow 3} \frac{4(t^2 - 9)t^2}{3(-3 + t)} = \lim_{x \rightarrow 3} \frac{4t^2(3 + t)}{3} = \frac{4 \times 9 \times 6}{3} = 72$$

**Increase Your Score  
for JEE Main April'2020**

**उत्कर्ष**  
15 JAN 2020

percentile between 97.0 to 98.99  
in JEE Main (Jan-2020)

Fees - ₹ 22000 Including GST

**उन्नति**  
17 JAN 2020

Below 97 percentile in JEE Main (Jan-2020)

Tenure: 62 Days | Schedule: 5 Classes Per Day

Fees - ₹ 27500 Including GST

**उत्थान**  
17 JAN 2020

99 percentile and above  
in JEE Main (Jan-2020)

Fees - ₹ 11000  
score 160-200

Fees - ₹ 5500  
score 200-240

Fees - ₹ 0  
score above 240

कर लो अब पूरी तैयारी

चूक ना जाये इस बारी

## INCREASE YOUR SCORE for JEE Main April 2020

**उत्थान** 17<sup>th</sup> JAN 2020

99 percentile and above  
in JEE Main (Jan-2020)

**उत्कर्ष** 15<sup>th</sup> JAN 2020

percentile between 97.0 to 98.99  
in JEE Main (Jan-2020)

**उन्नति** 17<sup>th</sup> JAN 2020

Below 97 percentile  
in JEE Main (Jan-2020)

**MOTION**<sup>TM</sup>

Nurturing potential through education

Toll Free : 1800-212-1799