

JEE | NEET | Foundation



29900+ SELECTIONS SINCE 2007

हो चुकी है ऑफलाइन क्लासरूम की शुरूआत अपने सपने को करो साकार, कोटा कोचिंग के साथ



Nitin Vijay (NV Sir) Managing Director Exp. : 18 yrs

Directors of Nucleus Education & Wizard of Mathematics

Now Offline associated with Motion Kota Classroom



Akhilesh Kanther (AKK Sir) Exp. : 17 yrs

Vishal Joshi (VJ Sir) Exp. : 18 yrs

Surendra K. Mishra (SKM Sir) Exp.: 16 yrs



Gavesh Bhardwaj (GB Sir) Exp. : 17 yrs

Academic Pillars of JEE Motion Kota



Ram Ratan Dwivedi (RRD Sir) Joint Director Exp.: 20 yrs



Amit Verma (AV Sir) Joint Director Exp.: 16 yrs



Vijay Pratap Singh (VPS Sir) Vice President Exp.: 20 yrs



Nikhil Srivastava (NS Sir) Head JEE Academics Exp. : 17 yrs



Aatish Agarwal (AA Sir) Sr. Faculty Exp. : 17 yrs



Jayant Chittora (JC Sir) Sr. Faculty Exp.: 16 yrs



Anurag Garg (AG Sir) Sr. Faculty Exp.: 17 yrs

Online + Offline Mode



Arjun Gupta (Árjun Sir) Sr. Faculty Exp.: 14 yrs



(DN Sir) Sr. Faculty Exp.: 13 yrs



Devki Nandan Pathak Avinash Kishore (AVN Sir) Sr. Faculty Exp.: 9 yrs



Vipin Sharma (VS Sir) Sr. Faculty Exp.: 12 yrs



Durgesh Pandey (Pandey Sir) Sr. Faculty Exp.: 8 yrs

Join English & Hindi Medium **EE DROPPER BATC**

Batch Starting from: 22nd Sept. 2021

SECTION - A

- Q.1 Which one of the following complexes is violet in colour :
 - (1) [Fe(CN)₅ NOS]⁴⁻

(2) $Fe_4[Fe(CN)_6]_3 \cdot H_2O$

(3) [Fe(SCN)₆]⁴⁻

(4) [Fe(CN)₆]⁴⁻

Sol. 1

- (1) $[Fe(CN)_6]^{4-} \rightarrow Pale yellow solution$
- (2) $[Fe(SCN)^6]^{4-} \rightarrow Blood red colour$
- (3) $Fe_4[Fe(CN_6)]_3.H_2O \rightarrow Prussian blue$
- (4) $[Fe(CN)_5NOS]^{4-} \rightarrow Violet colour$
- Q.2 The major product formed in the following reaction is :

COOH
$$NH_{2} \longrightarrow NH_{2} \longrightarrow Major product$$
H

(1)
$$CI$$
 N
 NH_2
 NH_2

$$(3) \qquad \begin{array}{c} CO_2CH_3 \\ NH_2 \cdot HCI \\ \end{array}$$

$$(4) \qquad \begin{matrix} CO_2CH_3 \\ NH_2 \cdot HCI \end{matrix}$$

COCI

Sol. 4




An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



- Q.3 The correct sequential addition of regents in the preparation of 3-nitrobenzoic acid from benzene is:
 - (1) Br₂/AlBr₃, NaCN, H₃O⁺, HNO₃/H₂SO₄
 - (2) HNO₃/H₂SO₄,Br₂/AlBr₃,Mg/ether,CO₂, H₃O⁺
 - (3) Br₂/AlBr₃,HNO₃/H₂SO₄,Mg/ether,CO₂, H3O⁺
 - (4) Br₂/AlBr₃,HNO₃/H₂SO₄,NaCN,H₃O⁺

Sol. 2



3- nitrobenzoic acid

Q.4 Given below are two statements:

Statements I: Frenkel defects are vacancy as well as interstitial defects.

Statements II: Frenkel defect leads of colour in ionic solids due to presence of F-centers.

Choose the most appropriate answer for the statements from the options given below:

- (1) Statement I is false but statements II is true.
- (2) Both Statement I and statements II are true.
- (3) Both Statement I and statements II are false.
- (4) Statement I is true but statements II is false.
- Sol.

Theory based.

Q.5 Excess of isobutane on reactions with Br₂ in presence of light at 125°C gives which one of the following, as the major product?

$$CH_3-CH-CH_2-Br$$
(2) CH_3

$$CH_3-CH-CH_2-Br$$

(3) CH_2-Br



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



MOTION JEE MAIN 2021

Sol. 1

$$\begin{array}{ccc} CH_3 & Br_2 & CH_3 \\ CH_3-CH-CH_3 & \longrightarrow & CH_3-C-CH_3 \\ excess isobutane & hv/\Delta & Br \end{array}$$

Q.6 Given below are two statements.

Statements I : The choice of reducing agents for metals extraction can be made by using Ellingham diagram, a plot of ΔG vs temperature .

Statements II : The value of ΔS increases from left to right in Ellingham diagram.

In the light of above statements, choose the most appropriate from the options given below:

- (1) Both Statement I and statements II are false.
- (2) Both Statement I and statements II are true.
- (3) Statement I is false but statements II is true.
- (4) Statement I is true but statements II is false.

Sol. 4

Given statement-I is true as in a number of processes, one element is used to reduce the oxide of another metal. Any element will reduce the oxide of other metal which lie above it in the Ellingham diagram because the free energy change will become more negative. Given statement-II is false as the value of ΔS is decreases from left to right in Ellingham diagram.

Q.7 What are the products formed in sequence when excess of CO₂ is passed in slaked lime?

(1) CaCO₃, Ca(HCO₃)₂

(2) $Ca(HCO_3)_2$, $CaCO_3$

(3) CaO, CaCO₃

(4) CaO, Ca $(HCO_3)_2$

Sol. 1

$$Ca(OH)_2 + CO_2 \longrightarrow CaCO_3 \downarrow +H_2O$$

 $CaCO_3 \downarrow +CO_2 + H_2O \rightarrow Ca(HCO_3)_2$

Q.8 The polymer formed on heating Novolac with formaldehyde is:

(1) Polyester

(2) Melamine

(3) Nylon 6, 6

(4) Bakelite

Sol. 4

Novolac + formaldehyde → Bakelite

Q.9 The incorrect statements is:

- (1) F_2 is stronger oxidizing agent than Cl_2 in aqueous solution.
- (2) Cl₂ is more reactive than CIF.
- (3) On hydrolysis CIF forms HOCl and HF.
- (4) F_2 is more reactive than CIF.

Sol. 2

(i) Reactivity order:

 $F_2 > CIF$ (inter halogen) $> Cl_2$

- (ii) CIF + $H_2O \rightarrow HOCI + HF$
- (iii) Oxidizing power in aqueous solution

 $F_2 > Cl_2 > Br_2 > I_2$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



MOTION JEE MAIN 2021

Q.10 The major product formed in the following reactions is :

$$(1) \xrightarrow{\text{HBr}} \text{Major Product}$$

$$(2) \xrightarrow{\text{Br}}$$

$$(3) \xrightarrow{\text{Br}}$$

$$(4) \xrightarrow{\text{Br}}$$

Sol. 1

- Q.11 Which one of the following, methods is most suitable for preparing deionized water?
 - (1) Calgon's method

(2) Clark's method

(3) Synthetic resin method

(4) Permutit method

Sol. 3

Pure demineralised (de-ionized) water free from all soluble mineral salts is obtained by passing water successively through a cation exchange (in the H^+ form) and an anion exchange (in the OH^- form) resins.

- Q.12 Which one of the following when dissolved in water gives coloured solution in nitrogen atmosphere?
 - (1) AgCl
- (2) CuCl₂
- (3) Cu₂Cl₂
- (4) ZnCl₂

Sol. 1

(1)
$$CuCl_2 + nH_2O \rightarrow Cu^{+2}_{(aq.)}$$

blue colour

- (2) AgCl + $nH_2O \rightarrow Insoluble$
- (3) $ZnCl_2 + nH_2O \rightarrow Zn_{(aq.)}^{+2}$

Colourless

- (4) $Cu_2Cl_2 + nH_2O \rightarrow Insoluble$
- Q.13 Among the following compounds I-IV, which one forms a yellow precipitate on reacting sequentially with (i) NaOH (ii) dil. HNO₃ (iii) AgNO₃?

An Unmatched Experience of Offline



KOTA CLASSROOM For JEE



MOTION™ JEE MAIN 2021

2 Sol.

Other compounds halide can't be removed because corresponding C⁺ is highly unstable.

Given below are two statements: 0.14


Statements I: In the titration between strong acid and weak base methyl orange is suitable as an indicator.

Statement II: For titration of acetic acid with NaOH phenolphthalein is not a suitable indicator. In the light of the above statements, choose the most appropriate answer from the options given below:


- (1) Statement I is true but statements II is false.
- (2) Both Statement I and statements II are false.
- (3) Statement I is false but statements II is true.
- (4) Both Statement I and statements II are true.

Sol.

Titration curve for strong acid and weak base initially a buffer of weak base and conjugate acid is:



Formed, thus pH falls slowly and after equivalence point, so the pH falls sharply so methyl arrange, having pH range of 3.2 to 4.4 will weak as indicator. So statement-I is correct.



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE





Titration curve for weak acid and strong base (NaOH) Initially weak acid will form a buffer so pH increases slowly but after equivalence point. It rises sharply covering range of phenolphthalein so it will be suitable indicator so statement-II is false.

- Q.15 Which one of the following is correct for the adsorption of a gas at a given temperature on a solid surface?
 - (1) $\Delta H < 0$, $\Delta S > 0$

(2)
$$\Delta H > 0$$
, $\Delta S < 0$

(3)
$$\Delta H < 0$$
, $\Delta S < 0$

(4)
$$\Delta H > 0$$
, $\Delta S > 0$

Sol.

- (i) Adsorption of gas at metal surface is an exothermic process so $\Delta H < 0$
- (ii) As the adsorption of gas on metal surface reduces the free movement of gas molecules thus restricting its randomness hence $\Delta S < 0$
- Q.16 Given below are two statements:

Statements I: The limiting molar conductivity of KCI (strong electrolyte) is higher compared to that of CH₃COOH (weak electrolyte)

Statement II: Molar conductivity decreases with decrease in concentration of electrolyte. In the light of the above statements, choose the most appropriate answer from the options given below:

- (1) Statement I is false but statements II is true.
- (2) Both Statement I and statements II are true.
- (3) Both Statement I and statements II are false.
- (4) Statement I is true but statements II is false.
- Sol.

Ion	H⁺	K ⁺	Cl⁻	CH₃COO⁻
$\Lambda_{mScm^2/mole}^{\infty}$	349.8	73.5	76.3	40.9

So,
$$\Lambda_{\text{mCH}_3\text{COOH}}^{\infty} = \Lambda_{\text{m(H}^+)}^{\infty} + \Lambda_{\text{mCH}_3\text{COO}^-}^{\infty}$$

$$= 349.8 + 40.9$$


$$= 390.7 \text{ Scm}^2/\text{mole}$$

$$\Lambda_{\mathsf{mKCI}}^{\infty} = \Lambda_{\mathsf{m}(\mathsf{K}^{\scriptscriptstyle{+}})}^{\infty} + \Lambda_{\mathsf{m}(\mathsf{CI}^{\scriptscriptstyle{-}})}^{\infty}$$

$$= 73.5 + 76.3 = 149.3 \text{ Scm}^2/\text{mole}$$

So statement-I is wrong or False.

As the concentration decreases, the dilution increases which increases the degree of dissociation, thus increasing the no. of ions, which increases the molar conductance. So statement-II is false.



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



Q.17 Given below are two statements:

Statements I: According to Bohr's model of an atom, qualitatively the megnitude of velocity of electron increases with decrease in positive charges on the nucleus as there is no strong hold on the electron by the nucleus.

Statement II: According to Bohr's model of an atom, qualitatively the megnitude of velocity of electron increases with decrease in principal quantum number.

In the light of the above statements, choose the most appropriate answer from the options given below:

- (1) Both Statement I and statements II are false.
- (2) Both Statement I and statements II are true.
- (3) Statement I is true but statements II is false.
- (4) Statement I is false but statements II is true.

Sol.

Velocity of electron in Bohr's atom is given by

$$V \propto \frac{Z}{n}$$

Z = atomic number of atom, corresponds to +ve charge so as Z increase velocity increases so statement-I is wrong.

and as 'n' decreases velocity increases so statement-II is correct.

The correct options for the products A and B of the following reactions are:

$$\mathbf{A} \xleftarrow{\operatorname{Br}_{2}(\operatorname{Excess})}{\operatorname{H}_{2}\operatorname{O}} \underbrace{\bigcirc \frac{\operatorname{Br}_{2}}{\operatorname{CS}_{2}, <5^{\circ}\operatorname{C}}} \mathbf{B}$$

$$\mathbf{A} = \bigcup_{\mathsf{Br}}^{\mathsf{OH}} \mathsf{Br} \quad , \quad \mathbf{B} = \bigcup_$$

Sol. 2

$$A \xrightarrow{Br} \xrightarrow{Br_2} \xrightarrow{Br_2} \xrightarrow{Br_2} \xrightarrow{Br_2} \xrightarrow{Br} \xrightarrow{Br}$$

An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



Q.19 The major products formed in the following reaction sequence A and B are:

$$CH_3 \xrightarrow{Br_2} A+B$$

(1)
$$\mathbf{A} = \begin{bmatrix} \mathbf{Br} & \mathbf{O} & \mathbf{Br} \\ \mathbf{II} & \mathbf{C} - \mathbf{CH}_3 \\ \mathbf{Br} & \mathbf{II} \end{bmatrix}$$

(2)
$$\mathbf{A} = \left\langle \begin{array}{c} 0 \\ \parallel \\ 0 \\ \text{OK} \end{array} \right\rangle$$
, $\mathbf{B} = \text{CHBr}_3$

(3)
$$\mathbf{A} = \left\langle \begin{array}{c} 0 \\ \parallel \\ \text{C-CBr}_3 \end{array} \right\rangle$$
, $\mathbf{B} = \left\langle \begin{array}{c} \\ \end{array} \right\rangle$ -CHO

(4)
$$A = C - CH_2 - Br$$
 , $B = C - CH_2 - OH_2 - O$

Sol. 2

$$C-CH_3 \xrightarrow{Br_2} A + B$$

$$(1) \bigcirc \bigcap_{C \text{ OK}} \bigcap_{C \text{ OK$$

- Q.20 The conversion of hydroxyapatite occure due to presence of F^- ions in water. The correct formula of hydroxyapatite is :
 - (1) $[3Ca_3(PO_4)_2 \cdot CaF_2]$

(2) $[Ca_3(PO_4)_2 \cdot CaF_2]$

(3) $[3Ca(OH)_2 \cdot CaF_2]$

(4) $[3Ca_3(PO_4)_2 \cdot Ca(OH)_2]$

Sol. 4

The F^{Θ} ions make the enamel on teeth much harder by converting hydroxyapatite, $[3(Ca_3(PO_4)_2].Ca(OH)_2]$, the enamel on the surface of the teeth into much harder fluroappatite. $[3Ca_3(PO_4)_2.CaF_2]$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



SECTION-B

- Q.1 Of the following four agueous solutions, total number of those solutions whose freezing points is lower than that of 0.10 M C_2H_5OH is (Integer answer)
 - (1) $0.10 \text{ M Ba}_3(PO_4)_2$

(2) 0.10 M Na₂SO₄

(3) 0.10 M KCI

(4) 0.10 M Li₃PO₄

Sol.

As 0.1 M C₂H₅OH is non-dissociative and rest all salt given are electrolyte so in each case effective molarity > 0.1 so each will have lower freezing point.

The following data was obtained for chemical reaction given below at 975 K. Q.2

$$2NO_{(g)} + 2H_{2_{(g)}} \rightarrow N_{2_{(g)}} + 2H_2O_{(g)}$$

[NO] $Mol L^{-1}$

 $Mol L^{-1}$

 $Mol L^{-1} s^{-1}$

- (1)
- 7×10^{-9}

- (2)
- 8×10^{-5} 8×10^{-5} 24×10^{-5} 8×10^{-5}
- 2.1×10^{-8}

- 24×10^{-5}
- 32×10^{-5}
- 8.4×10^{-8}

The order of the reaction with respect to NO is ______ (Integer answer)

Sol.

$$7 \times 10^{-9} = K \times (8 \times 10^{-5})^{x} (8 \times 10^{-5})^{y} \dots (1)$$

 $2.1 \times 10^{-8} = K \times (24 \times 10^{-5})^{x} (8 \times 10^{-5})^{y} \dots (2)$

$$\frac{1}{3} = \left(\frac{1}{3}\right)^{x} \Rightarrow x = 1$$

The Born-Haber cycle for KCl is evaluated with the following data: Q.3

$$\Delta_{\rm f}\overset{\Theta}{\rm H}$$
 for KCI = -436.7 kJ mol⁻¹; $\Delta_{\rm sub}\overset{\Theta}{\rm H}$ for K = 89.2 kJ mol⁻¹

$$\Delta_{\text{ionization}} \overset{\Theta}{H}$$
 for K = 419.0 kJ mol⁻¹; $\Delta_{\text{electrongain}} \overset{\Theta}{H}$ for Cl_(g) = -348.6 kJ mol⁻¹

$$\Delta_{\text{bond}} \overset{\Theta}{\text{H}}$$
 for Cl₂ = 243.0 kJ mol⁻¹

the magnitude of lattice enthalpy of KCl in kJ mol⁻¹ is _____ (Nearest integer)

Sol.

$$\Delta_{\rm f} H_{\rm K:CI}^{\odot} = \Delta_{\rm sub} H_{\rm (K)}^{\odot} + \Delta_{\rm ionization} H_{\rm (K)}^{\odot} + \frac{1}{2} \Delta_{\rm bond} H_{\rm (Cl_2)}^{\odot}$$

$$+ \Delta_{\rm electron \; gain} H^{\odot}_{\rm (CI)} + \Delta_{\rm lattice} H^{\odot}_{\rm (KCI)}$$

$$\Rightarrow -436.7 = 89.2 + 419.0 + \frac{1}{2} (243.0) + \{-348.6\}$$

$$+\Delta_{\text{lattice}}H^{\odot}_{(\text{KCI})}$$

$$\Rightarrow \Delta_{\text{lattice}} H_{(\text{KCI})}^{\odot} = -717.8 \text{kJ mol}^{-1}$$

The magnitude of lattice enthalpy of KCl in kJ mol⁻¹ is 718 (Nearest integer).



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



ANSWER KEY

Q.4 The number of 4f electrons in the ground state electronic configuration Gd^{2+} is _____ (Atomic number of Gd = 64)

Sol. 7

The electronic configuration of

 $_{64}$ Gd: [Xe] $4f^7 5d^1 6s^2$

So the electronic configuration of

 $_{64}\text{Gd}^{2+}$: [Xe] $4f^7 5d^1 6s^0$

i.e. the number of 4f electrons in the ground state electronic configuration of Gd^{2+} is 7.

Q.5 The ratio of number of water molecules in Mohr's salt and potash alum is $___ \times 10^{-1}$.

Sol. 5

Mohr's salt : $(NH_4)_2$ Fe $(SO_4)_2$.6H₂O

The number of water molecules in Mohr's salt = 6

Potash alum : $KAI(SO_4)_2.12H_2O$

The number of water molecules in potash alum = 12

So ratio of number of water molecules in Mohr's

salt and potash alum =
$$\frac{6}{12}$$

$$=\frac{1}{2}$$

$$=5\times10^{-1}$$

Q.6 These are physical properties of an elements.

- (1) Sublimation enthalpy
- (2) Ionisation enthalpy
- (3) Hydration enthalpy
- (4) Electron gain enthalpy

The total number of above properties that affect the reduction potential is _____ (Integer answer)


Sol. 3

Sublimation enthalpy, ionisation enthalpy and hydration enthalpy affect the reduction potential.

Q.7 AB $_3$ is an interhalogen T-shaped molecule. The number of lone pairs of electrons on A is _______. (Integer answer)

Sol. 2

T-Shaped molecule means 3 sigma bond and 2 lone pairs of electrons on central atom.





An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



MOTION JEE MAIN 2021

ANSWER KEY

Q.8 The total number of negative charge in the tetrapeptide, Gly-Glu-Asp-Tyr at pH 12.5 will be ______ . (Integer answers)

Sol. 4

Total negative charge produced = 4.

Q.9 An aqueous KCl solution of density 1.20 g mL^{-1} has a molality of 3.30 mol kg^{-1} . The molarity of the solution in mol L^{-1} is ______ . (Nearest integer)

Sol. 3

1000 kg solvent has 3.3 moles of KCl

1000 kg solvent \rightarrow 3.3 \times 74.5 gm KCl

$$\rightarrow 245.85$$

weight of solution = 1245.85 gm

volume of solution = $\frac{1245.85}{1.2}$ mI

So molarity = $\frac{3.3 \times 1.2}{1245.85} \times 1000 = 3.17$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



ANSWER KEY

- Q.10 The OH $^-$ concentration in a mixture of 5.0 mL of 0.0504 M NH $_4$ Cl and 2 mL of 0.0210 M NH $_3$ solution is x \times 10 $^{-6}$ M. The value of x is _____. (Nearest integer)
- Sol. 3

$$\left[NH_{4}^{+}\right] = 0.0504 \& \left[NH_{3}\right] = 0.0210$$

So, Kb =
$$\frac{\left[NH_{4}^{+}\right]\left[HO^{-}\right]}{\left[NH_{3}\right]}$$

[HO⁻] =
$$\frac{k_b \times [NH_3]}{[NH_4^+]}$$
 = 1.8 × 10⁻⁵ × $\frac{2}{5}$ × $\frac{210}{504}$

$$= 3 \times 10^{-6}$$

An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



हो चुकी है ऑफलाइन क्लासरूम की शुरूआत अपने सपने को करो साकार, कोटा कोचिंग के साथ



Nitin Vijay (NV Sir) Managing Director Exp. : 18 yrs

Directors of Nucleus Education & Wizard of Mathematics

Now Offline associated with Motion Kota Classroom



Akhilesh Kanther (AKK Sir) Exp. : 17 yrs

Vishal Joshi Surendra K. Mishra (VJ Sir) (SKM Sir) Exp. : 18 yrs Exp.: 16 yrs



Gavesh Bhardwai (GB Sir) Exp. : 17 yrs

Academic Pillars of JEE Motion Kota



Ram Ratan Dwivedi (RRD Sir) Joint Director Exp.: 20 yrs



Amit Verma (AV Sir) Joint Director Exp.: 16 yrs



Vijay Pratap Singh (VPS Sir) Vice President Exp.: 20 yrs



Nikhil Srivastava (NS Sir) Head JEE Academics Exp.: 17 yrs



Aatish Agarwal (AA Sir) Sr. Faculty Exp. : 17 yrs



Jayant Chittora (JC Sir) Sr. Faculty Exp. : 16 yrs



Anurag Garg (AG Sir) Sr. Faculty Exp.: 17 yrs



Arjun Gupta (Arjun Sir) Sr. Faculty Exp.: 14 yrs



Devki Nandan Pathak Avinash Kishore (DN Sir) Sr. Faculty Exp.: 13 yrs



(AVN Sir) Sr. Faculty Exp. : 9 yrs



Vipin Sharma (VS Sir) Sr. Faculty Exp.: 12 yrs



Durgesh Pandey (Pandey Sir) Sr. Faculty Exp. : 8 yrs

Join **English & Hindi Medium** DROPPER BATCH

Online + Offline Mode

Batch Starting from: 22nd Sept. 2021