

**JEE  
MAIN  
4<sup>th</sup>  
Attempt**

**CHEMISTRY**  
**27<sup>th</sup> August 2021 [SHIFT – 2]**  
**QUESTION WITH SOLUTION**

**JEE | NEET | Foundation**

**MOTION<sup>®</sup>**

**29900+**  
SELECTIONS SINCE 2007

हो चुकी है ऑफलाइन क्लासरूम की शुरुआत  
अपने सपने को करो साकार, कोटा कोचिंग के साथ

## Directors of Nucleus Education & Wizard of Mathematics

Now Offline associated with Motion Kota Classroom



**Nitin Vijay (NV Sir)**  
Managing Director  
Exp. : 18 yrs



**Akhilesh Kanther (AKK Sir)**  
Exp. : 17 yrs



**Vishal Joshi (VJ Sir)**  
Exp. : 18 yrs



**Surendra K. Mishra (SKM Sir)**  
Exp. : 16 yrs



**Gavesh Bhardwaj (GB Sir)**  
Exp. : 17 yrs

## Academic Pillars of JEE MOTION KOTA



**Ram Ratan Dwivedi (RRD Sir)**  
Joint Director  
Exp. : 20 yrs



**Amit Verma (AV Sir)**  
Joint Director  
Exp. : 16 yrs



**Vijay Pratap Singh (VPS Sir)**  
Vice President  
Exp. : 20 yrs



**Nikhil Srivastava (NS Sir)**  
Head JEE Academics  
Exp. : 17 yrs



**Aatish Agarwal (AA Sir)**  
Sr. Faculty  
Exp. : 17 yrs



**Jayant Chittora (JC Sir)**  
Sr. Faculty  
Exp. : 16 yrs



**Anurag Garg (AG Sir)**  
Sr. Faculty  
Exp. : 17 yrs



**Arjun Gupta (Arjun Sir)**  
Sr. Faculty  
Exp. : 14 yrs



**Devki Nandan Pathak (DN Sir)**  
Sr. Faculty  
Exp. : 13 yrs



**Avinash Kishore (AVN Sir)**  
Sr. Faculty  
Exp. : 9 yrs



**Vipin Sharma (VS Sir)**  
Sr. Faculty  
Exp. : 12 yrs



**Durgesh Pandey (Pandey Sir)**  
Sr. Faculty  
Exp. : 8 yrs

Join

**JEE DROPPER BATCH**

Online + Offline Mode

English & Hindi Medium

Batch Starting from :  
**22nd Sept. 2021**

### SECTION - A

1. Which one of the following tests used for the identification of functional groups in organic compounds does not use copper reagent?

- (1) Seliwanoff's test (2) Barfoed's test  
 (3) Biuret test for peptide bond (4) Benedict test

Sol. (1)

This test used to detect the presence of aldose and ketose reagent : resorcinol and conc hcl it is used to detect monosaccharide by seduction of cu(II) Cu(I).

Used to detect presence of peptide bond protein by seduction on of Cu(II) into Cu(I).

It is used to detect aldehyde and ketone by reduction of Cu(II) into Cu(I)

2. Choose the correct statement from the following :

- (1) Among the alkali metal halides, LiF is least soluble in water  
 (2) The low solubility of CsI in water is due to its high lattice enthalpy  
 (3) The standard enthalpy of formation for alkali metal bromides becomes less negative on descending the groups.  
 (4) LiF has least negative standard enthalpy of formation among alkali metal fluorides.

Sol. (1)






Standard enthalpy of formation for alkali metal bromides becomes more negative on desending down the group.

1. In case of CsI, lattice energy is less, but  $\text{Cs}^+$  is having less hydration enthalpy due to which it is less soluble in water.

2. For alkali metal fluorides, the solubility in water increases from lithium to caesium. LiF is least soluble in water.

2. Standard enthalpy of formation for LiF is most negative among alkali metal fluorides.

3. Which one of the following is the major product of the given reaction?




An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**




Sol. 1



4. In stratosphere most of the ozone formation is assisted by  
 (1) visible radiations (2) ultraviolet radiation  
 (3)  $\gamma$ -rays (4) cosmic rays

Sol. (2)  
 Ozone in the stratosphere is a product of UV radiations acting on dioxygen ( $\text{O}_2$ ) molecules.



5. Potassium permanganate on heating at 513K gives a product which is  
 (1) paramagnetic and green (2) diamagnetic and colourless  
 (3) paramagnetic and colourless (4) diamagnetic and green

Sol. (1)  
 $2\text{KMnO}_4 \xrightarrow[200^\circ\text{C}]{\Delta} \text{K}_2\text{MnO}_4 + \text{MnO}_2 + \text{O}_2$

In  $\text{K}_2\text{MnO}_4$ , manganese oxidation state is +6 and hence it has one unpaired e-.

6. Given below are two statements :

**Statement I :** Ethyl pent-4-yn-oate on reaction with  $\text{CH}_3\text{MgBr}$  gives a 3° alcohol.

**Statement II :** In this reaction one mole of ethyl pent-4-yn-oate utilizes two moles of  $\text{CH}_3\text{MgBr}$ .  
 In the light of the above statements choose the most appropriate answer from the options given below :

- (1) statement I is true but statement II is false  
 (2) both statement I and statement II are true  
 (3) both statement I and statement II are false  
 (4) statement I is false but statement II is true

An Unmatched Experience of Offline




**KOTA CLASSROOM** For JEE



New batch Starting from : **22nd Sept. 2021**



- Sol. (1)  
Statement 1 is true  
But it consume 3 moles of G R  
So statement 2 is false



7. The correct structures of A and B formed in the following reactions are :




An Unmatched Experience of Offline



**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**






- 8.** The oxide that gives  $H_2O_2$  most readily on treatment with  $H_2O$  is :  
 (1)  $Na_2O_2$       (2)  $BaO_2 \cdot 8H_2O$       (3)  $PbO_2$       (4)  $SnO_2$

- Sol.** (1)
- (1)  $PbO_2 + 2H_2O \rightarrow Pb(OH)_4$   
 (2)  $Na_2O_2 + 2H_2O \rightarrow 2NaOH + H_2O_2$   
 This reaction is possible at room temperature  
 3.  $SnO_2 + 2H_2O \rightarrow Sn(OH)_4$   
 4. Acidified  $BaO_2 \cdot 8H_2O$  gives  $H_2O_2$  after evaporation

- 9.** The compound/s which will show significant intermolecular H-bonding is/are :



- Sol.** (2)  
 (a) Shows intra molecular H-bonding  
 (b) Shows significant intermolecular H-bonding  
 (c) It do not show intermolecular H-bonding due to steric hindrance.

- 10.** The correct order of ionic radii for the ions,  $P^{3-}$ ,  $S^{2-}$ ,  $Ca^{2+}$ ,  $K^+$ ,  $Cl^-$  is :  
 (1)  $Cl^- > S^{2-} > P^{3-} > Ca^{2+} > K^+$       (2)  $P^{3-} > S^{2-} > Cl^- > K^+ > Ca^{2+}$   
 (3)  $P^{3-} > S^{2-} > Cl^- > Ca^{2+} > K^+$       (4)  $K^+ > Ca^{2+} > P^{3-} > S^{2-} > Cl^-$



An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



**Sol.** (2)  
 $P^{3-} > S^{2-} > Cl^{-} > K^{+} > Ca^{2+}$

(Correct order of ionic radii)

all the given species are isoelectronic species.





In isoelectronic species size increases with increase of negative charge and size decreases with increase in positive charge.

**11.** Which one of the following is formed (mainly) when red phosphorus is heated in sealed tube at 803K?


- (1) Yellow phosphorus (2) White phosphorus  
 (3)  $\beta$ -black phosphorus (4)  $\alpha$ -black phosphorus

**Sol.** (4)  
 When red phosphorus is heated in a sealed tube at 803 K,  $\alpha$ -black phosphorus is formed.

**Q.12** Which one of the following chemicals is responsible for the production of HCl in the stomach leading to irritation and pain?



**Sol.** (4)  
 Histamine stimulates the secretion of HCl



**Q.13** Lyophilic sols are more stable than lyophobic sols because,

- (1) The colloidal particles have positive charge.  
 (2) The Colloidal particles are solvated.  
 (3) There is a strong electrostatic repulsion between the negatively charged colloidal particles.  
 (4) The colloidal particles have no charge.

**Sol.** (2)  
 In the lyophilic colloids, the colloidal particles are extensively solvated.



An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**




**Q.14** Which one of the following reactions will not yield propionic acid?


- (1)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br} + \text{Mg}, \text{CO}_2$  dry ether/ $\text{H}_3\text{O}^+$
- (2)  $\text{CH}_3\text{CH}_2\text{CCl}_3 + \text{OH}^-/\text{H}_3\text{O}^+$
- (3)  $\text{CH}_3\text{CH}_2\text{COCH}_3 + \text{OI}^-/\text{H}_3\text{O}^+$
- (4)  $\text{CH}_3\text{CH}_2\text{CH}_3 + \text{KMnO}_4(\text{Heat}), \text{OH}^-/\text{H}_3\text{O}^+$





**Sol.** (1)

All gives propanoic acid as product but option 4 gives butanoic as product



**Q.15** The major product of the following reaction, if it occurs by  $\text{S}_{\text{N}}2$  mechanism is :



- (1) 
- (2) 
- (3) 
- (4) 




An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



Sol. (2)




**Q.16** Hydrolysis of sucrose gives :

- (1)  $\alpha$ -D-(+)-Glucose and  $\beta$ -D-(-)-Fructose
- (2)  $\alpha$ -D-(-)-Glucose and  $\alpha$ -D-(+)-Fructose
- (3)  $\alpha$ -D-(+)-Glucose and  $\alpha$ -D-(-)-Fructose
- (4)  $\alpha$ -D-(-)-Glucose and  $\beta$ -D-(-)-Fructose

Sol. (1)

Sucrose is formed by  $\alpha$ -D(+). Glucose +  $\beta$ -D (-)Fructose. we obtain these monomers on hydrolysis.

**Q.17** The major product (A) formed in the reaction given below is :




An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



Sol. (4)



**Q.18** Which one of the following is used to remove most of plutonium from spent nuclear fuel ?

- (1)  $I_2O_5$                       (2)  $ClF_3$                       (3)  $O_2F_2$                       (4)  $BrO_3$

Sol. (3)

$O_2F_2$  oxidises plutonium to  $PuF_6$  and the reaction is used in removing plutonium as  $PuF_6$  from spent nuclear fuel.

**Q.19** Match List- I with List – II:

List-I (Name of ore/mineral)		List-II (Chemical formula)	
(a)	Calamine	(i)	ZnS
(b)	Malachite	(ii)	$FeCO_3$
(c)	Siderite	(iii)	$ZnCO_3$
(d)	Sphalerite	(iv)	$CuCO_3 \cdot Cu(OH)_2$

Choose the most appropriate answer from the option given below

- (1) (a) – (iii), (b) – (ii), (c) – (iv), (d) – (i)  
 (2) (a) – (iv), (b) – (iii), (c) – (i), (d) – (ii)  
 (3) (a) – (iii), (b) – (iv), (c) – (ii), (d) – (i)  
 (4) (a) – (iii), (b) – (iv), (c) – (i), (d) – (ii)

Sol. (3)

(Name of ore/mineral)

- (a) Calamine                       $ZnCO_3$   
 (b) Malachite                       $CuCO_3 \cdot Cu(OH)_2$   
 (c) Siderite                       $FeCO_3$   
 (d) Sphalerite                      ZnS



An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**





If 'n' photons of  $\lambda = 1000 \text{ nm}$  are emitted, then :  $10^{-4} = n \times \frac{hc}{\lambda}$

$$\Rightarrow 10^{-4} = \frac{n \times 6.63 \times 10^{-34} \times 3 \times 10^8}{1000 \times 10^{-9}}$$

$$\Rightarrow n = 5.02 \times 10^{14} - 50.2 \times 10^{13}$$

$\Rightarrow 50$  (nearest integer)

Q.3 The resistance of a conductivity cell with cell constant  $1.14 \text{ cm}^{-1}$ , containing  $0.001 \text{ M KCl}$  at  $298 \text{ K}$  is  $1500 \Omega$ . The molar conductivity of  $0.001 \text{ M KCl}$  solution at  $298 \text{ K}$  is  $S \text{ cm}^2 \text{ mol}^{-1}$  is \_\_\_\_\_. (Integer answer)

Sol. **760**


$$K = \frac{1}{R} \times \ell / A = \left( \left( \frac{1}{1500} \right) \times 1.14 \right) S \text{ cm}^{-1}$$

$$\Rightarrow \wedge_m = 1000 \times \frac{\left( \frac{1.14}{1500} \right)}{0.001} S \text{ cm}^2 \text{ mol}^{-1}$$

$$= 760 S \text{ cm}^2 \text{ mol}^{-1}$$

$\Rightarrow 760$

Q.4 Two flasks I and II shown below are connected by a valve of negligible volume.



When the valve is opened, the final pressure of the system in bar is  $x \times 10^{-2}$ . The value of  $x$  is \_\_\_\_\_ (Integer answer)

[Assume - Ideal gas;  $1 \text{ bar} = 10^5 \text{ Pa}$ ; Molar mass of  $\text{N}_2 = 28.0 \text{ g mol}^{-1}$  ;

$R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$ ]

Sol. **84**

$\Rightarrow$  Assuming the system attains a final temperature of  $T$  (such that  $300 < T < 60$ )

$$\Rightarrow \left( \begin{array}{l} \text{Heat lost by} \\ N_2 \text{ of Container} \\ I \end{array} \right) = \left( \begin{array}{l} \text{Heat gained by} \\ H_2 \text{ of container} \\ II \end{array} \right)$$



An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



$$\Rightarrow n_l C_m (300 - T) = n_{II} C_m (T - 60)$$

$$\Rightarrow \left(\frac{2.8}{28}\right)(300 - T) = \frac{0.2}{28}(T - 60)$$

$$\Rightarrow 14(300 - T) = T - 60$$





$$\Rightarrow \Delta T_r = T_r - T_f' = 1.86 \times \frac{10}{9}$$

$$\Rightarrow T_r' = 273.15 - 1.86 \times \frac{10}{9}$$


$$= 271.08 \text{ K}$$

$$= 271 \text{ K (nearest-integer)}$$


Q.5 The number of species having non-pyramidal shape among the following is \_\_\_\_\_




**Sol.** 3




Trigonal planar



Trigonal planar



Trigonal planar



Pyramidal

Hence non-pyramidal species are  $\text{SO}_3$ ,  $\text{NO}_3^-$  and  $\text{CO}_3^{2-}$ .

Q.6 40 g of glucose (Molar mass = 180) is mixed with 200 mL of water. The freezing point of solution is \_\_\_\_\_ K. (Nearest integer)

[Given :  $K_f = 1.86 \text{ K kg mol}^{-1}$  ; Density of water =  $1.00 \text{ g cm}^{-3}$  ; Freezing point of water =  $273.15 \text{ K}$ ]

**Sol.** 271

$$w_{\text{C}_6\text{H}_{12}\text{O}_6} = 40\text{gm}$$

$$n_{\text{C}_6\text{H}_{12}\text{O}_6} = \frac{40}{180} \text{ mole}$$

An Unmatched Experience of Offline



**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



$$d_{\text{H}_2\text{O}} = 1\text{gm/ml}$$

$$W_{\text{H}_2\text{O}} = 200\text{gm} = 0.2\text{kg}$$

$$\Delta T_f = \text{kg m} = \frac{\text{kg} \times \text{mole}}{W_{\text{H}_2\text{O}} (\text{kg})}$$

$$\Delta T_f = \frac{1.86 \times 40}{180 \times 0.2}$$

$$\Delta T_f = 2.067$$

$$(T_f)_{\text{sol}} = (T_f^\circ) - \Delta T_f$$


$$= 273.15 - 2.067$$

$$(T_f)_{\text{sol}} = 271\text{K}$$

Q.7 The number of optical isomers possible for  $[\text{Cr}(\text{C}_2\text{O}_4)]^{-3}$  is \_\_\_\_\_

**Sol.** 2

The number of optical isomers for  $[\text{Cr}(\text{C}_2\text{O}_4)_3]^{-3}$  is two.



Q.8 When 5.1 g of solid  $\text{NH}_4\text{HS}$  is introduced into a two litre evacuated flask at  $27^\circ\text{C}$ , 20% of the solid decomposes into gaseous ammonia and hydrogen sulphide. The  $K_p$ , for the reaction at  $27^\circ\text{C}$  is  $x \times 10^{-2}$ . The value of  $x$  is \_\_\_\_\_. (Integer answer)

[Given  $R=0.082 \text{ L atm K}^{-1} \text{ mol}^{-1}$ ]

**Sol.** 6

$$n_{\text{NH}_4\text{HS}} = \frac{5.1}{51} = 0.1\text{mole}$$



t = 0	0.1	-	-
eq	0.1 - 0.02	0.02	0.02

$$pV = nRT$$



An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



$$P = \frac{nRT}{V}$$

$$K_p = (P_{\text{NH}_3})(P_{\text{H}_2\text{S}})$$


$$K_p = \left(\frac{0.02 \times RT}{V}\right) \left(\frac{0.02 \times RT}{V}\right)$$

$$K_p = \left(\frac{0.02 \times 0.0821 \times 300}{2}\right)^2$$

$$K_p = 6.07 \times 10^{-2}$$

$$x = 6$$

Q.9 Data given for the following reaction is as follows :



Substance	$\Delta_f H^\circ$ (KJ mol <sup>-1</sup> )	$\Delta S^\circ$ (J mol <sup>-1</sup> K <sup>-1</sup> )
FeO <sub>(s)</sub>	-266.3	57.49
C <sub>(graphite)</sub>	0	5.74
Fe <sub>(s)</sub>	0	27.28
CO <sub>(g)</sub>	-110.5	197.6

The minimum temperature in K at Which the reaction becomes spontaneous is \_\_\_\_ (integer answer)

**Sol.** 964

$$\Delta_r H^\circ = \sum \Delta H^\circ_p - \sum \Delta H^\circ_R$$

$$\Delta_r H^\circ = (0 - 110.5) - (-266.3 + 0)$$

$$\Delta_r H^\circ = 155.8 \text{ kJ/mole}$$

$$\Delta_r S^\circ = \sum \Delta_r S^\circ_p - \sum S^\circ_R$$

$$= (27.28 + 197.6) - (57.49 + 5.74)$$

$$\Delta_r S^\circ = 161.65 \text{ J/mol-k}$$

For spontaneous nature

$$\Delta G^\circ < 0$$

$$\Delta H^\circ - T\Delta S^\circ < 0$$

$$T > \frac{\Delta H^\circ}{\Delta S^\circ}$$



An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



$$T > \frac{155.8}{161.65 \times 10^{-3}}$$

$$T > 963.81$$

Minimum temperature = 964K

Q.10 100 g of propane is completely reacted with 1000 g of oxygen. The mole fraction of carbon dioxide in the resulting mixture is  $x \times 10^{-2}$ . The value of x is \_\_\_\_\_

(Nearest integer)


[Atomic weight: H = 1.008; C = 12.00; O = 16.00]

**Sol.** 19

$$n_{\text{C}_3\text{H}_8} = \frac{100}{44} \text{ mole}$$

$$n_{\text{O}_2} = \frac{1000}{32} \text{ mole}$$

O<sub>2</sub> is L.R.



An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



हो चुकी है ऑफलाइन क्लासरूम की शुरुआत  
अपने सपने को करो साकार, कोटा कोविंग के साथ

**Directors of Nucleus Education & Wizard of Mathematics**

Now Offline associated with Motion Kota Classroom



**Nitin Vijay (NV Sir)**  
Managing Director  
Exp. : 18 yrs



**Akhilesh Kanther (AKK Sir)**  
Exp. : 17 yrs



**Vishal Joshi (VJ Sir)**  
Exp. : 18 yrs



**Surendra K. Mishra (SKM Sir)**  
Exp. : 16 yrs



**Gavesh Bhardwaj (GB Sir)**  
Exp. : 17 yrs

## Academic Pillars of JEE MOTION KOTA



**Ram Ratan Dwivedi (RRD Sir)**  
Joint Director  
Exp. : 20 yrs



**Amit Verma (AV Sir)**  
Joint Director  
Exp. : 16 yrs



**Vijay Pratap Singh (VPS Sir)**  
Vice President  
Exp. : 20 yrs



**Nikhil Srivastava (NS Sir)**  
Head JEE Academics  
Exp. : 17 yrs



**Aatish Agarwal (AA Sir)**  
Sr. Faculty  
Exp. : 17 yrs



**Jayant Chittora (JC Sir)**  
Sr. Faculty  
Exp. : 16 yrs



**Anurag Garg (AG Sir)**  
Sr. Faculty  
Exp. : 17 yrs



**Arjun Gupta (Arjun Sir)**  
Sr. Faculty  
Exp. : 14 yrs



**Devki Nandan Pathak (DN Sir)**  
Sr. Faculty  
Exp. : 13 yrs



**Avinash Kishore (AVN Sir)**  
Sr. Faculty  
Exp. : 9 yrs



**Vipin Sharma (VS Sir)**  
Sr. Faculty  
Exp. : 12 yrs



**Durgesh Pandey (Pandey Sir)**  
Sr. Faculty  
Exp. : 8 yrs

Join

**JEE DROPPER BATCH**

Online + Offline Mode

English & Hindi Medium

Batch Starting from :  
**22nd Sept. 2021**