

# JEE | NEET | Foundation



29900+ SELECTIONS SINCE 2007

## हो चुकी है ऑफलाइन क्लासरूम की शुरूआत अपने सपने को करो साकार, कोटा कोचिंग के साथ



Nitin Vijay (NV Sir) Managing Director Exp. : 18 yrs

#### **Directors of Nucleus Education & Wizard of Mathematics** Now Offline associated with Motion Kota Classroom



Akhilesh Kanther (AKK Sir) Exp. : 17 yrs



Exp. : 18 yrs

Surendra K. Mishra (SKM Sir) Exp.: 16 yrs



Gavesh Bhardwaj (GB Sir) Exp. : 17 yrs

### Academic Pillars of JEE Motion Kota



Ram Ratan Dwivedi (RRD Sir) Joint Director Exp.: 20 yrs



**Amit Verma** (AV Sir) Joint Director Exp.: 16 yrs



Vijay Pratap Singh (VPS Sir) Vice President Exp.: 20 yrs



Nikhil Srivastava (NS Sir) Head JEE Academics Exp. : 17 yrs



**Aatish Agarwal** (AA Sir) Sr. Faculty Exp. : 17 yrs



**Jayant Chittora** (JC Sir) Sr. Faculty Exp.: 16 yrs



**Anurag Garg** (AG Sir) Sr. Faculty Exp.: 17 yrs

Online + Offline Mode



Arjun Gupta (Árjun Sir) Sr. Faculty Exp.: 14 yrs



(DN Sir) Sr. Faculty Exp.: 13 yrs



Devki Nandan Pathak Avinash Kishore (AVN Sir) Sr. Faculty Exp.: 9 yrs



Vipin Sharma (VS Sir) Sr. Faculty Exp.: 12 yrs



**Durgesh Pandey** (Pandey Sir) Sr. Faculty Exp.: 8 yrs

### Join English & Hindi Medium **EE DROPPER BATC**

Batch Starting from: 22nd Sept. 2021

#### **SECTION - A**

- 1. Which of the following is not correct for relation R on the set of real numbers?
  - (1)  $(x, y) \in R \Leftrightarrow |x y| \le 1$  is reflexive and symmetric.
  - (2)  $(x, y) \in R \Leftrightarrow 0 < |x y| \le 1$  is symmetric and transitive.
  - (3)  $(x, y) \in R \Leftrightarrow |x|-|y| \le 1$  is reflexive but not symmetric.
  - (4) (x, y)  $\in$  R  $\Leftrightarrow$   $0<\left|x\right|-\left|y\right|$   $\leq$  1 is neither transitive nor symmetric.

(2) Ans.

Note that (1, 2) and (2, 3) satisfy  $0 < |x-y| \le 1$ Sol.

but (1,3) does not satisfy it so

 $0 \le |x - y| \le 1$  is symmetric but not transitive

So, (2) is correct.

2. Three numbers are in an increasing geometric progression with common ratio r. If the middle number is doubled, then the new numbers are in an arithmetic progression with common difference d. If the fourth term of GP is  $3 r^2$ , then  $r^2$  – d is equal to:

$$(1) 7 + 3\sqrt{3}$$

(2) 
$$7 - \sqrt{3}$$

(2) 
$$7 - \sqrt{3}$$
 (3)  $7 - 7\sqrt{3}$  (4)  $7 + \sqrt{3}$ 

(4) 
$$7 + \sqrt{3}$$

(2) Ans.

Let numbers be  $\frac{a}{r}$ , a, ar  $\rightarrow$  G.P Sol.

$$\frac{a}{r}$$
, 2a,  $ar \rightarrow A.P \Rightarrow 4a = \frac{a}{r} + ar \Rightarrow r + \frac{1}{r} = 4$ 

$$r = 2 \pm \sqrt{3}$$

$$4^{th}$$
 form of G.P =  $3r^2 \Rightarrow ar^2 = 3r^2 \Rightarrow a = 3$ 

$$r = 2 + \sqrt{3}$$
,  $a = 3$ ,  $d = 2a - \frac{a}{r} = 3\sqrt{3}$ 

$$r^2 - d = (2 + \sqrt{3})^2 - 3\sqrt{3}$$

$$= 7 + 4\sqrt{3} - 3\sqrt{3}$$

$$= 7 + \sqrt{3}$$



An Unmatched Experience of Offline



**3.** The sum of 10 terms of the series

$$\frac{3}{1^2 \times 2^2} + \frac{5}{2^2 \times 3^2} + \frac{7}{3^2 \times 4^2} + \dots$$
 is:

- $(1) \frac{120}{121}$
- (2) 1
- (3)  $\frac{143}{144}$
- (4)  $\frac{99}{100}$

**Ans.** (1)

**Sol.** 
$$S = \frac{2^2 - 1^2}{1^2 \times 2^2} + \frac{3^2 - 2^2}{2^2 \times 3^2} + \frac{4^2 - 3^2}{3^2 \times 4^2} + \dots$$

$$= \left[\frac{1}{1^2} - \frac{1}{2^2}\right] + \left[\frac{1}{2^2} - \frac{1}{3^2}\right] + \left[\frac{1}{3^2} - \frac{1}{4^2}\right] + \dots + \left[\frac{1}{10^2} - \frac{1}{11^2}\right]$$

$$= 1 - \frac{1}{121}$$

$$=\frac{120}{121}$$

**4.** The number of real roots of the equation  $e^{4x} + 2e^{3x} - e^x - 6 = 0$  is:


- (1) 0
- (2) 1
- (3) 4
- (4) 2

**Ans.** (2)

**Sol.** Let 
$$e^x = t > 0$$

$$f(t) = t^4 + 2t^3 - t - 6 = 0$$

$$f'(t) = 4t^3 + 6t^2 - 1$$



$$f''(t) = 12t^2 + 12t > 0$$

$$f(0) = -6$$
,  $f(1) = -4$ ,  $f(2) = 24$ 

$$\Rightarrow$$
 Number of real roots = 1



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



The integral  $\int \frac{1}{\sqrt[4]{(x-1)^3(x+2)^5}} dx$  is equal to: 5.

(where C is a constant of integration)

(1) 
$$\frac{3}{4} \left( \frac{x+2}{x-1} \right)^{\frac{5}{4}} + c$$

$$(1) \frac{3}{4} \left( \frac{x+2}{x-1} \right)^{\frac{5}{4}} + c \qquad (2) \frac{3}{4} \left( \frac{x+2}{x-1} \right)^{\frac{1}{4}} + c \qquad (3) \frac{4}{3} \left( \frac{x-1}{x+2} \right)^{\frac{1}{4}} + c \qquad (4) \frac{4}{3} \left( \frac{x-1}{x+2} \right)^{\frac{5}{4}} + c$$

(3) 
$$\frac{4}{3} \left( \frac{x-1}{x+2} \right)^{\frac{1}{4}} +$$

(4) 
$$\frac{4}{3} \left( \frac{x-1}{x+2} \right)^{\frac{5}{4}} + 0$$

Ans.

**Sol.** 
$$\int \frac{dx}{(x-1)^{3/4} (x+2)^{5/4}}$$

$$= \int \frac{dx}{\left(\frac{x+2}{x-1}\right)^{5/4}.(x-1)^2}$$

$$put \ \frac{x+2}{x-1} = t$$

$$=-\frac{1}{3}\int\frac{dt}{t^{5/4}}$$

$$=\frac{4}{3}\cdot\frac{1}{t^{1/4}}+C$$

$$=\frac{4}{3}\left(\frac{x-1}{x+2}\right)^{1/4}+C$$


- A vertical pole fixed to the horizontal ground is divided in the ratio 3:7 by a mark on it with 6. lower part shorter than the upper part. If the two parts subtend equal angles at a point on the ground 18 m away from the base of the pole, then the height of the pole (in meters) is:
  - (1)  $12\sqrt{15}$
- (2)  $8\sqrt{10}$
- (3)  $6\sqrt{10}$
- (4)  $12\sqrt{10}$

(4) Ans.

An Unmatched Experience of Offline



Sol.



Let height of pole =  $10 \ell$ 

$$\tan\alpha = \frac{3\ell}{18} = \frac{\ell}{6}$$

$$tan\,2\alpha=\frac{10\ell}{18}$$

$$\frac{2\tan\alpha}{1-\tan^2\alpha} = \frac{10\ell}{18}$$

Use 
$$\tan \alpha = \frac{\ell}{6} \Rightarrow \ell = \sqrt{\frac{72}{5}}$$

height of pole =  $=10\ell = 12\sqrt{10}$ 

Let f be a non-negative function in [0, 1] and twice differentiable in (0, 1). If 7.

$$\int_0^x \sqrt{1 - \left(f'(t)\right)^2} \ dt \ = \ \int_0^x f(t) \, dt, \\ 0 \le x \le 1 \ \text{ and } f(0) \ = \ 0, \ then \ \lim_{x \to 0} \frac{1}{x^2} \int_0^x f(t) \, dt :$$

- (1) equals 1
- (2) does not exist (3) equals  $\frac{1}{2}$
- (4) equals 0

Ans. (3)

 $\int_{0}^{x} \sqrt{1 - \left(f'(t)\right)^{2}} dt = \int_{0}^{x} f(t) dt 0 \le x \le 1$ Sol.

differentiating both the sides

$$\sqrt{1-\left(f'(x)\right)^2} = f(x)$$

$$\Rightarrow 1 - (f'(x))^2 = f^2(x)$$

An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



# MOTION JEE MAIN 2021

$$\frac{f'(x)}{\sqrt{1-f^2(x)}}=1$$

$$\sin^{-1} f(x) = x + c$$

$$f(0) = 0 \Rightarrow C = 0 \Rightarrow f(x) = \sin x$$

Now 
$$\lim_{x\to 0} \frac{\int_0^x \sin t \, dt}{x^2} \left(\frac{0}{0}\right) = \frac{1}{2}$$

8. If the function 
$$f(x) = \begin{cases} \frac{1}{x} log_e \left( \frac{1 + \frac{x}{a}}{1 - \frac{x}{b}} \right) & , & x < 0 \\ k, & , & x = 0 \end{cases}$$
$$\frac{cos^2 x - sin^2 x - 1}{\sqrt{x^2 + 1} - 1} , & x > 0$$

is continuous at x=0, then  $\frac{1}{a}+\frac{1}{b}+\frac{4}{k}$  is equal to :

$$(1) -4$$

$$(2) -5$$

**Ans.** (2)

**Sol.** If f(x) is continuous at x = 0, RHL = LHL = f(0)

$$\lim_{x \to 0^+} f(x) = \lim_{x \to 0^+} \frac{\cos^2 x - \sin^2 x - 1}{\sqrt{x^2 + 1} - 1} \cdot \frac{\sqrt{x^2 + 1} + 1}{\sqrt{x^2 + 1} + 1} \text{ (Rationalisation)}$$

$$= \lim_{x \to 0^+} -\frac{2\sin^2 x}{x^2} \cdot \left(\sqrt{x^2 + 1} + 1\right) = -4$$

$$\lim_{x\to 0^{-}} f(x) = \lim_{x\to 0^{-}} \frac{1}{x} \ln \left( \frac{1+\frac{x}{a}}{1-\frac{x}{b}} \right)$$

$$\lim_{x\to 0^{-}} \frac{\ell n \left(1 + \frac{x}{a}\right)}{\left(\frac{x}{a}\right).a} + \frac{\ell n \left(1 - \frac{x}{b}\right)}{\left(-\frac{x}{b}\right).b}$$

$$=\frac{1}{a}+\frac{1}{b}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



So 
$$\frac{1}{a} + \frac{1}{b} = -4 = k$$

$$\Rightarrow \frac{1}{a} + \frac{1}{b} + \frac{4}{k} = -4 - 1 = -5$$

- The function  $f(x) = |x^2 2x 3|$ .  $e^{|9x^2 12x + 4|}$  is not differentiable at exactly : 9.
  - (1) two points
- (2) one point
- (3) four points
- (4) three points

Ans. (1)

**Sol.** 
$$f(x) = |(x-3)(x+1)| \cdot e^{(3x-2)^2}$$

$$f(x) = \begin{cases} (x-3)(x+1).e^{(3x-2)^2} & ; \ x \in \ (3,\infty) \\ -(x-3)(x+1).e^{(3x-2)^2} & ; \ x \in \ [-1,3] \\ (x-3).(x+1).e^{(3x-2)^2} & ; \ x \in \ (-\infty,-1) \end{cases}$$

Clearly, non-differentiable at x = -1 & x = 3.

- If  $a_r = cos \frac{2r\pi}{q} + i sin \frac{2r\pi}{q}$ ,  $r = 1, 2, 3, ..., i = \sqrt{-1}$ , then the determinant 10.

  - $\begin{vmatrix} a_1 & a_2 & a_3 \\ a_4 & a_5 & a_6 \\ a_7 & a_8 & a_9 \end{vmatrix}$  is equal to :

  - (1)  $a_2a_6 a_4a_8$  (2)  $a_1a_9 a_3a_7$
- $(3) a_5$
- $(4) a_9$

**Ans.** (2)

**Sol.** 
$$a_r = e^{\frac{i2\pi r}{9}}$$
,  $r = 1, 2, 3, ... . a_1, a_2, a_3, ... . are in G.P.$ 

$$\begin{vmatrix} a_1 & a_2 & a_3 \\ a_n & a_5 & a_6 \\ a_7 & a_8 & a_9 \end{vmatrix} = \begin{vmatrix} a_1 & a_2^2 & a_1^3 \\ a_1^4 & a_1^5 & a_1^6 \\ a_1^7 & a_1^8 & a_1^9 \end{vmatrix} = a_1.a_1^4.a_1^7 \begin{vmatrix} 1 & a_1 & a_1^2 \\ 1 & a_1 & a_1^2 \\ 1 & a_1 & a_1^2 \end{vmatrix} = 0$$

Now 
$$a_1 a_0 - a_3 a_7 = a_1^{10} - a_1^{10} = 0$$

- 11. Let the equation of the plane, that passes through the point (1, 4, -3) and contains the line of intersection of the planes 3x - 2y + 4z - 7 = 0 and x + 5y - 2z + 9 = 0, be  $\alpha x + \beta y + \gamma z + 3 = 0$ 0 then  $\alpha+\beta+\gamma$  is equal to:
  - (1) 15
- (2)15
- (3) -23
- (4)23

An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



(3)Ans.

Equation of plane is Sol.

$$3x - 2y + 4z - 7 + \lambda(x + 5y - 2z + 9) = 0$$

$$(3 + \lambda)x + (5\lambda - 2)y + (4 - 2\lambda)z + 9\lambda - 7 = 0$$

passing through (1, 4, -3)

$$\Rightarrow$$
 3 +  $\lambda$  + 20  $\lambda$  - 8 - 12 + 6  $\lambda$  + 9  $\lambda$  - 7 = 0

$$\Rightarrow \lambda = \frac{2}{3}$$

⇒ equation of plane is

$$-11x - 4y - 8z + 3 = 0$$

$$\Rightarrow \alpha + \beta + \gamma = -23$$

12. The line  $12x \cos\theta + 5y \sin\theta = 60$  is tangent to which of the following curves?

$$(1) 25x^2 + 12y^2 = 3600$$

(2) 
$$144x^2 + 25y^2 = 3600$$

(3) 
$$x^2 + y^2 = 169$$

$$(4) x^2 + y^2 = 60$$

(2) Ans.

Sol.  $12x\cos\theta + 5y\sin\theta = 60$ 

$$\frac{x\cos\theta}{5} + \frac{y\sin\theta}{12} = 1$$

is tangent to 
$$\frac{x^2}{25} + \frac{y^2}{144} = 1$$

$$144x^2 + 25y^2 = 3600$$

If p and q are the lengths of the perpendiculars from the origin on the lines, x cosec  $\alpha$  – y sec  $\alpha$ 13. = k cot  $2\alpha$  and x sin  $\alpha$  + y cos  $\alpha$  = k sin  $2\alpha$  respectively, then  $k^2$  is equal to:

(1) 
$$p^2 + 2q^2$$

(2) 
$$2p^2 + q^2$$

$$(3) p^2 + 4q^3$$

(3) 
$$p^2 + 4q^2$$
 (4)  $4p^2 + q^2$ 

(1) Ans.

First line is  $\frac{x}{\sin \alpha} - \frac{y}{\cos \alpha} = \frac{k \cos 2\alpha}{\sin 2\alpha}$ Sol.

$$\Rightarrow x \cos \alpha - y \sin \alpha = \frac{k}{2} \cos 2\alpha$$

$$\Rightarrow p = \left| \frac{k}{2} \cos \alpha \right| \Rightarrow 2P = |k\cos 2\alpha| \dots (1)$$

second line is  $x \sin \alpha + y \cos \alpha = k \sin 2\alpha$ 



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



$$\Rightarrow$$
 q = | ksin2 $\alpha$  | ...(ii)

Hence 
$$4p^2 + q^2 = k^2$$
 (from (i) & (ii))

$$\textbf{14.} \qquad \lim_{x \to 0} \frac{\sin^2\left(\pi \ cos^4x\right)}{x^4} \ \text{is equal to} \ :$$

(1) 
$$2\pi^2$$

(2) 
$$4\pi$$

(3) 
$$\pi^2$$

(4) 
$$4\pi^2$$

$$\textbf{Sol.} \quad \lim_{x \to 0} \frac{\sin^2\left(\pi \cos^4 x\right)}{x^4}$$

$$=\lim_{x\to 0}\frac{1-\cos\left(2\pi\cos^4x\right)}{2x^4}$$

$$= \lim_{x \to 0} \frac{1 - \cos\left(2\pi - 2\pi\cos^4 x\right)}{\left[2\pi\left(1 - \cos^4 x\right)\right]^2} 4\pi^2 \cdot \frac{\sin^4 x}{2x^4} (1 + \cos^2 x)^2$$

$$=\frac{1}{4}.4\pi^2\,\frac{1}{2}(2)^2\,=4\pi^2$$

15. cosec18° is a root of the equation:

$$(1) x^2 - 2x - 4 = 0$$

$$(2) x^2 - 2x + 4 = 0$$

(1) 
$$x^2 - 2x - 4 = 0$$
 (2)  $x^2 - 2x + 4 = 0$  (3)  $4x^2 + 2x - 1 = 0$  (4)  $x^2 + 2x - 4 = 0$ 

(-) Ans.

**Sol.** 
$$\cos \text{ec} 18^{\circ} = \frac{1}{\sin 18^{\circ}} = \frac{4}{\sqrt{5} - 1} = \sqrt{5} + 1$$

Let 
$$\cos ec 18^0 = x = \sqrt{5} + 1$$

$$\Rightarrow \quad x-1=\sqrt{5}$$

Squaring both sides, we get

$$x^2 - 2x + 1 = 5$$

$$\Rightarrow x^2 - 2x - 4 = 0$$

16. The length of the latus rectum of a parabola, whose vertex and focus are on the positive x-axis at a distance R and S (>R) respectively from the origin, is:

$$(1) 4(S + R)$$

$$(2) 2(S + R)$$

$$(3) 2(S - R)$$

(3) 
$$2(S - R)$$
 (4)  $4(S - R)$ 

Ans. (4)




An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



Sol.



 $V \rightarrow Vertex$ 

 $F \rightarrow focus$ 

$$VF = S - R$$

So latus rectum = 4(S-R)

**17.** If the following system of linear equations

$$2x + y + z = 5$$

$$x - y + z = 3$$

$$x + y + az = b$$

has no solution, then:

(1) 
$$a \neq \frac{1}{3}, b = \frac{7}{3}$$

(2) 
$$a \neq -\frac{1}{3}, b = \frac{7}{3}$$

(3) 
$$a = \frac{1}{3}, b \neq \frac{7}{3}$$

(1) 
$$a \neq \frac{1}{3}$$
,  $b = \frac{7}{3}$  (2)  $a \neq -\frac{1}{3}$ ,  $b = \frac{7}{3}$  (3)  $a = \frac{1}{3}$ ,  $b \neq \frac{7}{3}$  (4)  $a = -\frac{1}{3}$ ,  $b \neq \frac{7}{3}$ 

Ans.

Here D=  $\begin{vmatrix} 2 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & a \end{vmatrix} = 2(-a-1) - 1(a-1) + 1 + 1 = 1 - 3a$ Sol.

$$D_3 = \begin{vmatrix} 2 & 1 & 5 \\ 1 & -1 & 3 \\ 1 & 1 & b \end{vmatrix} = 2(-b-3) - 1(b-3) + 5(1+1) = 7 - 3b$$

for  $a = \frac{1}{3}$ ,  $b \neq \frac{7}{3}$ , system has no solutions



An Unmatched Experience of Offline



**ANSWER KEY** 

Let  $\vec{a}$  and  $\vec{b}$  be two vectors such that  $|2\vec{a}+3\vec{b}|=|3\vec{a}+\vec{b}|$  and the angle between  $\vec{a}$  and  $\vec{b}$  is 60°. 18.

If  $\frac{1}{8}\vec{a}$  is a unit vector, then  $|\vec{b}|$  is equal to:

(1) Ans.

**Sol.** 
$$|3\vec{a} + \vec{b}|^2 = |2\vec{a} + 3\vec{b}|^2$$

$$(3\vec{a} + \vec{b}).(3\vec{a} + \vec{b}) = (2\vec{a} + 3\vec{b}).(2\vec{a} + 3\vec{b})$$

$$9\vec{a}.\vec{a} + 6\vec{a}.\vec{b} + \vec{b}.\vec{b} = 4\vec{a}.\vec{a} + 12\vec{a}.\vec{b} + 9.\vec{b}.\vec{b}$$

$$5|\vec{a}|^2 - 6\vec{a}.\vec{b} = 8|\vec{b}|^2$$

$$5(8)^2 - 6.8. \ \left| \vec{b} \right| \ cos60^\circ = 8 \ \left| \vec{b} \right|^2 \ \left( \because \frac{1}{8} \left| \vec{a} \right| = 1 \right)$$
$$\Rightarrow \left| \vec{a} \right| = 8$$

$$40 - 3\left|\vec{b}\right| = \left|\vec{b}\right|^2$$

$$\Rightarrow \left| \vec{b} \right|^2 + 3 \left| \vec{b} \right| - 40 = 0$$

$$|\vec{\mathbf{b}}| = -8, \quad |\vec{\mathbf{b}}| = 5$$

(rejected)

- If  $\frac{dy}{dx} = \frac{2^{x+y} 2^x}{2^y}$ , y(0) = 1, then y(1) is equal to : 19.
  - (1)  $\log_2(2 + e)$  (2)  $\log_2(1 + e)$
- $(3) log_2(2e)$
- (4)  $\log_2(1 + e^2)$

**Ans.** (2)

$$\textbf{Sol.} \qquad \frac{dy}{dx} = \frac{2^x 2^y - 2^x}{2^y}$$

$$2^{y} \frac{dy}{dx} = 2^{x} (2^{y} - 1)$$

$$\int \frac{2^y}{2^y - 1} dy = \int 2^x dx$$

$$\frac{\ell n(2^y-1)}{\ell n2} = \frac{2^x}{\ell n2} + C$$

$$\Rightarrow \log_2(2^{y}-1) = 2^{x} \log_2 e + C$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



$$y(0) = 1 \Rightarrow 0 = \log_2 e + C$$

$$C = -log_2 e$$

$$\Rightarrow \log_2(2^y - 1) = (2^x - 1)\log_2 e$$

Put 
$$x = 1, \log_2(2^y - 1) = \log_2 e$$

$$2^{y} = e + 1$$

$$y = log_2(e + 1)$$

Let \*,  $\square \in \{\land, \lor\}$  be such that the Boolean expression  $(p * \sim q) \Rightarrow (p \square q)$  is a tautology. 20.

(1) 
$$* = \vee . \square = \wedge$$

(2) 
$$* = \land$$
,  $\square = \lor$ 

$$(3) * = \land \square = \land$$

$$(1) *= \lor, \square = \land \qquad (2) *= \land, \square = \lor \qquad (3) *= \land, \square = \land \qquad (4) *= \lor, \square = \lor$$

Ans.

 $(p \land \neg q) \rightarrow (p \lor q)$  is tautology Sol.

р	q	~q	p∧ ~ q	p∨q	$(p \land \sim q) \to (p \lor q)$
Т	Т	F	F	Т	Т
Т	F	Т	Т	Т	Т
F	Т	F	F	Т	Т
F	F	Т	F	F	Т



An Unmatched Experience of Offline



#### **SECTION-B**

**1.** If  $x \phi(x) = \int_5^x (3t^2 - 2\phi'(t)) dt$ , x > -2, and  $\phi(0) = 4$ , then  $\phi(2)$  is \_\_\_\_\_

**Ans.** (4)

**Sol.** 
$$x\phi(x) = \int_{5}^{x} 3t^2 - 2\phi'(t)dt$$

$$x\phi(x) = x^3 - 125 - 2\lceil \phi(x) - \phi(5) \rceil$$

$$x\phi(x) = x^3 - 125 - 2\phi(x) - 2\phi(5)$$

$$\phi(0)=4 \Rightarrow \phi(5)=-\frac{133}{2}$$

$$\phi(x) = \frac{x^3 + 8}{x + 2}$$

$$\phi(2) = 4$$

- 2. The square of the distance of the point of intersection of the line  $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z+1}{6}$  and the plane 2x y + z = 6 from the point (-1, -1, 2) is \_\_\_\_\_\_.
- **Ans.** (61

**Sol.** 
$$\frac{x-1}{2} = \frac{y-2}{3} = \frac{z+1}{6} = \lambda$$

$$x=2\lambda+1\text{, }y=3\lambda+2\text{, }z=6\lambda-1$$

for point of intersection of line & plane

$$2(2\lambda + 1) - (3\lambda + 2) + (6\lambda - 1) = 6$$

$$7\lambda = 7 \Rightarrow \lambda = 1$$

point: (3,5,5)

$$(distance)^2 = (3+1)^2 + (5+1)^2 + (5-2)^2$$

$$= 16 + 36 + 9 = 61$$

- 3. If  $\left(\frac{3^6}{4^4}\right)k$  is the term, independent of x, in the binomial expansion of  $\left(\frac{x}{4} \frac{12}{x^2}\right)^{12}$ , then k is equal to \_\_\_\_\_.
- **Ans.** (55)



An Unmatched Experience of Offline

## KOTA CLASSROOM For JEE



**Sol.** 
$$\left(\frac{x}{4} - \frac{12}{x^2}\right)^{12}$$

$$T_{r+1} = \left(-1\right)^r \cdot {}^{12}C_r \left(\frac{x}{4}\right)^{12-r} \left(\frac{12}{x^2}\right)^r$$

$$T_{r+1} = (-1)^r \cdot {}^{12}C_r \left(\frac{1}{4}\right)^{12-r} (12)^r \cdot (x)^{12-3r}$$

Term independent of  $x \Rightarrow 12 - 3r = 0 \Rightarrow r = 4$ 

$$T_5 = (-1)^4 \cdot {}^{12}C_4 \left(\frac{1}{4}\right)^8 (12)^4 = \frac{3^6}{4^4} \cdot k$$

$$\Rightarrow$$
 k = 55

- 4. An electric instrument consists of two units. Each unit must function independently for the instrument to operate. The probability that the first unit functions is 0.9 and that of the second unit is 0.8. The instrument is switched on and it fails to operate. If the probability that only the first unit failed and second unit is functioning is p, then 98 p is equal to \_\_\_\_\_\_\_.
- **Ans.** (28)
- **Sol.**  $I_1$  = first unit is functioning

 $I_2$  = second unit is functioning

$$P(I_1) = 0.9, P(I_2) = 0.8$$

$$P(\bar{I}_1) = 0.1, P(\bar{I}_2) = 0.2$$

$$P = \frac{0.8 \times 0.1}{0.1 \times 0.2 + 0.9 \times 0.2 + 0.1 \times 0.8} = \frac{8}{28}$$

$$98P=\frac{8}{28}\times 98=28$$

- A point z moves in the complex plane such that  $\arg\left(\frac{z-2}{z+2}\right) = \frac{\pi}{4}$ , then the minimum value of  $\left|z-9\sqrt{2}-2i\right|^2$  is equal to \_\_\_\_\_\_.
- **Ans.** (98)

**Sol.** Let 
$$z = x + iy$$

$$arg\left(\frac{x-2+iy}{x+2+iy}\right) = \frac{\pi}{4}$$

arg 
$$(x - 2 + iy) - arg (x + 2 + iy) = \frac{\pi}{4}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



$$tan^{-1}\left(\frac{y}{x-2}\right) - tan^{-1}\left(\frac{y}{x+2}\right) = \frac{\pi}{4}$$


$$\frac{\frac{y}{x-2} - \frac{y}{x+2}}{1 + \left(\frac{y}{x-2}\right) \cdot \left(\frac{y}{x+2}\right)} = \tan \frac{\pi}{4} = 1$$

$$\frac{xy + 2y - xy + 2y}{x^2 - 4 + y^2} = 1$$

$$4y = x^2 - 4 + y^2$$

$$x^2 + v^2 - 4v - 4 = 0$$

locus is a circle with centre (0, 2) & radius =  $2\sqrt{2}$ 



min. value =  $(AP)^2 = (OP-OA)^2$ 


$$= \left(9\sqrt{2} - 2\sqrt{2}\right)^2$$

$$= \left(7\sqrt{2}\right)^2 = 98$$

**6.** The number of six letter words (with or without meaning), formed using all the letters of the word 'VOWELS', so that all the consonants never come together, is \_\_\_\_\_.

**Ans.** (576)

Sol.



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE



All Consonants should not be together

- = Total All consonants together,
- = 6! 3! 4! = 576
- **7.** The mean of 10 numbers  $7 \times 8$ ,  $10 \times 10$ ,  $13 \times 12$ ,  $16 \times 14$ , ... is

**Ans.** (398)

**Sol.**  $7 \times 8$ ,  $10 \times 10$ ,  $13 \times 12$ ,  $16 \times 14$  ...

$$T_n = (3n + 4)(2n + 6) = 2(3n + 4)(n + 3)$$

$$= 2(3n^2 + 13n + 12) = 6n^2 + 26n + 24$$

$$S_{10} = \sum_{n=1}^{10} T_n = 6\sum_{n=1}^{10} n^2 + 26\sum_{n=1}^{10} n + 24\sum_{n=1}^{10} 1$$

$$= \frac{6(10 \times 1 \times 21)}{6} + 26 \times \frac{10 \times 11}{2} + 24 \times 10$$

$$= 10 \times 11 (21 + 13) + 240$$

= 3980

Mean = 
$$\frac{S_{10}}{10} = \frac{3980}{10} = 398$$

8. If 'R' is the least value of 'a' such that the function  $f(x) = x^2 + ax + 1$  is increasing on [1, 2] and 'S" is the greatest value of 'a' such that the function  $f(x) = x^2 + ax + 1$  is decreasing on [1, 2], then the value of |R - S| is \_\_\_\_\_\_.

**Ans.** (2)

**Sol.**  $f(x) = x^2 + ax + 1$ 

$$f'(x) = 2x + a$$

when f(x) is increasing on [1, 2]

$$2x + a \ge 0 \ \forall \ x \in [1, 2]$$

$$a \ge -2x \ \forall \ x \in [1, 2]$$

$$R = -4$$

when f(x) is decreasing on [1, 2]

$$2x + a \le 0 \forall x \in [1, 2]$$

$$a \leq -2 \ \forall \ x \in [1, 2]$$

$$S = -2$$

$$|R - S| = |-4 + 2| = 2$$

- **9.** Let [t] denote the greatest integer  $\leq$  t. Then the value of 8.  $\int_{-\frac{1}{2}}^{1} ([2x] + |x|) dx$  is \_\_\_\_\_\_.
- **Ans.** (5)




An Unmatched Experience of Offline


KOTA CLASSROOM For JEE



### **ANSWER KEY**

# Motion<sup>™</sup> JEE MAIN 2021





$$= \int_{-1/2}^{1} \left[ 2x \right] dx + \int_{-1/2}^{1} |x| dx$$

$$= \left( -\frac{x^2}{2} \right)_{-1/2}^{0} + \left( \frac{x^2}{2} \right)_{0}^{1}$$

$$= \left( 0 + \frac{1}{8} \right) + \frac{1}{2}$$

$$= \frac{5}{8}$$

10. If the variable line  $3x + 4y = \alpha$  lies between the two circles  $(x-1)^2 + (y-1)^2 = 1$  and  $(x-9)^2 + (y-1)^2 = 4$ , without intercepting a chord on either circle, then the sum of all the integral values of  $\alpha$  is \_\_\_\_\_\_.

**Ans.** (165)


8I = 5



An Unmatched Experience of Offline



Sol.



Both centres should lie on either side of the line as well as line can be tangent to circle.

$$(3 + 4 - \alpha) \cdot (27 + 4 - \alpha) < 0$$

$$(7 - \alpha) \cdot (31 - \alpha) < 0 \Rightarrow \alpha \in (7, 31)$$
 ...(1)

$$d_1$$
 = distance of (1, 1) from line

$$d_2$$
 = distance of (9, 1) from line

$$d_1 \ge r_1 \Rightarrow \frac{|7-\alpha|}{5} \ge 1 \Rightarrow \alpha \in (-\infty,2) \cup [12,\infty)$$

$$d_1 \ge r_2 \Rightarrow \frac{|31 - \alpha|}{5} \ge 2 \Rightarrow \alpha \in (-\infty, 21) \cup [41, \infty) \quad ...(3)$$

$$(1) \cap (2) \cap (3) \Rightarrow \alpha \in [12, 21]$$

Sum of integers = 165





## हो चुकी है ऑफलाइन क्लासरूम की शुरूआत अपने सपने को करो साकार, कोटा कोचिंग के साथ



Nitin Vijay (NV Sir) Managing Director Exp. : 18 yrs

### **Directors of Nucleus Education & Wizard of Mathematics** Now Offline associated with Motion Kota Classroom



Akhilesh Kanther (AKK Sir) Exp. : 17 yrs

Vishal Joshi (VJ Sir) Exp. : 18 yrs

Surendra K. Mishra (SKM Sir) Exp.: 16 yrs



Gavesh Bhardwai (GB Sir) Exp. : 17 yrs

### Academic Pillars of JEE Motion Kota



Ram Ratan Dwivedi (RRD Sir) Joint Director Exp.: 20 yrs



**Amit Verma** (AV Sir) Joint Director Exp.: 16 yrs



Vijay Pratap Singh (VPS Sir) Vice President Exp.: 20 yrs



Nikhil Srivastava (NS Sir) Head JEE Academics Exp.: 17 yrs



**Aatish Agarwal** (AA Sir) Sr. Faculty Exp. : 17 yrs



Jayant Chittora (JC Sir) Sr. Faculty Exp. : 16 yrs



**Anurag Garg** (AG Sir) Sr. Faculty Exp.: 17 yrs



Arjun Gupta (Arjun Sir) Sr. Faculty Exp.: 14 yrs



Devki Nandan Pathak Avinash Kishore (DN Sir) Sr. Faculty Exp.: 13 yrs



(AVN Sir) Sr. Faculty Exp. : 9 yrs



Vipin Sharma (VS Sir) Sr. Faculty Exp.: 12 yrs



**Durgesh Pandey** (Pandey Sir) Sr. Faculty Exp. : 8 yrs

### Join **English & Hindi Medium** DROPPER BATCH

Online + Offline Mode

Batch Starting from: 22nd Sept. 2021



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

