



**JEE
MAIN
4th
Attempt**

MATHEMATICS
31st August 2021 [SHIFT - 2]
QUESTION WITH SOLUTION

JEE | NEET | Foundation

MOTION[®]

29900+
SELECTIONS SINCE 2007

हो चुकी है ऑफलाइन क्लासरूम की शुरुआत
अपने सपने को करो साकार, कोटा कोचिंग के साथ

Directors of Nucleus Education & Wizard of Mathematics

Now Offline associated with Motion Kota Classroom



Nitin Vijay (NV Sir)
Managing Director
Exp. : 18 yrs



Akhilesh Kanther (AKK Sir)
Exp. : 17 yrs

Vishal Joshi (VJ Sir)
Exp. : 18 yrs

Surendra K. Mishra (SKM Sir)
Exp. : 16 yrs



Gavesh Bhardwaj (GB Sir)
Exp. : 17 yrs

Academic Pillars of JEE MOTION KOTA



Ram Ratan Dwivedi (RRD Sir)
Joint Director
Exp. : 20 yrs



Amit Verma (AV Sir)
Joint Director
Exp. : 16 yrs



Vijay Pratap Singh (VPS Sir)
Vice President
Exp. : 20 yrs



Nikhil Srivastava (NS Sir)
Head JEE Academics
Exp. : 17 yrs



Aatish Agarwal (AA Sir)
Sr. Faculty
Exp. : 17 yrs



Jayant Chittora (JC Sir)
Sr. Faculty
Exp. : 16 yrs



Anurag Garg (AG Sir)
Sr. Faculty
Exp. : 17 yrs



Arjun Gupta (Arjun Sir)
Sr. Faculty
Exp. : 14 yrs



Devki Nandan Pathak (DN Sir)
Sr. Faculty
Exp. : 13 yrs



Avinash Kishore (AVN Sir)
Sr. Faculty
Exp. : 9 yrs



Vipin Sharma (VS Sir)
Sr. Faculty
Exp. : 12 yrs



Durgesh Pandey (Pandey Sir)
Sr. Faculty
Exp. : 8 yrs

Join

JEE DROPPER BATCH

Online + Offline Mode

English & Hindi Medium

Batch Starting from :
22nd Sept. 2021

SECTION - A

1. The sum of the roots of the equation, $x + 1 - 2 \log_2 (3 + 2^x) + 2 \log_4 (10 - 2^{-x}) = 0$, is:
 (1) $\log_2 12$ (2) $\log_2 14$ (3) $\log_2 11$ (4) $\log_2 13$

Ans. (3)

Sol. $x + 1 - 2 \log_2 (3 + 2^x) + 2 \log_4 (10 - 2^{-x}) = 0$
 $\log_2 (2^{x+1}) - \log_2 (3 + 2^x)^2 + \log_2 (10 - 2^{-x}) = 0$

$$\log_2 \left(\frac{2^{x+1} \cdot (10 - 2^{-x})}{(3 + 2^x)^2} \right) = 0$$

$$\frac{2(10 \cdot 2^x - 1)}{(3 + 2^x)^2} = 1$$

$$\Rightarrow 20 \cdot 2^x - 2 = 9 + 2^{2x} + 6 \cdot 2^x$$

$$\therefore (2^x)^2 - 14(2^x) + 11 = 0$$

Roots are 2^{x_1} & 2^{x_2}

$$\therefore 2^{x_1} \cdot 2^{x_2} = 11$$

$$x_1 + x_2 = \log_2(11)$$

2. The distance of the point $(-1, 2, -2)$ from the line of intersection of the planes $2x + 3y + 2z = 0$ and $x - 2y + z = 0$ is:

- (1) $\frac{\sqrt{42}}{2}$ (2) $\frac{5}{2}$ (3) $\frac{1}{\sqrt{2}}$ (4) $\frac{\sqrt{34}}{2}$

Ans. (4)

Sol. $P_1 : 2x + 3y + 2z = 0$

$$\Rightarrow \vec{n}_1 = 2\hat{i} + 3\hat{j} + 2\hat{k}$$

$$P_2 : x - 2y + z = 0$$

$$\Rightarrow \vec{n}_2 = \hat{i} - 2\hat{j} + \hat{k}$$

Direction vector of line L which is line of intersection of P_1 & P_2

$$\vec{r} = \vec{n}_1 \times \vec{n}_2 = 7\hat{i} - 7\hat{k}$$

DR's of L are $(1, 0, -1)$

$$\Rightarrow \text{Equation of L : } \frac{x}{1} = \frac{y}{0} = \frac{z}{-1} = \lambda$$




An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**





DR's of $\overline{PQ} = (\lambda + 1, -2, 2 - \lambda)$

$\therefore \overline{PQ} \perp \vec{r}$

$$\Rightarrow (\lambda + 1)(1) + (-2)(0) + (2 - \lambda)(-1) = 0$$

$$\Rightarrow \lambda = \frac{1}{2} \Rightarrow Q\left(\frac{1}{2}, 0, \frac{-1}{2}\right)$$

$$\Rightarrow PQ = \frac{\sqrt{34}}{2}$$

3. If $\alpha = \lim_{x \rightarrow \pi/4} \frac{\tan^3 x - \tan x}{\cos\left(x + \frac{\pi}{4}\right)}$ and $\beta = \lim_{x \rightarrow 0} (\cos x)^{\cot x}$ are the roots of the equation, $ax^2 + bx - 4 = 0$,

then the ordered pair (a, b) is:

(1) (-1, -3)

(2) (-1, 3)

(3) (1, 3)

(4) (1, -3)

Ans. (3)

Sol. $\alpha = \lim_{x \rightarrow \frac{\pi}{4}} \frac{\tan^3 x - \tan x}{\cos\left(x + \frac{\pi}{4}\right)}$; $\frac{0}{0}$ form

Using L Hospital rule

$$\alpha = \lim_{x \rightarrow \frac{\pi}{4}} \frac{3 \tan^2 x \sec^2 x - \sec^2 x}{-\sin\left(x + \frac{\pi}{4}\right)}$$

$$\Rightarrow \alpha = -4$$

$$\beta = \lim_{x \rightarrow 0} (\cos x)^{\cot x} = e^{\lim_{x \rightarrow 0} \frac{(\cos x - 1)}{\tan x}}$$

$$\beta = e^{\lim_{x \rightarrow 0} \frac{-(1 - \cos x)}{x^2} \cdot \frac{x^2}{\left(\frac{\tan x}{x}\right)^x}}$$

$$\beta = e$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



$$\beta = e^{\lim_{x \rightarrow 0} \left(\frac{-1}{2}\right)^{\frac{x}{1}}} = e^0 \Rightarrow \beta = 1$$

$$\alpha = -4; \beta = 1$$

If $ax^2 + bx - 4 = 0$ are the roots then

$$16a - 4b - 4 = 0 \text{ \& } a + b - 4 = 0$$

$$\Rightarrow a = 1 \text{ \& } b = 3$$

4. Let f be any continuous function on $[0, 2]$ and twice differentiable on $(0, 2)$. If $f(0) = 0$, $f(1) = 1$ and $f(2) = 2$, then:

(1) $f''(x) > 0$ for all $x \in (0, 2)$

(2) $f'(x) = 0$ for some $x \in [0, 2]$

(3) $f''(x) = 0$ for all $x \in (0, 2)$

(4) $f''(x) = 0$ for some $x \in (0, 2)$

Ans. (4)

Sol. $f(0) = 0$, $f(1) = 1$ and $f(2) = 2$

Let $h(x) = f(x) - x$ has three roots

By Rolle's theorem $h'(x) = f'(x) - 1$ has at least two roots

$h''(x) = f''(x) = 0$ has at least one roots.

5. Negation of the statement $(p \vee r) \Rightarrow (q \vee r)$ is:

(1) $p \wedge q \wedge r$

(2) $\sim p \wedge q \wedge r$

(3) $p \wedge \sim q \wedge \sim r$

(4) $\sim p \wedge q \wedge \sim r$

Ans. (3)

Sol. $\therefore \sim(A \Rightarrow B) = A \wedge \sim B$

$$\therefore \sim((p \vee r) \Rightarrow (q \vee r))$$

$$= (p \vee r) \wedge (\sim q \wedge \sim r)$$

$$= ((p \vee r) \wedge (\sim r)) \wedge (\sim q)$$

$$= p \wedge (\sim r) \wedge (\sim q)$$

6. If $\alpha + \beta + \gamma = 2\pi$, then the system of equations

$$x + (\cos \gamma) y + (\cos \beta)z = 0$$

$$(\cos \gamma) x + y + (\cos \alpha)z = 0$$

$$(\cos \beta) x + (\cos \alpha)y + z = 0$$

has:

(1) exactly two solutions

(2) a unique solution

(3) no solution

(4) infinitely many solutions

Ans. (4)



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



Sol. $\alpha + \beta + \gamma = 2\pi$

$$\begin{vmatrix} 1 & \cos \gamma & \cos \beta \\ \cos \gamma & 1 & \cos \alpha \\ \cos \beta & \cos \alpha & 1 \end{vmatrix}$$

$$\begin{aligned} &= 1 + 2\cos\alpha \cdot \cos\beta \cdot \cos\gamma - \cos^2\alpha - \cos^2\beta - \cos^2\gamma \\ &= \sin^2\gamma - \cos^2\alpha - \cos^2\beta + (\cos(\alpha + \beta) + \cos(\alpha - \beta)) \cos\gamma \\ &= \sin^2\gamma - \cos^2\alpha - \cos^2\beta + \cos^2\gamma + \cos(\alpha - \beta) \cos\gamma \\ &= \sin^2\alpha - \cos^2\beta + \cos(\alpha - \beta) \cos(\alpha + \beta) \\ &= \sin^2\alpha - \cos^2\beta + \cos^2\alpha - \sin^2\beta = 0 \end{aligned}$$

7. Let a_1, a_2, a_3, \dots be an A.P. If $\frac{a_1 + a_2 + \dots + a_{10}}{a_1 + a_2 + \dots + a_p} = \frac{100}{p^2}$, $p \neq 10$, then $\frac{a_{11}}{a_{10}}$ is equal to:

- (1) $\frac{121}{100}$ (2) $\frac{100}{121}$ (3) $\frac{19}{21}$ (4) $\frac{21}{19}$

Ans. (4)

Sol.
$$\frac{\frac{10}{2}(2a_1 + 9d)}{\frac{p}{2}(2a_1 + (p-1)d)} = \frac{100}{p^2}$$

$$(2a_1 + 9d)p = 10(2a_1 + (p-1)d)$$

$$9dp = 20a_1 - 2pa_1 + 10d(p-1)$$

$$9p = (20 - 2p) \frac{a_1}{d} + 10(p-1)$$

$$\frac{a_1}{d} = \frac{(10-p)}{2(10-p)} = \frac{1}{2}$$

$$\therefore \frac{a_{11}}{a_{10}} = \frac{a_1 + 10d}{a_1 + 9d} = \frac{\frac{1}{2} + 10}{\frac{1}{2} + 9} = \frac{21}{19}$$

8. Let $\vec{a}, \vec{b}, \vec{c}$ be three vectors mutually perpendicular to each other and have same magnitude. If a vector \vec{r} satisfies

$$\vec{a} \times \{(\vec{r} - \vec{b}) \times \vec{a}\} + \vec{b} \times \{(\vec{r} - \vec{c}) \times \vec{b}\} + \vec{c} \times \{(\vec{r} - \vec{a}) \times \vec{c}\} = \vec{0}, \text{ then } \vec{r} \text{ is equal to:}$$

- (1) $\frac{1}{2}(\vec{a} + \vec{b} + 2\vec{c})$ (2) $\frac{1}{3}(2\vec{a} + \vec{b} - \vec{c})$ (3) $\frac{1}{3}(\vec{a} + \vec{b} + \vec{c})$ (4) $\frac{1}{2}(\vec{a} + \vec{b} + \vec{c})$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



Ans. (4)

Sol. Suppose, $\vec{r} = x\vec{a} + y\vec{b} + 2\vec{c}$

$$\text{and } |\vec{a}| = |\vec{b}| = |\vec{c}| = k$$

$$\vec{a} \times \{(\vec{r} - \vec{b}) \times \vec{a}\} + \vec{b} \times \{(\vec{r} - \vec{c}) \times \vec{b}\} + \vec{c} \times \{(\vec{r} - \vec{a}) \times \vec{c}\} = \vec{0}$$

$$\Rightarrow k^2(\vec{r} - \vec{b}) - k^2x\vec{a} + k^2(\vec{r} - \vec{c}) - k^2y\vec{b} + k^2(\vec{r} - \vec{a}) - k^2z\vec{c} = \vec{0}$$

$$\Rightarrow 3\vec{r} - (\vec{a} + \vec{b} + \vec{c}) - \vec{r} = \vec{0}$$

$$\Rightarrow \vec{r} = \frac{\vec{a} + \vec{b} + \vec{c}}{2}$$

9. The locus of mid-points of the line segments joining $(-3, -5)$ and the points on the ellipse

$$\frac{x^2}{4} + \frac{y^2}{9} = 1 \text{ is:}$$

$$(1) 36x^2 + 16y^2 + 90x + 56y + 145 = 0 \quad (2) 36x^2 + 16y^2 + 108x + 80y + 145 = 0$$

$$(3) 36x^2 + 16y^2 + 72x + 32y + 145 = 0 \quad (4) 9x^2 + 4y^2 + 18x + 8y + 145 = 0$$

Ans. (2)

Sol. General point on $\frac{x^2}{4} + \frac{y^2}{9} = 1$, is A $(2\cos\theta, 3\sin\theta)$

Given, B $(-3, -5)$

$$\text{Midpoint C} \left(\frac{2\cos\theta - 3}{2}, \frac{3\sin\theta - 5}{2} \right)$$

$$h = \frac{2\cos\theta - 3}{2}; k = \frac{3\sin\theta - 5}{2}$$

$$\Rightarrow \left(\frac{2h+3}{2} \right)^2 + \left(\frac{2k+5}{3} \right)^2 = 1$$

$$\Rightarrow 36x^2 + 16y^2 + 108x + 80y + 145 = 0$$

10. If $\frac{dy}{dx} = \frac{2^x y + 2^y \cdot 2^x}{2^x + 2^{x+y} \log_e 2}$, $y(0) = 0$, then for $y = 1$, the value of x lies in the interval:

$$(1) \left[\frac{1}{2}, 1 \right]$$

$$(2) \left[0, \frac{1}{2} \right]$$

$$(3) (1, 2)$$

$$(4) (2, 3)$$

Ans. (3)



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



Sol. $\frac{dy}{dx} = \frac{2^x(y+2^y)}{2^x(1+2^y \ln 2)}$

$$\Rightarrow \int \frac{(1+2^y)\ln 2}{(y+2^y)} dy = \int dx$$

$$\Rightarrow \ln|y+2^y| = x + c$$

$$x = 0; y = 0 \Rightarrow c = 0$$

$$\Rightarrow x = \ln|y+2^y|$$

$$\Rightarrow \text{at } y = 1, x = \ln 3$$

$$\therefore 3 \in (e, e^2) \Rightarrow x \in (1, 2)$$

11. If z is a complex number such that $\frac{z-i}{z-1}$ is purely imaginary, then the minimum value of $|z - (3 + 3i)|$ is

- (1) $3\sqrt{2}$ (2) $6\sqrt{2}$ (3) $2\sqrt{2}$ (4) $2\sqrt{2} - 1$

Ans. (3)

Sol. $\frac{z-i}{z-1}$ is purely Imaginary number


Let $z = x + iy$

$$\therefore \frac{x + i(y-1)}{(x-1) + iy} \times \frac{(x-1) - iy}{(x-1) - iy}$$

$$\Rightarrow \frac{x(x-1) + y(y-1) + i(-y-x+1)}{(x-1)^2 + y^2} \text{ is purely Imaginary number}$$

$$\Rightarrow x(x-1) + y(y-1) = 0$$

$$\Rightarrow \left(x - \frac{1}{2}\right)^2 + \left(y - \frac{1}{2}\right)^2 = \frac{1}{2}$$



$$\therefore |z - (3 + 3i)|_{\min} = |PC| - \frac{1}{\sqrt{2}}$$

$$= \frac{5}{\sqrt{2}} - \frac{1}{\sqrt{2}} = 2\sqrt{2}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



12. The mean and variance of 7 observations are 8 and 16 respectively. If two observations are 6 and 8, then the variance of the remaining 5 observations is:

- (1) $\frac{92}{5}$ (2) $\frac{134}{5}$ (3) $\frac{536}{25}$ (4) $\frac{112}{5}$

Ans. (3)

Sol. Let, 8, 16, x_1, x_2, x_3, x_4, x_5 be the observations.

$$\text{Now, } \frac{x_1 + x_2 + \dots + x_5 + 14}{7} = 8$$

$$\Rightarrow \sum_{i=1}^5 x_i = 42 \quad \dots(1)$$

$$\text{Also, } \frac{x_1^2 + x_2^2 + \dots + x_5^2 + 8^2 + 6^2}{7} - 64 = 16$$

$$\Rightarrow \sum_{i=1}^5 x_i^2 = 560 - 100 = 460 \quad \dots(2)$$

So, variance of x_1, x_2, \dots, x_5

$$= \frac{460}{5} - \left(\frac{42}{5}\right)^2 = \frac{2300 - 1764}{25} = \frac{536}{25}$$

13. The number of solutions of the equation $32^{\tan^2 x} + 32^{\sec^2 x} = 81, 0 \leq x \leq \frac{\pi}{4}$ is:

- (1) 0 (2) 1 (3) 2 (4) 3

Ans. (2)

Sol. $(32)^{\tan^2 x} + (32)^{\sec^2 x} = 81$

$$\Rightarrow (32)^{\tan^2 x} + (32)^{1+\tan^2 x} = 81$$

$$\Rightarrow (32)^{\tan^2 x} = \frac{81}{33}$$

In interval $\left[0, \frac{\pi}{4}\right]$ only one solution.

14. Let $S = \{1, 2, 3, 4, 5, 6\}$. Then the probability that a randomly chosen onto function g from S to S satisfies $g(3) = 2g(1)$ is:

- (1) $\frac{1}{5}$ (2) $\frac{1}{30}$ (3) $\frac{1}{15}$ (4) $\frac{1}{10}$

Ans. (4)



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**




Sol. $g(3) = 2g(1)$ can be defined in 3 ways
 number of onto functions in this condition = $3 \times 4!$
 Total number of onto functions = $6!$
 Required probability = $\frac{3 \times 4!}{6!} = \frac{1}{10}$

15. Let A be the set of all points (α, β) such that the area of triangle formed by the points (5,6), (3, 2) and (α, β) is 12 square units. Then the least possible length of a line segment joining the origin to a point in A, is:

- (1) $\frac{16}{\sqrt{5}}$ (2) $\frac{12}{\sqrt{5}}$ (3) $\frac{8}{\sqrt{5}}$ (4) $\frac{4}{\sqrt{5}}$

Ans. (3)

Sol.



$$\left| \begin{array}{ccc} 5 & 6 & 1 \\ 3 & 2 & 1 \\ \alpha & \beta & 1 \end{array} \right| = 12$$

$$4\alpha - 2\beta = \pm 24 + 8$$

$$\Rightarrow 4\alpha - 2\beta = +24 + 8 \Rightarrow 2\alpha - \beta = 16$$

$$2x - y - 16 = 0 \quad \dots(1)$$

$$\Rightarrow 4\alpha - 2\beta = -24 + 8 \Rightarrow 2\alpha - \beta = -8$$

$$2x - y + 8 = 0 \quad \dots(2)$$

Perpendicular distance of (1) from (0, 0)

$$\left| \frac{0 - 0 - 16}{\sqrt{5}} \right| = \frac{16}{\sqrt{5}}$$

Perpendicular distance of (2) from (0, 0) is $\left| \frac{0 - 0 + 8}{\sqrt{5}} \right| = \frac{8}{\sqrt{5}}$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



16. An angle of intersection of the curves, $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and $x^2 + y^2 = ab$, $a > b$, is:

- (1) $\tan^{-1}\left(\frac{a-b}{2\sqrt{ab}}\right)$ (2) $\tan^{-1}\left(\frac{a+b}{\sqrt{ab}}\right)$ (3) $\tan^{-1}\left(\frac{a-b}{\sqrt{ab}}\right)$ (4) $\tan^{-1}(2\sqrt{ab})$

Ans. (3)

Sol. $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, x^2 + y^2 = ab$

$$\frac{2x_1}{a^2} + \frac{2y_1 y'}{b^2} = 0$$

$$\Rightarrow y_1' = \frac{-x_1 b^2}{a^2 y_1} \quad \dots(1)$$

$$\therefore 2x_1 + 2y_1 y' = 0$$

$$\Rightarrow y_2' = \frac{-x_1}{y_1} \quad \dots(2)$$

Here (x_1, y_1) is point of intersection of both curves

$$\therefore x_1^2 = \frac{a^2 b}{a+b}, y_1^2 = \frac{ab^2}{a+b}$$

$$\therefore \tan \theta = \left| \frac{y_1' - y_2'}{1 + y_1' y_2'} \right| = \left| \frac{\frac{-x_1 b^2}{a^2 y_1} + \frac{x_1}{y_1}}{1 + \frac{x_1^2 b^2}{a^2 y_1^2}} \right|$$

$$\tan \theta = \left| \frac{-b^2 x_1 y_1 + a^2 x_1 y_1}{a^2 y_1^2 + b^2 x_1^2} \right|$$

$$\tan \theta = \left| \frac{a-b}{\sqrt{ab}} \right|$$

17. If $[x]$ is the greatest integer $\leq x$, then $\pi^2 \int_0^2 \left(\sin \frac{\pi x}{2} \right) (x - [x])^{[x]} dx$ is equal to:

- (1) $4(\pi - 1)$ (2) $2(\pi - 1)$ (3) $2(\pi + 1)$ (4) $4(\pi + 1)$

Ans. (1)

Sol. $\pi^2 \left[\int_0^1 \sin \frac{\pi x}{2} dx + \int_1^2 \sin \frac{\pi x}{2} (x-1) dx \right]$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



$$\begin{aligned}
 &= \pi^2 \left[-\frac{2}{\pi} \left(\cos \frac{\pi x}{2} \right) + \left((x-1) \left(-\frac{2}{\pi} \cos \frac{\pi x}{2} \right) \right)_1^2 - \int_1^2 -\frac{2}{\pi} \cos \frac{\pi x}{2} dx \right] \\
 &= \pi^2 \left[0 + \frac{2}{\pi} + \frac{2}{\pi} + \frac{2}{\pi} \cdot \frac{2}{\pi} \left(\sin \frac{\pi x}{2} \right)_1^2 \right] \\
 &= 4\pi - 4 = 4(\pi - 1)
 \end{aligned}$$

18. Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be a function such that $f(m+n) = f(m) + f(n)$ for every $m, n \in \mathbb{N}$. If $f(6) = 18$, then $f(2) \cdot f(3)$ is equal to:

- (1) 36 (2) 6 (3) 18 (4) 54

Ans. (4)

Sol. $f(m+n) = f(m) + f(n)$

Put $m = 1, n = 1$

$f(2) = 2f(1)$

Put $m = 2, n = 1$

$f(3) = f(2) + f(1) = 3f(1)$

Put $m = 3, n = 3$

$f(6) = 2f(3) \Rightarrow f(3) = 9$

$\Rightarrow f(1) = 3, f(2) = 6$

$f(2) \cdot f(3) = 6 \times 9 = 54$

19. The domain of the function

$f(x) = \sin^{-1} \left(\frac{3x^2 + x - 1}{(x-1)^2} \right) + \cos^{-1} \left(\frac{x-1}{x+1} \right)$ is:

- (1) $\left[\frac{1}{4}, \frac{1}{2} \right] \cup \{0\}$ (2) $[-2, 0] \cup \left[\frac{1}{4}, \frac{1}{2} \right]$ (3) $\left[0, \frac{1}{4} \right]$ (4) $\left[0, \frac{1}{2} \right]$

Ans. (1)

Sol. $f(x) = \sin^{-1} \left(\frac{3x^2 + x - 1}{(x-1)^2} \right) + \cos^{-1} \left(\frac{x-1}{x+1} \right)$

$$-1 \leq \frac{x-1}{x+1} \leq 1 \Rightarrow 0 \leq x < \infty \quad \dots(1)$$

$$-1 \leq \frac{3x^2 + x - 1}{(x-1)^2} \leq 1 \Rightarrow x \in \left[\frac{-1}{4}, \frac{1}{2} \right] \cup \{0\} \quad \dots(2)$$

Equation (1) & (2)

$$\Rightarrow \text{Domain} = \left[\frac{1}{4}, \frac{1}{2} \right] \cup \{0\}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



20. If $y \frac{dy}{dx} = x \left[\frac{y^2}{x^2} + \frac{\phi\left(\frac{y^2}{x^2}\right)}{\phi'\left(\frac{y^2}{x^2}\right)} \right]$, $x > 0$, $\phi > 0$ and $y(1) = -1$, then $\phi\left(\frac{y^2}{4}\right)$ is equal to:

(1) $\phi(1)$

(2) $4\phi(2)$

(3) $2\phi(1)$

(4) $4\phi(1)$

Ans. (4)

Sol. Let, $y = tx$

$$\frac{dy}{dx} = t + x \frac{dt}{dx}$$

$$\therefore tx \left(t + x \frac{dt}{dx} \right) = x \left(t^2 + \frac{\phi(t^2)}{\phi'(t^2)} \right)$$

$$t^2 + xt \frac{dt}{dx} = t^2 + \frac{\phi(t^2)}{\phi'(t^2)}$$

$$\int \frac{t\phi'(t^2)}{\phi(t^2)} dt = \int \frac{dx}{x}$$

Let $\phi(t^2) = p$

$$\therefore \phi'(t^2) 2t dt = dp$$

$$\Rightarrow \int \frac{dy}{2p} = \int \frac{dx}{x}$$

$$\frac{1}{2} \ln \phi(t^2) = \ln x + \ln c$$

$$\phi(t^2) = x^2 k$$

$$\phi\left(\frac{y^2}{x^2}\right) = kx^2, \phi(1) = k$$

$$\phi\left(\frac{y^2}{4}\right) = 4\phi(1)$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



SECTION-B

1. Suppose the line $\frac{x-2}{\alpha} = \frac{y-2}{-5} = \frac{z+2}{2}$ lies on the plane $x + 3y - 2z + \beta = 0$, then $(\alpha+\beta)$ is equal to ____

Ans. (7)

Sol. Point $(2, 2, -2)$ also lies on given plane

$$\text{So, } 2 + 3 \times 2 - 2(-2) + \beta = 0$$

$$\Rightarrow 2 + 6 + 4 + \beta = 0 \Rightarrow \beta = -12$$

$$\text{Also, } \alpha \times 1 - 5 \times 3 + 2 \times -2 = 0$$

$$\Rightarrow \alpha - 15 - 4 = 0 \Rightarrow \alpha = 19$$

$$\therefore \alpha + \beta = 19 - 12 = 7$$

2. If $\int \frac{\sin x}{\sin^3 x + \cos^3 x} dx = \alpha \log_e |1 + \tan x| + \beta \log_e |1 - \tan x + \tan^2 x| + \gamma \tan^{-1} \left(\frac{2 \tan x - 1}{\sqrt{3}} \right) + C$,

When C is constant of integration, then the value of $18(\alpha + \beta + \gamma^2)$ is _____

Ans. (3)

Sol.
$$\int \frac{\sin x}{\sin^3 x + \cos^3 x} dx = \int \frac{\frac{\sin x}{\cos^3 x}}{1 + \tan^3 x} dx = \int \frac{\tan x \cdot \sec^2 x}{(\tan x + 1)(1 + \tan^2 x - \tan x)} dx$$

Let $\tan x = t \Rightarrow \sec^2 x \cdot dx = dt$

$$= \int \frac{t}{(t+1)(t^2-t+1)} dt$$

$$= \int \left(\frac{A}{t+1} + \frac{B(2t-1)}{t^2-t+1} + \frac{C}{t^2-t+1} \right) dx$$

$$\Rightarrow A(t^2 - t + 1) + B(2t - 1)(t^2 - t + 1) + C(t + 1) = t$$

$$\Rightarrow t^2(A + 2B) + t(-A + B + C) + A - B + C = 1$$

$$\therefore A + 2B = 0 \quad \dots(1)$$

$$-A + B + C = 1 \quad \dots(2)$$

$$A - B + C = 1 \quad \dots(3)$$

$$\Rightarrow C = \frac{1}{2} \Rightarrow A - B = -\frac{1}{2} \quad \dots(4)$$

$$A + 2B = 0$$

$$A - B = -\frac{1}{2}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



$$\Rightarrow 3B = \frac{1}{2} \Rightarrow B = \frac{1}{6}$$

$$A = -\frac{1}{3}$$

$$I = -\frac{1}{3} \int \frac{dt}{1+t} + \frac{1}{6} \int \frac{2t-1}{t^2-t+1} dt + \frac{1}{2} \int \frac{dt}{t^2-t+1}$$

$$= -\frac{1}{3} \ln|(1+\tan x)| + \frac{1}{6} \ln|\tan^2 x - \tan x + 1| + \frac{1}{2} \cdot \frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{\tan x - \frac{1}{2}}{\frac{\sqrt{3}}{2}} \right)$$

$$= -\frac{1}{3} \ln|1+\tan x| + \frac{1}{6} \ln|\tan^2 x - \tan x + 1| + \frac{1}{\sqrt{3}} \tan^{-1} \left(\frac{2\tan x - 1}{\sqrt{3}} \right) + C$$

$$\alpha = -\frac{1}{3}, \beta = \frac{1}{6}, \gamma = \frac{1}{\sqrt{3}}$$

$$18(\alpha + \beta + \gamma^2) = 18 \left(-\frac{1}{3} + \frac{1}{6} + \frac{1}{3} \right) = 3$$

3. A tangent line L is drawn at the point (2, -4) on the parabola $y^2 = 8x$. If the line L is also tangent to the circle $x^2 + y^2 = a$, then 'a' is equal to _____.

Ans. (2)

Sol. Tangent of $y^2 = 8x$ is $y = mx + \frac{2}{m}$

$$P(2, -4) \Rightarrow -4 = 2m + \frac{2}{m}$$

$$\Rightarrow m + \frac{1}{m} = -2 \Rightarrow m = -1$$

\therefore tangent is $y = -x - 2$

$$\Rightarrow x + y + 2 = 0 \quad \dots(1)$$

(1) is also tangent to $x^2 + y^2 = a$

$$\text{So, } \frac{2}{\sqrt{2}} = \sqrt{a} \Rightarrow \sqrt{a} = \sqrt{2}$$

$$\Rightarrow a = 2$$

4. The number of 4-digit numbers which are neither multiple of 7 nor multiple of 3 is _____

Ans. (5143)

Sol. A = 4 - digit numbers divisible by 3



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



$$A = 1002, 1005, \dots, 9999.$$

$$9999 = 1002 + (n - 1)3$$

$$\Rightarrow (n - 1)3 = 8997 \Rightarrow n = 3000$$

B = 4 - digit numbers divisible by 7

$$B = 1001, 1008, \dots, 9996$$

$$\Rightarrow 9996 = 1001 + (n - 1)7$$

$$\Rightarrow n = 1286$$

$$A \cap B = 1008, 1029, \dots, 9996$$

$$9996 = 1008 + (n - 1)21$$

$$\Rightarrow n = 429$$

So, no divisible by either 3 or 7

$$= 3000 + 1286 - 429 = 3857$$


total 4-digits numbers = 9000

$$\text{required numbers} = 9000 - 3857 = 5143$$

5. If the line $y = mx$ bisects the area enclosed by the lines $x = 0$, $y = 0$, $x = \frac{3}{2}$ and the curve $y = 1 + 4x - x^2$, then $12m$ is equal to _____.

Ans. (26)

Sol.



$$\text{Total area} = \int_0^{3/2} (1 + 4x - x^2) dx$$

$$= x + 2x^2 - \frac{x^3}{3} \Big|_0^{3/2} = \frac{39}{8}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**




$$\& \frac{39}{16} = \frac{1}{2} \cdot \frac{3}{2} \cdot \frac{3}{2} m$$

$$\Rightarrow 3m = \frac{13}{2} \Rightarrow 12m = 26$$

6. Let B be the centre of the circle $x^2 + y^2 - 2x + 4y + 1 = 0$. Let the tangents at two points P and Q on the circle intersect at the point A(3, 1). Then $8 \cdot \left(\frac{\text{area } \Delta APQ}{\text{area } \Delta BPQ} \right)$ is equal to _____.

Ans. (18)

Sol.



$$\tan \theta = \frac{3}{2}$$

$$\frac{\text{Area } \Delta APQ}{\text{Area } \Delta BPQ} = \frac{AR}{RB} = \frac{3 \sin \theta}{2 \cos \theta} = \frac{9}{4}$$

$$8 \left(\frac{\text{Area } \Delta APQ}{\text{Area } \Delta BPQ} \right) = 18$$

7. Let $f(x)$ be a cubic polynomial with $f(1) = -10$, $f(-1) = 6$, and has a local minima at $x = 1$, and $f'(x)$ has a local minima at $x = -1$. Then $f(3)$ is equal to _____.

Ans. (22)

Sol. $F'(x) = a(x - 1)(x + 3)$

$$F''(x) = 6a(x + 1)$$

$$F'(x) = 3a(x + 1)^2 + b$$

$$F'(1) = 0 \Rightarrow b = -12a$$

$$F(x) = a(x + 1)^3 - 12ax + c$$

$$= (x + 1)^3 - 12x - 6$$

$$F(3) = 64 - 36 - 6 = 22$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



8. If $S = \frac{7}{5} + \frac{9}{5^2} + \frac{13}{5^3} + \frac{19}{5^4} + \dots$, then $160S$ is equal to_____.

Ans. (305)

Sol. $S = \frac{7}{5} + \frac{9}{5^2} + \frac{13}{5^3} + \frac{19}{5^4} + \dots$

$$\frac{1}{5}S = \frac{7}{5^2} + \frac{9}{5^3} + \frac{13}{5^4} + \dots$$

On subtracting

$$\frac{4}{5}S = \frac{7}{5} + \frac{2}{5^2} + \frac{4}{5^3} + \frac{6}{5^4} + \dots$$

$$S = \frac{7}{4} + \frac{1}{10} \left(1 + \frac{1}{2} + \frac{3}{5^2} + \dots \right)$$

$$S = \frac{7}{4} + \frac{1}{10} \left(1 - \frac{1}{5} \right)^{-2}$$

$$= \frac{7}{4} + \frac{1}{10} \times \frac{25}{16} = \frac{61}{32}$$

$$\Rightarrow 160S = 5 \times 61 = 305$$

9. If the coefficient of a^7b^8 in the expansion of $(a + 2b + 4ab)^{10}$ is $K \cdot 2^{16}$, then K is equal to_____.

Ans. (315)

Sol. $\frac{10!}{\alpha! \beta! \gamma!} a^\alpha (2b)^\beta \cdot (4ab)^\gamma$

$$\frac{10!}{\alpha! \beta! \gamma!} a^{\alpha+\beta} \cdot b^{\beta+\gamma} \cdot 2^\beta \cdot 4^\gamma$$

$$\alpha + \beta + \gamma = 10 \quad \dots(1)$$

$$\alpha + \gamma = 7 \quad \dots(2)$$

$$\beta + \gamma = 8 \quad \dots(3)$$

$$\text{Equation (2) + (3) - (1)} \Rightarrow \gamma = 5$$

$$\alpha = 2$$

$$\beta = 3$$

$$\text{So, coefficients} = \frac{10!}{2!3!5!} 2^3 \cdot 2^{10}$$

$$= \frac{10 \times 9 \times 8 \times 7 \times 6 \times 5}{2 \times 3 \times 2 \times 5!} \times 2^{13}$$

$$= 315 \times 2^{16} \Rightarrow k = 315$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



10. The number of elements in the set

$$\left\{ A = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix}; a, b, d \in \{-1, 0, 1\} \text{ and } (I-A)^3 = I - A^3 \right\}, \text{ where } I \text{ is } 2 \times 2 \text{ identity matrix, is } \underline{\hspace{2cm}}$$

Ans. (8)

Sol. $(I - A)^3 = I^3 - A^3 - 3A(I - A) = I - A^3$

$$\Rightarrow 3A(I - A) = 0 \text{ or } A^2 = A$$

$$\Rightarrow \begin{bmatrix} a^2 & ab + bd \\ 0 & d^2 \end{bmatrix} = \begin{bmatrix} a & b \\ 0 & d \end{bmatrix}$$

$$\Rightarrow a^2 = a, b(a + d - 1) = 0, d^2 = d$$

$$\text{If } b \neq 0, a + d = 1 \Rightarrow 4 \text{ ways}$$

$$\text{If } b = 0, a = 0, 1 \text{ \& } d = 0, 1 \Rightarrow 4 \text{ ways}$$

$$\Rightarrow \text{Total 8 matrices}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



हो चुकी है ऑफलाइन क्लासरूम की शुरुआत
अपने सपने को करो साकार, कोटा कोचिंग के साथ

Directors of Nucleus Education & Wizard of Mathematics

Now Offline associated with Motion Kota Classroom



Nitin Vijay (NV Sir)
Managing Director
Exp. : 18 yrs



Akhilesh Kanther (AKK Sir)
Exp. : 17 yrs



Vishal Joshi (VJ Sir)
Exp. : 18 yrs



Surendra K. Mishra (SKM Sir)
Exp. : 16 yrs



Gavesh Bhardwaj (GB Sir)
Exp. : 17 yrs

Academic Pillars of JEE MOTION KOTA



Ram Ratan Dwivedi (RRD Sir)
Joint Director
Exp. : 20 yrs



Amit Verma (AV Sir)
Joint Director
Exp. : 16 yrs



Vijay Pratap Singh (VPS Sir)
Vice President
Exp. : 20 yrs



Nikhil Srivastava (NS Sir)
Head JEE Academics
Exp. : 17 yrs



Aatish Agarwal (AA Sir)
Sr. Faculty
Exp. : 17 yrs



Jayant Chittora (JC Sir)
Sr. Faculty
Exp. : 16 yrs



Anurag Garg (AG Sir)
Sr. Faculty
Exp. : 17 yrs



Arjun Gupta (Arjun Sir)
Sr. Faculty
Exp. : 14 yrs



Devki Nandan Pathak (DN Sir)
Sr. Faculty
Exp. : 13 yrs



Avinash Kishore (AVN Sir)
Sr. Faculty
Exp. : 9 yrs



Vipin Sharma (VS Sir)
Sr. Faculty
Exp. : 12 yrs



Durgesh Pandey (Pandey Sir)
Sr. Faculty
Exp. : 8 yrs

Join

JEE DROPPER BATCH

Online + Offline Mode

English & Hindi Medium

Batch Starting from :
22nd Sept. 2021