

**JEE  
MAIN  
4<sup>th</sup>  
Attempt**

**PHYSICS**

**27<sup>th</sup> August 2021 [SHIFT – 1]**

**QUESTION WITH SOLUTION**

**JEE | NEET | Foundation**

**MOTION<sup>®</sup>**

**29900+**  
SELECTIONS SINCE 2007

हो चुकी है ऑफलाइन क्लासरूम की शुरुआत  
अपने सपने को करो साकार, कोटा कोचिंग के साथ

**Directors of Nucleus Education & Wizard of Mathematics**

Now Offline associated with Motion Kota Classroom



**Nitin Vijay (NV Sir)**  
Managing Director  
Exp. : 18 yrs



**Akhilesh Kanther (AKK Sir)**  
Exp. : 17 yrs



**Vishal Joshi (VJ Sir)**  
Exp. : 18 yrs



**Surendra K. Mishra (SKM Sir)**  
Exp. : 16 yrs



**Gavesh Bhardwaj (GB Sir)**  
Exp. : 17 yrs

**Academic Pillars of JEE MOTION KOTA**



**Ram Ratan Dwivedi (RRD Sir)**  
Joint Director  
Exp. : 20 yrs



**Amit Verma (AV Sir)**  
Joint Director  
Exp. : 16 yrs



**Vijay Pratap Singh (VPS Sir)**  
Vice President  
Exp. : 20 yrs



**Nikhil Srivastava (NS Sir)**  
Head JEE Academics  
Exp. : 17 yrs



**Aatish Agarwal (AA Sir)**  
Sr. Faculty  
Exp. : 17 yrs



**Jayant Chittora (JC Sir)**  
Sr. Faculty  
Exp. : 16 yrs



**Anurag Garg (AG Sir)**  
Sr. Faculty  
Exp. : 17 yrs



**Arjun Gupta (Arjun Sir)**  
Sr. Faculty  
Exp. : 14 yrs



**Devki Nandan Pathak (DN Sir)**  
Sr. Faculty  
Exp. : 13 yrs



**Avinash Kishore (AVN Sir)**  
Sr. Faculty  
Exp. : 9 yrs



**Vipin Sharma (VS Sir)**  
Sr. Faculty  
Exp. : 12 yrs



**Durgesh Pandey (Pandey Sir)**  
Sr. Faculty  
Exp. : 8 yrs

Join

**JEE DROPPER BATCH**

Online + Offline Mode

English & Hindi Medium

Batch Starting from :  
**22nd Sept. 2021**

### SECTION – A

1. In a photoelectric experiment, increasing the intensity of incident light :
- (1) increases the frequency of photons incident and increases the K.E. of the ejected electrons
  - (2) increases the number of photons incident and also increases the K.E. of the ejected electrons
  - (3) increases the number of photons incident and the K.E. of the ejected electrons remains unchanged.
  - (4) increases the frequency of photons incident and the K.E. of the ejected electrons remains unchanged.

**Sol. 3**  
 → Increasing intensity means number of incident photons are increased.  
 → Kinetic energy of ejected electrons depends on the frequency of incident photons, not the intensity.

2. Two ions of masses 4 amu and 16 amu have charges +2e and +3e respectively. These ions pass through the region of constant perpendicular magnetic field. The kinetic energy of both ions is same. Then -
- (1) Both ions will be deflected equally
  - (2) no ion will be deflected
  - (3) lighter ion will be deflected more than heavier ion
  - (4) lighter ion will be deflected less than heavier ion


**Sol. 3**

$$r = \frac{P}{qB} = \frac{\sqrt{2mk}}{qB}$$

Given they have same kinetic energy

$$r \propto \frac{\sqrt{m}}{q}$$

$$\frac{r_1}{r_2} = \frac{\sqrt{4}}{2} \times \frac{3}{\sqrt{16}} = \frac{3}{4}$$

$$r_2 = \frac{4r_1}{3} \quad (r_2 \text{ is for heavier ion and } r_1 \text{ is for lighter ion})$$


$$\sin \theta = \frac{d}{R}$$

$\theta \rightarrow$  Deflection

$$\theta \propto \frac{1}{R}$$

(R  $\rightarrow$  radius of path)

$\therefore R_2 > R_1 \Rightarrow \theta_2 < \theta_1$

3. Which of the following is not a dimensionless quantity ?
- (1) Power factor
  - (2) Quality factor
  - (3) Permeability of free space ( $\mu_0$ )
  - (4) Relative magnetic permeability ( $\mu_r$ )



An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



**Sol. 3**

$$[\mu_r] = 1 \text{ as } \mu_r = \frac{\mu}{\mu_m}$$

$$[\text{Power factor } (\cos \phi)] = 1$$

$$\mu_0 = \frac{B_0}{H} \text{ (unit = NA}^{-2}\text{) : Not dimensionless}$$

$$[\mu_0] = [\text{MLT}^{-2}\text{A}^{-2}]$$

$$\text{Quality factor (Q)} = \frac{\text{Energy stored}}{\text{energy dissipated per cycle}}$$

So Q is unitless and dimensionless.

**4.** Electric field in a plane electromagnetic wave is given by

$$E = 50 \sin (500x - 10 \times 10^{10} t) \text{ V/m}$$

The velocity of electromagnetic wave in this medium is -

(Given C = speed of light in vacuum)

- (1)  $\frac{2}{3}C$                       (2) C                      (3)  $\frac{C}{2}$                       (4)  $\frac{3}{2}C$


**Sol. 1**

$$v = \frac{\omega}{k} = \frac{10 \times 10^{10}}{500} = 2 \times 10^8$$

$$v = \frac{2C}{3}$$

**5.** The resultant of these force  $\overline{OP}, \overline{OQ}, \overline{OR}, \overline{OS}$  and  $\overline{OT}$  is approximately \_\_\_\_\_ N.

[Take  $\sqrt{3} = 1.7, \sqrt{2} = 1.4$  Given  $\hat{i}$  and  $\hat{j}$  unit vector along x,y axis]



- (1)  $-1.5\hat{i} - 15.5\hat{j}$       (2)  $9.25\hat{i} + 5\hat{j}$       (3)  $3\hat{i} + 15\hat{j}$       (4)  $2.5\hat{i} - 14.5\hat{j}$




An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



Sol. 2



$$\begin{aligned}\bar{F}_x &= \left( 10 \times \frac{\sqrt{3}}{2} + 20 \left( \frac{1}{2} \right) + 20 \left( \frac{1}{\sqrt{2}} \right) - 15 \left( \frac{1}{\sqrt{2}} \right) - 15 \left( \frac{\sqrt{3}}{2} \right) \right) \hat{i} \\ &= 9.25 \hat{i} \\ \bar{F}_y &= \left( 15 \left( \frac{1}{2} \right) + 20 \left( \frac{\sqrt{3}}{2} \right) + 10 \left( \frac{1}{2} \right) - 15 \left( \frac{1}{\sqrt{2}} \right) - 20 \left( \frac{1}{\sqrt{2}} \right) \right) \hat{j} \\ &= 5 \hat{j}\end{aligned}$$

6. These are  $10^{10}$  radioactive nuclei in a given radioactive element. Its half-life time is 1 minute. How many nuclei will remain after 30 seconds ? ( $\sqrt{2} = 1.414$ )

- (1)  $7 \times 10^9$       (2)  $2 \times 10^{10}$       (3)  $10^5$       (4)  $4 \times 10^{10}$

Sol. 1

$$\frac{N}{N_0} = \left( \frac{1}{2} \right)^{t_{1/2}}$$

$$\frac{N}{10^{10}} = \left( \frac{1}{2} \right)^{\frac{30}{60}}$$

$$\Rightarrow N = 10^{10} \times \left( \frac{1}{2} \right)^{\frac{1}{2}} = \frac{10^{10}}{\sqrt{2}} \approx 7 \times 10^9$$

7. For a transistor in CE mode to be used as an amplifier, it must be operated in -

- (1) Both cut-off and Saturation  
(2) Saturation region only  
(3) Cut-off region only  
(4) The active region only

Sol. 4

Active region of the CE transistor is linear region and is best suited for its use as an amplifier.




An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE


New batch Starting from : **22nd Sept. 2021**



8. A bar magnet is passing through a conducting loop of radius  $R$  with velocity  $v$ . The radius of the bar magnet is such that it just passes through the loop. The induced e.m.f. in the loop can be represented by the approximate curve -



Sol. 4



- When magnet passes through centre region of solenoid, no current/Emf is induced in loop.
- While entering flux increases so negative induced emf
- While leaving flux decreases so positive induced emf.

9. An object is placed beyond the centre of curvature  $C$  of the given concave mirror. If the distance of the object is  $d_1$  from  $C$  and the distance of the image formed is  $d_2$  from  $C$ , the radius of curvature of this mirror is -

(1)  $\frac{2d_1d_2}{d_1 - d_2}$       (2)  $\frac{2d_1d_2}{d_1 + d_2}$       (3)  $\frac{d_1d_2}{d_1 - d_2}$       (4)  $\frac{d_1d_2}{d_1 + d_2}$



An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE


New batch Starting from : **22nd Sept. 2021**



**Sol. 1**

Using newton's formula  
 $(f + d_1)(f - d_2) = f^2$   
 $f^2 + fd_1 - fd_2 - d_1d_2 = f^2$   
 $f = \frac{d_1d_2}{d_1 - d_2}$   
 $\therefore R = \frac{2d_1d_2}{d_1 - d_2}$

**10.** Find the distance of the image from object O, formed by the combination of lenses in the figure –



(1) infinity

(2) 10 cm

(3) 75 cm

(4) 20 cm

**Sol. 3**

$$\frac{1}{V_1} + \frac{1}{30} = \frac{1}{10}$$

$$\frac{1}{V_1} = \frac{2}{30} \Rightarrow V_1 = 15 \text{ cm}$$

$$\frac{1}{V_2} - \frac{1}{10} = -\frac{1}{10}$$

$$\frac{1}{V_2} = 0$$

$$V_2 = \infty$$

$$V_3 = 30 \text{ cm}$$

Distance of the image from object O is = 75 cm

**11.** A huge circular arc of length 4.4 ly subtends an angle '4s' at the centre of the circle. How long it would take for a body to complete 4 revolution if its speed is 8AU per second ?

Given : 1 ly =  $9.46 \times 10^{15}$  m

1 AU =  $1.5 \times 10^{11}$  m

(1)  $4.1 \times 10^8$  s      (2)  $3.5 \times 10^6$  s      (3)  $7.2 \times 10^8$  s      (4)  $4.5 \times 10^{10}$  s

**Sol. 4**

$$R = \frac{\ell}{\theta}$$

$$\text{Time} = \frac{4 \times 2\pi R}{v} = \frac{4 \times 2\pi}{v} \left( \frac{\ell}{\theta} \right)$$

put  $\ell = 4.4 \times 9.46 \times 10^{15}$

$v = 8 \times 1.5 \times 10^{11}$

we get time =  $4.5 \times 10^{10}$  sec



An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



- 12.** If E and H represents the intensity of electric field and magnetising field respectively, then the unit of E/H will be -  
 (1) newton                      (2) ohm                              (3) mho                              (4) joule

**Sol. 2**

$$\text{Unit of } \frac{E}{H} \text{ is } \frac{\text{volt / metre}}{\text{Ampere / metre}} = \frac{\text{volt}}{\text{Ampere}} = \text{ohm}$$

- 13.** A balloon carries a total load of 185 kg at normal pressure and temperature of 27°C. What load will the balloon carry on rising to a height at which the barometric pressure is 45 cm of Hg and the temperature is -7°C. Assuming the volume constant ?  
 (1) 181.46 kg                      (2) 219.07 kg                      (3) 214.15 kg                      (4) 123.54 kg

**Sol. 4**

$$P_m = \rho RT$$

$$\therefore \frac{P_1}{P_2} = \frac{\rho_1 T_1}{\rho_2 T_2}$$

$$\frac{\rho_1}{\rho_2} \Rightarrow \frac{P_1 T_2}{P_2 T_1} = \left( \frac{76}{45} \right) \times \frac{266}{300}$$

$$\frac{\rho_1}{\rho_2} \Rightarrow \frac{M_1}{M_2} = \frac{76 \times 266}{45 \times 300}$$

$$\therefore M_2 \Rightarrow \frac{45 \times 300 \times 185}{76 \times 266} = 123.54 \text{ kg}$$

- 14.** In Millikan's oil drop experiment, what is viscous force acting on an uncharged drop of radius  $2.0 \times 10^{-5}$  m and density  $1.2 \times 10^3 \text{ kgm}^{-3}$  ? Take viscosity of liquid =  $1.8 \times 10^{-5} \text{ Nsm}^{-2}$ .  
 (Neglect buoyancy due to air)  
 (1)  $5.8 \times 10^{-10} \text{ N}$       (2)  $1.8 \times 10^{-10} \text{ N}$       (3)  $3.9 \times 10^{-10} \text{ N}$       (4)  $3.8 \times 10^{-11} \text{ N}$


**Sol. 3**

Viscous force = Weight

$$= \rho \times \left( \frac{4}{3} \pi r^3 \right) g$$

$$= 3.9 \times 10^{-10}$$

- 15.** The variation of displacement with time of a particle executing free simple harmonic motion is shown in the figure.



The potential energy U(x) versus time (t) plot of the particle is correctly shown to figure -  
 An Unmatched Experience of Offline




**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**








**Sol. 2**  
Potential energy is maximum at maximum distance from mean.

**16.** Moment of inertia of a square plate of side  $l$  about the axis passing through one of the corner and perpendicular to the plane of square plate is given by -

- (1)  $MI^2$                       (2)  $\frac{MI^2}{12}$                       (3)  $\frac{MI^2}{6}$                       (4)  $\frac{2}{3}MI^2$

**Sol. 4**  
According to perpendicular axis theorem.




$$I_x + I_y = I_z$$

$$I_z \Rightarrow \frac{ml^2}{3} + \frac{ml^2}{3} = \frac{2ml^2}{3}$$

**17.** A uniform charged disc of radius  $R$  having surface charge density  $\sigma$  is placed in the  $xy$  plane with its center at the origin. Find the electric field intensity along the  $z$ -axis at a distance  $Z$  from origin -

- (1)  $E = \frac{\sigma}{2\epsilon_0} \left( 1 + \frac{Z}{(Z^2 + R^2)^{1/2}} \right)$                       (2)  $E = \frac{\sigma}{2\epsilon_0} \left( 1 - \frac{Z}{(Z^2 + R^2)^{1/2}} \right)$   
 (3)  $E = \frac{2\epsilon_0}{\sigma} \left( \frac{1}{(Z^2 + R^2)^{1/2}} + Z \right)$                       (4)  $E = \frac{\sigma}{2\epsilon_0} \left( \frac{1}{(Z^2 + R^2)} + \frac{1}{Z^2} \right)$

**Sol. 2**  
Consider a small ring of radius  $r$  and thickness  $dr$  on disc.



An Unmatched Experience of Offline



**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



area of elemental ring on disc

$$dA = 2\pi r dr$$

charge on this ring  $dq = \sigma dA$

$$dE_z = \frac{k dq z}{(z^2 + r^2)^{3/2}}$$

$$E = \int_0^R dE_z = \frac{\sigma}{2 \epsilon_0} \left[ 1 - \frac{z}{\sqrt{R^2 + z^2}} \right]$$

- 18.** An ideal gas is expanding such that  $PT^3 = \text{constant}$ . The coefficient of volume expansion of the gas is -

(1)  $\frac{4}{T}$                       (2)  $\frac{1}{T}$                       (3)  $\frac{2}{T}$                       (4)  $\frac{3}{T}$

**Sol. 1**

$$PT^3 = \text{constant}$$

$$\left( \frac{nRT}{V} \right) T^3 = \text{constant}$$

$$T^4 V^{-1} = \text{constant}$$

$$T^4 = kV$$

$$\Rightarrow 4 \frac{\Delta T}{T} = \frac{\Delta V}{V} \quad \dots(1)$$

$$\Delta V = V\gamma \Delta T \quad \dots(2)$$

comparing (1) and (2)

we get

$$\gamma = \frac{4}{T}$$

- 19.** Five identical cells each of internal resistance  $1\Omega$  and emf  $5V$  are connected in series and in parallel with an external resistance 'R'. For what value of 'R', current in series and parallel combination will remain the same ?

(1)  $10\Omega$                       (2)  $25\Omega$                       (3)  $1\Omega$                       (4)  $5\Omega$

**Sol. 3**

$$i_1 = \frac{25}{5 + R}$$

$$i_2 = \frac{5}{R + \frac{1}{5}}$$

$$i_1 = i_2 \Rightarrow 5 \left( R + \frac{1}{5} \right) = 5 + R$$

$$4R = 4$$

$$R = 1\Omega$$




An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**




20. calculate the amount of charge on capacitor of  $4\mu\text{F}$ . The internal resistance of battery is  $1\Omega$ .



- (1) zero                      (2)  $4\mu\text{C}$                       (3)  $16\mu\text{C}$                       (4)  $8\mu\text{C}$

**Sol. 4**

On simplifying circuit we get



No current in upper wire.


$$\therefore V_{AB} = \frac{5}{4+1} \times 4 = 4 \text{ v}$$

$$\therefore \theta = (C_{eq})v$$

$$\Rightarrow 2 \times 4 = 8\mu\text{C}$$

### Section - B

1. A circular is arranged as shown in figure. The output voltage  $V_0$  is equal to \_\_\_\_\_ V.



An Unmatched Experience of Offline


**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



**Sol. 5**

As diodes  $D_1$  and  $D_2$  are in forward bias, so they acted as negligible resistance  
 $\Rightarrow$  Input voltage become zero



$\Rightarrow$  Input current is zero  
 $\Rightarrow$  Output current is zero  
 $\Rightarrow V_0 = 5$  volt

- 2.** First, a set of  $n$  equal resistors of  $10\Omega$  each are connected in series to a battery of emf  $20V$  and internal resistance  $10\Omega$ . A current  $I$  is observed to flow. Then, the  $n$  resistors are connected in parallel to the same battery. It is observed that the current is increased 20 times, then the value of  $n$  is \_\_\_\_\_.

**Sol. 20**

In series

$$R_{eq} = nR = 10n$$

$$i_s = \frac{20}{10 + 10n} = \frac{2}{1 + n}$$

in parallel

$$R_{eq} = \frac{10}{n}$$

$$i_p = \frac{20}{\frac{10}{n} + 10} = \frac{2n}{1 + n}$$

$$\frac{i_p}{i_s} = 20$$

$$\frac{\left(\frac{2n}{1+n}\right)}{\left(\frac{2}{1+n}\right)} = 20$$

$$n = 20$$

- 3.** Two cars X and Y are approaching each other with velocities  $36 \text{ km/h}$  and  $72 \text{ km/h}$  respectively. The frequency of a whistle sound as emitted by a passenger in car X, heard by the passenger in car Y is  $1320 \text{ Hz}$ . If the velocity of sound in air is  $340 \text{ m/s}$ , the actual frequency of the whistle sound produced is \_\_\_\_\_ Hz.




An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



Sol. 1210



$$V_x = 36 \text{ km/hr} = 10 \text{ m/s}$$


$$V_y = 72 \text{ km/hr} = 20 \text{ m/s}$$

by doppler's effect


$$F' = F_0 \left( \frac{V \pm V_0}{V \pm V_s} \right)$$

$$1320 = F_0 \left( \frac{340 + 20}{340 - 10} \right) \Rightarrow F_0 = 1210 \text{ Hz}$$

4. A rod CD of thermal resistance  $10.0 \text{ KW}^{-1}$  is joined at the middle of an identical rod AB as shown in figure. The ends A,B and D are maintained at  $200^\circ\text{C}$ ,  $100^\circ\text{C}$  and  $125^\circ\text{C}$  respectively. The heat current in CD is P watt. The value of P is \_\_\_\_\_.



Sol. 2



Rods are identical so

$$R_{AB} = R_{CD} = 10 \text{ Kw}^{-1}$$

C is mid-point of AB, so

$$R_{AC} = R_{CB} = 5 \text{ Kw}^{-1}$$

at point C

$$\frac{200 - T}{5} = \frac{T - 125}{10} + \frac{T - 100}{5}$$

$$2(200 - T) = T - 125 + 2(T - 100)$$

$$400 - 2T = T - 125 + 2T - 200$$

$$T = \frac{725}{5} = 145^\circ\text{C}$$

$$I_h = \frac{145 - 125}{10} \text{ w} = \frac{20}{10} \text{ w}$$

$$I_h = 2\text{w}$$



An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



5. A uniform conducting wire of length is  $24a$ , and resistance  $R$  is wound up as a current carrying coil in the shape of an equilateral triangle of side 'a' and then in the form of a square of side 'a'. The coil is connected to a voltage source  $V_0$ . The ratio of magnetic moment of the coil in case of equilateral triangle to that for square is  $1 : \sqrt{y}$  where  $y$  is \_\_\_\_\_.

**Sol. 3**

$$\text{In triangle shape } N_t = \frac{24a}{3a} = 8$$

$$\text{In square } N_s = \frac{24a}{4a} = 6$$

$$\frac{M_t}{M_s} = \frac{N_t I A_t}{N_s I A_s} \quad [I \text{ will be same in both}]$$

$$\frac{8 \times \frac{\sqrt{3}}{4} \times a^2}{6 \times a^2}$$

$$\frac{M_t}{M_s} = \frac{1}{\sqrt{3}}$$

$$y = 3$$

6. The alternating current is given by

$$i = \left\{ \sqrt{42} \sin\left(\frac{2\pi}{T} t\right) + 10 \right\} \text{ A}$$

The r.m.s. value of this current is \_\_\_\_\_ A.

**Sol. 11**

$$f_{\text{rms}}^2 = f_{1\text{rms}}^2 + f_{2\text{rms}}^2$$

$$\left(\frac{\sqrt{42}}{\sqrt{2}}\right)^2 + 10^2$$

$$= 121 \Rightarrow f_{\text{rms}} = 11 \text{ A}$$

7. Two persons A and B perform same amount of work in moving a body through a certain distance  $d$  with application of force acting at angles  $45^\circ$  and  $60^\circ$  with the direction of displacement respectively. The ratio of force applied by person A to the force applied by person B is  $\frac{1}{\sqrt{x}}$ . The value of  $x$  is \_\_\_\_\_.

**Sol. 2**

$$\text{Given } W_A = W_B$$

$$F_A d \cos 45^\circ = F_B d \cos 60^\circ$$

$$F_A \times \frac{1}{\sqrt{2}} = F_B \times \frac{1}{2}$$

$$\frac{F_A}{F_B} = \frac{\sqrt{2}}{2} = \frac{1}{\sqrt{2}}$$

$$x = 2$$



An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



8. If the velocity of a body related to displacement  $x$  is given by  $v = \sqrt{5000 + 24x}$  m/s, then the acceleration of the body is \_\_\_\_\_ m/s<sup>2</sup>

Sol. 12

$$V = \sqrt{5000 + 24x}$$

$$\frac{dV}{dx} = \frac{1}{2\sqrt{5000 + 24x}} \times 24 = \frac{12}{\sqrt{5000 + 24x}}$$

$$\text{now } a = V \frac{dV}{dx}$$

$$= \sqrt{5000 + 24x} \times \frac{12}{\sqrt{5000 + 24x}}$$

$$a = 12 \text{ m/s}^2$$

9. A transmitting antenna has a height of 320 m and that of receiving antenna is 2000 m. The maximum distance between them for satisfactory communication in line of sight mode is 'd'. The value of 'd' is \_\_\_\_\_ km.


Sol. 224

$$d_m = \sqrt{2Rh_T} + \sqrt{2Rh_R}$$

$$d_m = \left( \sqrt{2 \times 6400 \times 10^3 \times 320} + \sqrt{2 \times 6400 \times 10^3 \times 2000} \right) \text{m}$$

$$d_m = 224 \text{ km}$$

10. A body of mass (2M) splits into four masses (m, M - m, m, M - m), which are rearranged to form a square as shown in the figure. The ratio of  $\frac{M}{m}$  for which, the gravitational potential energy of the system becomes maximum is  $x : 1$ . The value of  $x$  is \_\_\_\_\_.



Sol. 2

Energy is maximum when mass is split equally so  $\frac{M}{m} = 2$



An Unmatched Experience of Offline

**KOTA CLASSROOM** For JEE

New batch Starting from : **22nd Sept. 2021**



हो चुकी है ऑफलाइन क्लासरूम की शुरुआत  
अपने सपने को करो साकार, कोटा कोविंग के साथ

**Directors of Nucleus Education & Wizard of Mathematics**

Now Offline associated with Motion Kota Classroom



**Nitin Vijay (NV Sir)**  
Managing Director  
Exp. : 18 yrs



**Akhilesh Kanther (AKK Sir)**  
Exp. : 17 yrs



**Vishal Joshi (VJ Sir)**  
Exp. : 18 yrs



**Surendra K. Mishra (SKM Sir)**  
Exp. : 16 yrs



**Gavesh Bhardwaj (GB Sir)**  
Exp. : 17 yrs

## Academic Pillars of JEE MOTION KOTA



**Ram Ratan Dwivedi (RRD Sir)**  
Joint Director  
Exp. : 20 yrs



**Amit Verma (AV Sir)**  
Joint Director  
Exp. : 16 yrs



**Vijay Pratap Singh (VPS Sir)**  
Vice President  
Exp. : 20 yrs



**Nikhil Srivastava (NS Sir)**  
Head JEE Academics  
Exp. : 17 yrs



**Aatish Agarwal (AA Sir)**  
Sr. Faculty  
Exp. : 17 yrs



**Jayant Chittora (JC Sir)**  
Sr. Faculty  
Exp. : 16 yrs



**Anurag Garg (AG Sir)**  
Sr. Faculty  
Exp. : 17 yrs



**Arjun Gupta (Arjun Sir)**  
Sr. Faculty  
Exp. : 14 yrs



**Devki Nandan Pathak (DN Sir)**  
Sr. Faculty  
Exp. : 13 yrs



**Avinash Kishore (AVN Sir)**  
Sr. Faculty  
Exp. : 9 yrs



**Vipin Sharma (VS Sir)**  
Sr. Faculty  
Exp. : 12 yrs



**Durgesh Pandey (Pandey Sir)**  
Sr. Faculty  
Exp. : 8 yrs

Join

**JEE DROPPER BATCH**

Online + Offline Mode

English & Hindi Medium

Batch Starting from :  
**22nd Sept. 2021**