

**JEE
MAIN
4th
Attempt**

PHYSICS

27th August 2021 [SHIFT – 2]

QUESTION WITH SOLUTION

JEE | NEET | Foundation

MOTION[®]

29900+
SELECTIONS SINCE 2007

हो चुकी है ऑफलाइन क्लासरूम की शुरुआत
अपने सपने को करो साकार, कोटा कोचिंग के साथ

Directors of Nucleus Education & Wizard of Mathematics

Now Offline associated with Motion Kota Classroom



Nitin Vijay (NV Sir)
Managing Director
Exp. : 18 yrs



Akhilesh Kanther (AKK Sir)
Exp. : 17 yrs



Vishal Joshi (VJ Sir)
Exp. : 18 yrs



Surendra K. Mishra (SKM Sir)
Exp. : 16 yrs



Gavesh Bhardwaj (GB Sir)
Exp. : 17 yrs

Academic Pillars of JEE MOTION KOTA



Ram Ratan Dwivedi (RRD Sir)
Joint Director
Exp. : 20 yrs



Amit Verma (AV Sir)
Joint Director
Exp. : 16 yrs



Vijay Pratap Singh (VPS Sir)
Vice President
Exp. : 20 yrs



Nikhil Srivastava (NS Sir)
Head JEE Academics
Exp. : 17 yrs



Aatish Agarwal (AA Sir)
Sr. Faculty
Exp. : 17 yrs



Jayant Chittora (JC Sir)
Sr. Faculty
Exp. : 16 yrs



Anurag Garg (AG Sir)
Sr. Faculty
Exp. : 17 yrs



Arjun Gupta (Arjun Sir)
Sr. Faculty
Exp. : 14 yrs



Devki Nandan Pathak (DN Sir)
Sr. Faculty
Exp. : 13 yrs



Avinash Kishore (AVN Sir)
Sr. Faculty
Exp. : 9 yrs



Vipin Sharma (VS Sir)
Sr. Faculty
Exp. : 12 yrs



Durgesh Pandey (Pandey Sir)
Sr. Faculty
Exp. : 8 yrs

Join

JEE DROPPER BATCH


Online + Offline Mode

English & Hindi Medium

Batch Starting from :
22nd Sept. 2021

SECTION - A

1. Curved surfaces of a plano-convex lens of refractive index μ_1 and a plano-concave lens of refractive index μ_2 have equal radius of curvature as shown in figure. Find the ratio of radius of curvature to the focal length of the combined lens.



- (1) $\frac{1}{\mu_1 - \mu_2}$ (2) $\mu_1 - \mu_2$ (3) $\mu_2 - \mu_1$ (4) $\frac{1}{\mu_2 - \mu_1}$

Sol. 2



$$\frac{1}{f_1} = (\mu_1 - 1) \left(\frac{1}{R} \right)$$

$$\frac{1}{f_2} = (\mu_2 - 1) \left(-\frac{1}{R} \right)$$

$$\frac{1}{f_1} + \frac{1}{f_2} = \frac{1}{f_{eq}} = \frac{(\mu_1 - 1) - (\mu_2 - 1)}{R}$$

$$\frac{1}{f_{eq}} = \frac{(\mu_1 - \mu_2)}{R}$$

$$\frac{R}{f_{eq}} = (\mu_1 - \mu_2)$$

2. Two discs have moments of inertia I_1 and I_2 about their respective axes perpendicular to the plane and passing through the centre. They are rotating with angular speeds, ω_1 and ω_2 respectively and are brought into contact face to face with their axes of rotation coaxial. The loss in kinetic energy of the system in the process is given by :

- (1) $\frac{I_1 I_2}{(I_1 + I_2)} (\omega_1 - \omega_2)^2$ (2) $\frac{I_1 I_2}{2(I_1 + I_2)} (\omega_1 - \omega_2)^2$ (3) $\frac{(I_1 - I_2)^2 \omega_1 \omega_2}{2(I_1 + I_2)}$ (4) $\frac{(\omega_1 - \omega_2)^2}{2(I_1 + I_2)}$

Sol. 2

From conservation of angular momentum we get

$$I_1 \omega_1 + I_2 \omega_2 = (I_1 + I_2) \omega$$

$$\omega = \frac{I_1 \omega_1 + I_2 \omega_2}{I_1 + I_2}$$

$$K_i = \frac{1}{2} I_1 \omega_1^2 + \frac{1}{2} I_2 \omega_2^2$$

$$K_f = \frac{1}{2} (I_1 + I_2) \omega^2$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



$$k_i - k_f = \frac{1}{2} \left[I_1 \omega_1^2 + I_2 \omega_2^2 - \frac{(I_1 \omega_1 + I_2 \omega_2)^2}{I_1 + I_2} \right]$$

Solving above we get

$$k_i - k_f = \frac{1}{2} \left(\frac{I_1 I_2}{I_1 + I_2} \right) (\omega_1 - \omega_2)^2$$

3. The height of Victoria falls is 63m. What is the difference in temperature of water at the top and at the bottom of fall ?

[Given 1 cal = 4.2J and specific heat of water = 1 cal g⁻¹ °C⁻¹]

- (1) 14.76°C (2) 1.476°C (3) 0.147°C (4) 0.014°C

Sol. 3

Change in P.E. = Heat energy


$$mgh = mS\Delta T$$

$$\Delta T = \frac{gh}{S}$$

$$= \frac{10 \times 63}{4200 \text{ J/kgC}}$$

$$= 0.147^\circ\text{C}$$

4. The boxes of masses 2 kg and 8 kg are connected by a massless string passing over smooth pulleys. Calculate the time taken by box of mass 8 kg to strike the ground starting from rest. (use g = 10m/s²)



- (1) 0.25s (2) 0.34s (3) 0.2s (4) 0.4s

Sol. 4




An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**





$$(m_1g - 2T) = m_1a \dots (1)$$

$$T - m_2g = m_2(2a)$$

$$2T - 2m_2g = 4m_2a \dots (2)$$

$$m_1g - 2m_2g = (m_1 + 4m_2)a$$

$$a = \frac{(8 - 4)g}{(8 + 8)} = \frac{4}{16}g = \frac{g}{4}$$

$$a = \frac{10}{4} \text{ m/s}^2$$

$$S = \frac{1}{2}at^2$$

$$\frac{0.2 \times 2 \times 4}{10} = t^2$$

$$t = 0.4 \text{ sec}$$

5. A coaxial cable consists of an inner wire of radius 'a' surrounded by an outer shell of inner and outer radii 'b' and 'c' respectively. The inner wire carries an electric current i_0 , which is distributed uniformly across cross-sectional area. The outer shell carries an equal current in opposite direction and distributed uniformly. What will be the ratio of the magnetic field at a distance x from the axis when (i) $x < a$ and (ii) $a < x < b$?


(1) $\frac{b^2 - a^2}{x^2}$

(2) $\frac{a^2}{x^2}$

(3) $\frac{x^2}{b^2 - a^2}$

(4) $\frac{x^2}{a^2}$

Sol. 4



When $x < a$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



$$B_1(2\pi x) = \mu_0 \left(\frac{i_0}{\pi a^2} \right) \pi x^2$$

$$B(2\pi x) = \frac{\mu_0 i_0 x^2}{a^2}$$

$$B_1 = \frac{\mu_0 i_0 x}{2\pi a^2} \quad \dots (a)$$


when $a < x < b$

$$B_2(2\pi x) = \mu_0 i_0$$

$$B_2 = \frac{\mu_0 i_0}{2\pi x} \quad \dots (2)$$

$$\frac{B_1}{B_2} = \frac{\mu_0 i_0 \frac{x}{2\pi a^2}}{\frac{\mu_0 i_0}{2\pi x}} = \frac{x^2}{a^2}$$

6. The colour coding on a carbon resistor is shown in the given figure. The resistance value of the given resistor is :



- (1) $(5700 \pm 285)\Omega$ (2) $(5700 \pm 375)\Omega$ (3) $(7500 \pm 375)\Omega$ (4) $(7500 \pm 750)\Omega$

Sol. 3


$$R = 75 \times 10^2 \pm 5\% \text{ of } 7500$$

$$R = (7500 \pm 375)\Omega$$

7. A mass of 50kg is placed at the centre of a uniform spherical shell of mass 100 kg and radius 50m. If the gravitational potential at a point, 25m from the centre is V kg/m. The value of V is :

- (1) $-20G$ (2) $+2G$ (3) $-4G$ (4) $-60G$

Sol. 3



$$\begin{aligned} V_A &= \left[-\frac{GM_1}{r} - \frac{GM_2}{R} \right] \\ &= \left[-\frac{50}{25}G - \frac{100}{50}G \right] \\ &= -4G \end{aligned}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



8. Match List-I with List- II

List – I

- (a) R_H (Rydberg constant)
- (b) h (Planck's constnat)
- (c) μ_B (Magnetic field energy density)
- (d) η (Coefficient of viscocity)

List – II

- (i) $\text{kg m}^{-1}\text{s}^{-1}$
- (ii) $\text{kg m}^2\text{s}^{-1}$
- (iii) m^{-1}
- (iv) $\text{kg m}^{-1}\text{s}^{-2}$

Choose the most appropriate answer from the options given below :

- (1) (a)-(iii), (b)-(ii), (c)-(i),(d)-(iv)
- (2) (a)-(iii), (b)-(ii), (c)-(iv),(d)-(i)
- (3) (a)-(ii), (b)-(iii), (c)-(iv),(d)-(i)
- (4) (a)-(iv), (b)-(ii), (c)-(i),(d)-(iii)

Sol. 2

SI unit of Rydberg const. = m^{-1}
 SI unit of Plank's const. = $\text{kg m}^2\text{s}^{-1}$
 SI unit of Magnetic field energy density = $\text{kg m}^{-1}\text{s}^{-2}$
 SI unit of coeff. of viscosity = $\text{kg m}^{-1}\text{s}^{-1}$

9. If the rms speed of oxygen molecules at 0°C is 160 m/s, find the rms speed of hydrogen molecules at 0°C .

- (1) 640m/s
- (2) 80m/s
- (3) 332m/s
- (4) 40m/s

Sol. 1

$$V_{\max} = \sqrt{\frac{3KT}{M}}$$


$$\frac{(V_{\text{rms}})_{\text{O}_2}}{(V_{\text{rms}})_{\text{H}_2}} = \sqrt{\frac{M_{\text{H}_2}}{M_{\text{O}_2}}} = \sqrt{\frac{2}{32}}$$

$$(V_{\text{rms}})_{\text{H}_2} = 4 \times (V_{\text{rms}})_{\text{O}_2}$$

$$= 4 \times 160$$

$$= 640 \text{ m/s}$$

10. Three capacitors $C_1 = 2\mu\text{F}$, $C_2 = 6\mu\text{F}$, and $C_3 = 12\mu\text{F}$ are connected as shown in figure. Find the ratio of the charges on capacitor C_1 , C_2 and C_3 respectively :



- (1) 2:1:1
- (2) 2:3:3
- (3) 3:4:4
- (4) 1:2:2

Sol. 4




An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**





$$(V_D - V)C_2 + (V_D - 0)C_3 = 0$$

$$(V_D - V)6 + (V_D - 0)12 = 0$$

$$V_D - V + 2V_D = 0$$

$$V_D = \frac{V}{3}$$

$$q_2 = (V - V_D)C_2 = \left(V - \frac{V}{3}\right)(6\mu\text{F})$$

$$q_2 = (4V)\mu\text{F}$$

$$q_3 = (V_D - 0)C_3 = \frac{V}{3} \times 12\mu\text{F} = 4V\mu\text{F}$$

$$q_1 = (V - 0)C_1 = V(2\mu\text{F})$$

$$q_1 : q_2 : q_3 = 2 : 4 : 4$$

$$q_1 : q_2 : q_3 = 1 : 2 : 2$$

- 11.** For a transistor α and β are given as $\alpha = \frac{I_C}{I_E}$ and $\beta = \frac{I_C}{I_B}$. Then the correct relation between α and β will be:

(1) $\alpha = \frac{1-\beta}{\beta}$

(2) $\alpha\beta = 1$

(3) $\beta = \frac{\alpha}{1-\alpha}$

(4) $\alpha = \frac{\beta}{1-\beta}$

Sol. 3

$$\alpha = \frac{I_C}{I_E}, \beta = \frac{I_C}{I_B}; I_E = I_C + I_B$$

$$\alpha = \frac{I_C}{I_C + I_B} = \frac{I_C / I_B}{\frac{I_C}{I_B} + 1} = \frac{\beta}{\beta + 1}$$

$$1 + \frac{1}{\beta} = \frac{1}{\alpha}$$

$$\frac{1}{\beta} = \frac{1-\alpha}{\alpha}$$

$$\beta = \frac{\alpha}{1-\alpha}$$




An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**




- 12.** A constant magnetic field of 1T is applied in the $x > 0$ region. A metallic circular ring of radius 1m is moving with a constant velocity of 1m/s along the x-axis. At $t=0$ s, the centre O of the ring is at $x = -1$ m. What will be the value of the induced emf in the ring at $t = 1$ s ? (Assume the velocity of ring does not change.)



- Sol. 1**
- (1) 2 V (2) $2\pi V$ (3) 0 V (4) 1 V
- Emf = BLV
 = 1.(2R).1
 = 2V

- 13.** Water drops are falling, from a nozzle of a shower onto the floor, from a height of 9.8m. The drops fall at a regular interval of time. When the first drop strikes the floor, at the instant, the third drop begins to fall. Locate the position of second drop from the floor when the first drop strikes the floor.

- (1) 4.18m (2) 2.94m (3) 2.45m (4) 7.35m
- Sol. 4**



$$H = \frac{1}{2}gt^2$$

$$\frac{9.8 \times 2}{9.8} = t^2$$

$$t = \sqrt{2} \text{ sec}$$

Δt : time interval between drops

$$h = \frac{1}{2}g(\sqrt{2} - \Delta t)^2$$

$$0 = \frac{1}{2}g(\sqrt{2} - 2\Delta t)^2$$

$$\Delta t = \frac{1}{\sqrt{2}}$$

$$h = \frac{1}{2}g\left(\sqrt{2} - \frac{1}{\sqrt{2}}\right)^2 = \frac{1}{2} \times 9.8 \times \frac{1}{2} = \frac{9.8}{4} = 2.45\text{m}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



$$H - h = 9.8 - 2.45 \\ = 7.35\text{m}$$

- 14.** An antenna is mounted on a 400m tall building. What will be the wavelength of signal that can be radiated effectively by the transmission tower upto a range of 44 km?
 (1) 605m (2) 75.6m (3) 37.8m (4) 302m

Sol. 1

h : height of antenna
 λ : wavelength of signal
 $h < \lambda$
 $\lambda > h$
 $\lambda > 400\text{m}$

- 15.** For full scale deflection of total 50 divisions, 50mV voltage is required in galvanometer. The resistance of galvanometer if its current sensitivity is 2 div/ mA will be :
 (1) 2 Ω (2) 5 Ω (3) 1 Ω (4) 4 Ω

Sol. 1

$$I_{\max} = \frac{50}{2} = 25\text{mA}$$

$$R = \frac{V}{I} = \frac{50\text{mV}}{25\text{mA}} = 2\Omega$$

- 16.** A player kicks a football with an initial speed of 25ms^{-1} at an angle of 45° from the ground. What are the maximum height and the time taken by the football to reach at the highest point during motion ? (Take $g = 10 \text{ ms}^{-2}$)
 (1) $h_{\max} = 10 \text{ m}$ $T = 2.5 \text{ s}$ (2) $h_{\max} = 15.625\text{m}$ $T = 3.54\text{s}$
 (3) $h_{\max} = 15.625\text{m}$ $T = 1.77\text{s}$ (4) $h_{\max} = 3.54\text{m}$ $T = 0.125\text{s}$

Sol. 3

$$H = \frac{U^2 \sin^2 \theta}{2g}$$

$$= \frac{(25)^2 \cdot (\sin 45)^2}{2 \times 10}$$

$$= 15.625\text{m}$$

$$T = \frac{U \sin \theta}{g}$$

$$= \frac{25 \times \sin 45^\circ}{10}$$

$$= 2.5 \times 0.7$$

$$= 1.77\text{s}$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



- 17.** A monochromatic neon lamp with wavelength of 670.5 nm illuminates a photo-sensitive material which has a stopping voltage of 0.48V. What will be the stopping voltage if the sources light is changed with another source of wavelength of 474.6nm?

(1) 0.96V (2) 1.5V (3) 1.25V (4) 0.24V

Sol. 3

$$kE_{\max} = \frac{hc}{\lambda_1} + \phi$$

$$\text{or } eV_0 = \frac{hc}{\lambda_1} + \phi$$

$$\text{when } \lambda_1 = 670.5\text{nm}; \quad V_0 = 0.48$$

$$\text{when } \lambda_1 = 474.6\text{nm}; \quad V_0 = ?$$

$$\text{So, } e(0.48) = \frac{1240}{670.5} + \phi \quad \dots(1)$$

$$e(V_0) = \frac{1240}{474.6} + \phi \quad \dots(2)$$

$$(2) - (1)$$

$$e(V_0 - 0.48) = 1240 \left(\frac{1}{474.6} - \frac{1}{670.5} \right) eV$$

$$V_0 = 0.48 + 1240 \left(\frac{670.5 - 474.6}{474.6 \times 670.5} \right) \text{Volt}$$

$$V_0 = 0.48 + 0.76$$

$$V_0 = 1.24V \approx 1.25V$$

- 18.** The light waves from two coherent sources have same intensity $I_1 = I_2 = I_0$. In interference pattern the intensity of light at minima is zero. What will be the intensity of light at maxima?

(1) $5 I_0$ (2) $4 I_0$ (3) $2 I_0$ (4) I_0

Sol. 2

$$I_{\max} = (\sqrt{I_1} + \sqrt{I_2})^2$$

$$4I_0$$

- 19.** If force (F), length (L) and time (T) are taken as the fundamental quantities. Then what will be the dimension of density :

(1) $[FL^{-4}T^2]$ (2) $[FL^{-3}T^3]$ (3) $[FL^{-3}T^2]$ (4) $[FL^{-5}T^2]$

Sol. 1

$$\text{Density} = [F^a L^b T^c]$$

$$[ML^{-3}] = [M^a L^a T^{-2a} L^b T^c]$$

$$[M^1 L^{-3}] = [M^a L^{a+b} T^{-2a+c}]$$

$$a = 1 ; \quad a + b = -3 ; \quad -2a + c = 0$$

$$1 + b = -3 \quad c = 2a$$

$$b = -4 \quad c = 2$$

$$\text{So, density} = [F^1 L^{-4} T^2]$$




An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



20. Figure shows a rod AB, which is bent in a 120° circular arc of radius R. A charge $(-Q)$ is uniformly distributed over rod AB. What is the electric field \vec{E} at the centre of curvature O?



- (1) $\frac{3\sqrt{3}Q}{8\pi\epsilon_0 R^2}(\hat{i})$ (2) $\frac{3\sqrt{3}Q}{8\pi^2\epsilon_0 R^2}(\hat{i})$ (3) $\frac{3\sqrt{3}Q}{16\pi^2\epsilon_0 R^2}(\hat{i})$ (4) $\frac{3\sqrt{3}Q}{8\pi^2\epsilon_0 R^2}(-\hat{i})$

Sol. 2

$$\epsilon = \frac{2k\lambda}{R} \sin\left(\frac{\theta}{2}\right)(-\hat{i})$$

$$\lambda = \left(\frac{-Q}{R\theta}\right) = \left(\frac{-Q}{R \cdot \frac{2\pi}{3}}\right)$$

$$\lambda = \frac{-3Q}{2\pi R}$$

$$\epsilon = \frac{2k}{R} \cdot \frac{-3Q}{2\pi R} \cdot \sin(60^\circ)(-\hat{i})$$

$$\epsilon = \frac{3\sqrt{3}Q}{8\pi^2\epsilon_0 R^2}(\hat{i})$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



Section B

1. A plane electromagnetic wave with frequency of 30 MHz travels in free space. At particular point in space and time, electric field is 6 V/m. The magnetic field at this point will be $x \times 10^{-8}$ T. The value of x is ____.

Sol. 2

$$|B| = \frac{|E|}{c} = \frac{6}{3 \times 10^8}$$

$$= 2 \times 10^{-8} \text{ T}$$

$$\therefore x = 2$$

2. A heat engine operates between a cold reservoir at temperature $T_2 = 400$ K and a hot reservoir at temperature T_1 . It takes 300 J of heat from the hot reservoir and delivers 240 J of heat to the cold reservoir in a cycle. The minimum temperature of the hot reservoir has to be ____ K.

Sol. 500

$$Q_{in} = 300 \text{ J} ; Q_{out} = 240 \text{ J}$$

$$\text{Work done} = Q_{in} - Q_{out} = 300 - 240 = 60 \text{ J}$$

$$\text{Efficiency} = \frac{W}{Q_{in}} = \frac{60}{300} = \frac{1}{5}$$

$$\text{efficiency} = 1 - \frac{T_2}{T_1}$$

$$\frac{1}{5} = 1 - \frac{400}{T_1} \Rightarrow \frac{400}{T_1} = \frac{4}{5}$$

$$T_1 = 500 \text{ k}$$


3. X different wave lengths may be observed in the spectrum from a hydrogen sample if the atoms are excited to states with principal quantum number $n = 6$? The value of X is ____.

Sol. 15

$$\text{No. of different wavelengths} = \frac{n(n-1)}{2}$$

$$= \frac{6 \times (6-1)}{2} = \frac{6 \times 5}{2} = 15$$

4. A bullet of 10g, moving with velocity v , collides head-on with the stationary bob of a pendulum and recoils with velocity 100 m/s. The length of the pendulum is 0.5m and mass of the bob is 1 kg. The minimum value of $v =$ ____ m/s so that the pendulum describes a circle. (Assume the string to be inextensible and $g = 10 \text{ m/s}^2$)



Sol. 400




An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**





$$V' = \sqrt{5gR} = \sqrt{5 \times 10 \times 0.5}$$


$$V' = 5 \text{ m/s}$$

$$m_1 V = m_2 \times 5 - m_1 \times 100$$

$$\frac{10}{1000} \times V = 5 - \frac{10}{1000} \times 100$$

$$V = 400 \text{ m/s}$$

5. A zener diode of power rating 2W is to be used as a voltage regulator. If the zener diode has a breakdown of 10V and it has to regulate voltage fluctuated between 6 V and 14 V, the value of R_s for safe operation should be _____ Ω .



Sol. 20

When unregulated voltage is 14 V voltage across zener diode must be 10 V So potential difference across resistor $\Delta V_{R_s} = 4V$

and $P_{\text{zener}} = 2W$

$$VI = 2$$

$$I = \frac{2}{10} = 0.2A$$

$$\Delta V_{R_s} = IR_s$$

$$4 \times 0.2R_s \Rightarrow R_s = \frac{40}{2} = 20\Omega$$

6. Two simple harmonic motion, are represented by the equations

$$y_1 = 10 \sin\left(3\pi t + \frac{\pi}{3}\right)$$

$$y_2 = 5(\sin 3\pi t + \sqrt{3} \cos 3\pi t)$$

Ratio of amplitude of y_1 to $y_2 = x : 1$. The value of x is _____.

Sol. 1

$$y_1 = 10 \sin\left(3\pi t + \frac{\pi}{3}\right) \Rightarrow \text{Amplitude} = 10$$

$$y_2 = 5(\sin 3\pi t + \sqrt{3} \cos 3\pi t)$$

$$y_2 = 10\left(\frac{1}{2} \sin 3\pi t + \frac{\sqrt{3}}{2} \cos 3\pi t\right)$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**




$$y_2 = 10 \left(\cos \frac{\pi}{3} \sin 3\pi t + \sin \frac{\pi}{3} \cos 3\pi t \right)$$

$$y_2 = 10 \sin \left(3\pi t + \frac{\pi}{3} \right) \Rightarrow \text{Amplitude} = 10$$

$$\text{So ratio of amplitudes} = \frac{10}{10} = 1$$

7. The ratio of the equivalent resistance of the network (shown in figure) between the points a and b when switch is open and switch is closed is $x : 8$. The value of x is _____.




Sol. 9

$$R_{\text{eq open}} = \frac{3R}{2}$$


$$R_{\text{eq closed}} = 2 \times \frac{R \times 2R}{3R} = \frac{4R}{3}$$

$$\frac{R_{\text{eq open}}}{R_{\text{eq closed}}} = \frac{3R}{2} \times \frac{3}{4R} = \frac{9}{8}$$

$$\therefore x = 9$$



8. Wires W_1 and W_2 are made of same material having the breaking stress of $1.25 \times 10^9 \text{ N/m}^2$. W_1 and W_2 have cross-sectional area of $8 \times 10^{-7} \text{ m}^2$ and $4 \times 10^{-7} \text{ m}^2$, respectively. Masses of 20kg and 10 kg hang from them as shown in the figure. The maximum mass that can be placed in the pan without breaking the wires is _____ kg. (Use $g = 10 \text{ m/s}^2$)



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



Sol. 40

$$B.S_1 = \frac{T_{1\max}}{8 \times 10^{-7}} \Rightarrow T_{1\max} = 8 \times 1.25 \times 100$$

$$= 1000\text{N}$$


$$B.S_2 = \frac{T_{2\max}}{4 \times 10^{-7}} \Rightarrow T_{2\max} = 4 \times 1.25 \times 100$$

$$= 500\text{N}$$

$$m = \frac{500 - 100}{10} = 40\text{kg}$$

- 9.** A tuning fork is vibrating at 250 Hz. The length of the shortest closed organ pipe that will resonate with the tuning fork will be _____ cm.
(Take speed of sound in air as 340ms^{-1})

Sol. 34



$$\frac{\lambda}{4} = l \Rightarrow \lambda = 4l$$

$$f = \frac{v}{\lambda} = \frac{v}{4l}$$


$$\Rightarrow 250 = \frac{340}{4l}$$

$$\Rightarrow l = \frac{34}{4 \times 25} = 0.34\text{m}$$

$$l = 34\text{cm}$$

- 10.** An ac circuit has an inductor and a resistor of resistance R in series, such that $X_L = 3R$. Now, a capacitor is added in series such that $X_C = 2R$. The ratio of new power factor with the old power factor of the circuit is $\sqrt{5} : x$. The value of x is _____.

Sol. 1



$$\cos \phi = \frac{R}{\sqrt{R^2 + 3R^2}}$$

$$= \frac{1}{\sqrt{10}}$$

$$\frac{\cos \phi'}{\cos \phi} = \frac{\sqrt{10}}{\sqrt{2}} = \frac{\sqrt{5}}{1}$$

$$\cos \phi' = \frac{R}{\sqrt{R^2 + R^2}}$$

$$= \frac{1}{\sqrt{2}}$$

$$\therefore x = 1$$



An Unmatched Experience of Offline

KOTA CLASSROOM For JEE

New batch Starting from : **22nd Sept. 2021**



हो चुकी है ऑफलाइन क्लासरूम की शुरुआत
अपने सपने को करो साकार, कोटा कोविंग के साथ

Directors of Nucleus Education & Wizard of Mathematics

Now Offline associated with Motion Kota Classroom



Nitin Vijay (NV Sir)
Managing Director
Exp. : 18 yrs



Akhilesh Kanther (AKK Sir)
Exp. : 17 yrs



Vishal Joshi (VJ Sir)
Exp. : 18 yrs



Surendra K. Mishra (SKM Sir)
Exp. : 16 yrs



Gavesh Bhardwaj (GB Sir)
Exp. : 17 yrs

Academic Pillars of JEE MOTION KOTA



Ram Ratan Dwivedi (RRD Sir)
Joint Director
Exp. : 20 yrs



Amit Verma (AV Sir)
Joint Director
Exp. : 16 yrs



Vijay Pratap Singh (VPS Sir)
Vice President
Exp. : 20 yrs



Nikhil Srivastava (NS Sir)
Head JEE Academics
Exp. : 17 yrs



Aatish Agarwal (AA Sir)
Sr. Faculty
Exp. : 17 yrs



Jayant Chittora (JC Sir)
Sr. Faculty
Exp. : 16 yrs



Anurag Garg (AG Sir)
Sr. Faculty
Exp. : 17 yrs



Arjun Gupta (Arjun Sir)
Sr. Faculty
Exp. : 14 yrs



Devki Nandan Pathak (DN Sir)
Sr. Faculty
Exp. : 13 yrs



Avinash Kishore (AVN Sir)
Sr. Faculty
Exp. : 9 yrs



Vipin Sharma (VS Sir)
Sr. Faculty
Exp. : 12 yrs



Durgesh Pandey (Pandey Sir)
Sr. Faculty
Exp. : 8 yrs

Join

JEE DROPPER BATCH

Online + Offline Mode

English & Hindi Medium

Batch Starting from :
22nd Sept. 2021