



**JEE
MAIN
July
2021**

CHEMISTRY
27th July 2021 [SHIFT - 1]
QUESTION WITH SOLUTION

JEE | NEET | Foundation

MOTIONTM

25000+
SELECTIONS SINCE 2007

अब मोशन ही है, सर्वश्रेष्ठ विकल्प!

MOTION welcomes

Directors of Nucleus Education & Wizard of Mathematics
Now Offline associated with Motion Kota Classroom



Nitin Vijay (NV Sir)
Managing Director
Exp. : 18 yrs



Akhilesh Kanther
(AKK Sir)
Exp. : 17 yrs



Vishal Joshi
(VJ Sir)
Exp. : 18 yrs



Surendra K. Mishra
(SKM Sir)
Exp. : 16 yrs



Gavesh Bhardwaj
(GB Sir)
Exp. : 17 yrs

Academic Pillars of JEE Motion Kota



Ram Ratan Dwivedi
(RRD Sir)
Joint Director
Exp. : 20 yrs



Amit Verma
(AV Sir)
Joint Director
Exp. : 16 yrs



Vijay Pratap Singh
(VPS Sir)
Vice President
Exp. : 20 yrs



Nikhil Srivastava
(NS Sir)
Head JEE Academics
Exp. : 17 yrs



Aatish Agarwal
(AA Sir)
Sr. Faculty
Exp. : 17 yrs



Jayant Chittora
(JC Sir)
Sr. Faculty
Exp. : 16 yrs



Anurag Garg
(AG Sir)
Sr. Faculty
Exp. : 17 yrs



Arjun Gupta
(Arjun Sir)
Sr. Faculty
Exp. : 14 yrs



Devki Nandan Pathak
(DN Sir)
Sr. Faculty
Exp. : 13 yrs



Avinash Kishore
(AVN Sir)
Sr. Faculty
Exp. : 9 yrs



Vipin Sharma
(VS Sir)
Sr. Faculty
Exp. : 12 yrs



Durgesh Pandey
(Pandey Sir)
Sr. Faculty
Exp. : 8 yrs

Join **JEE DROPPER BATCH** Kota Classroom

English & Hindi Medium

Online + Offline Mode

Batch Starting from: **4th August 2021**

Time to use all your skills, Now!

CRASH COURSE

JEE Advanced 2021

Starting from: **9th August '21**

Why should you choose?

- ◆ Live Lectures
- ◆ Best JEE Faculties
- ◆ Doubt Support on learning app
- ◆ Online test with discussion
- ◆ Personalized performance analysis




1. Which one of the following statements is **NOT** correct?
- (1) The dissolved oxygen concentration below 6 ppm inhibits fish growth
 - (2) Eutrophication indicates that water body is polluted
 - (3) Eutrophication leads to increase in the oxygen level in water
 - (4) Eutrophication leads to anaerobic conditions

Sol. (3)

Eutrophication leads to decrease in oxygen level of water.

3rd statement is incorrect


2.



The correct order of stability of given carbocation is :

- (1) C > A > D > B
- (2) D > B > C > A
- (3) A > C > B > D
- (4) D > B > A > C

Sol. (3)



Stable due to Resonance

3. Given below are two statements: One is labelled as **Assertion A** and the other is labelled as **Reason R**.

Assertion A :Lithium halides are some what covalent in nature.


Reason R : Lithium possess high polarization capability.

In the light of the above statements, choose the most appropriate answer from the options given below:


- (1) Both **A** and **R** are true but **R** is NOT the correct explanation of **A**
- (2) **A** is true but **R** is false
- (3) **A** is false but **R** is true
- (4) Both **A** and **R** are true and **R** is the correct explanation of **A**

Sol. (4)


Lithium due to small size has very high polarization capability and thus increases covalent nature in Halides.



Consider the above reaction and identify the product P :



Sol. (3)



(Anti - markownikov addition)

5. The statement that is INCORRECT about Ellingham diagram is :
- (1) Provides idea about the reaction rate.
 - (2) provides idea about free energy change.
 - (3) Provide idea about reduction of metal oxide.
 - (4) Provides idea about changes in the phase during the reaction.

Sol. (1)
Ellingham diagram is a plot between ΔG° and T and does not give any information regarding rate of reaction

6. For a reaction of order n, the unit of the rate constant is :

- (1) $\text{mol}^{1-n} \text{L}^{1-n} \text{s}^{-1}$
- (2) $\text{mol}^{1-n} \text{L}^{n-1} \text{s}^{-1}$
- (3) $\text{mol}^{1-n} \text{L}^{1-n} \text{s}$
- (4) $\text{mol}^{1-n} \text{L}^{2n} \text{s}^{-1}$

Sol. (2)
Rate = $k[A]^n$
comparing units


$$\frac{\left(\frac{\text{mol}}{\ell}\right)}{\text{sec}} = k \left(\frac{\text{mol}}{\ell}\right)^n$$

$$\Rightarrow k = \text{mol}^{(1-n)} \ell^{(n-1)} \text{s}^{-1}$$

7. The product obtained from the electrolytic oxidation of acidified sulphate solution, is:
- (1) $\text{HO}_2\text{SOSO}_2\text{H}$
 - (2) $\text{HO}_3\text{SOOSO}_3\text{H}$
 - (3) HSO_4^-
 - (4) $\text{HO}_3\text{SOSO}_3\text{H}$

Sol. (2)
anode : $2\text{SO}_4^{2-}(\text{aq}) \rightarrow (\text{S}_2\text{O}_8)^{-2} + 2\text{e}^-$
Cathode : $2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$
Electrolysis of concentrated solution of acidified sulphate solution yields $\text{H}_2\text{S}_2\text{O}_8$.

8. Presence of which reagent will affect the reversibility of the following reaction, and change it to a irreversible reaction:



- (1) HOCl
- (2) Liquid NH_3
- (3) dilute HNO_2
- (4) Concentrated HIO_3

Sol. (4)

Iodination of alkane is reversible reaction.

It can be irreversible in the presence of strong oxidising agent like conc. HNO_3 or conc. HIO_3

9. Match List-I with List-II :

List – I	List – II
(a) NaOH	(i) Acidic
(b) $\text{Be}(\text{OH})_2$	(ii) Basic
(c) $\text{Ca}(\text{OH})_2$	(iii) Amphoteric
(d) $\text{B}(\text{OH})_3$	
(e) $\text{Al}(\text{OH})_3$	

Choose the most appropriate answer from the option given below :

- (1) (a)-(ii), (b)-(ii), (c)-(iii), (d)-(ii), (e)-(iii)
- (2) (a)-(ii), (b)-(iii), (c)-(ii), (d)-(i), (e)-(iii)
- (3) (a)-(ii), (b)-(ii), (c)-(iii), (d)-(i), (e)-(iii)
- (4) (a)-(ii), (b)-(i), (c)-(ii), (d)-(iii), (e)-(iii)

Sol. (2)

$\text{NaOH} \longrightarrow$ Basic

$\text{Be}(\text{OH})_2 \longrightarrow$ Amphoteric

$\text{Ca}(\text{OH})_2 \longrightarrow$ Basic

$\text{B}(\text{OH})_3 \longrightarrow$ Acidic

$\text{Al}(\text{OH})_3 \longrightarrow$ Amphoteric

10. Given below are two statements:

Statement I :Aniline is less basic than acetamide.

Statement II :In aniline, the lone pair of electrons on nitrogen atom is delocalised over benzene ring due to resonance and hence less available to a proton.

Choose the **most appropriate** option;


- (1) Both statement I and statement II are true.
- (2) Both statement I and statement II are false.
- (3) Statement I is true but statement II is false.
- (4) Statement I is false but statement II is true.

Sol. (4)


Explanation :- aniline is more basic than acetamide because in acetamide, lone pair of nitrogen is delocalized to more electronegative element oxygen.

In Aniline lone pair of nitrogen delocalised over benzene ring.

11. Which one of the following compounds will give orange precipitate when treated with 2, 4-dinitrophenyl hydrazine?



Sol. (4)



Explanation \Rightarrow 2-4-D.N.P test is used for carbonyl compound (aldehyde & ketone)

12. Staggered and eclipsed conformers of ethane are:

- (1) Enantiomers (2) Rotamers
(3) Mirror images (4) Polymers

Sol. (2)


Staggered and eclipsed conformers of ethane also known as rotamers

13. The number of geometrical isomers found in the metal complexes $[\text{PtCl}_2(\text{NH}_3)_2]$, $[\text{Ni}(\text{CO})_4]$, $[\text{Ru}(\text{H}_2\text{O})_3\text{Cl}_3]$ and $[\text{CoCl}_2(\text{NH}_3)_4]^+$ respectively, are :



- (1) 1, 1, 1, 1 (2) 2, 1, 2, 2
(3) 2, 1, 2, 1 (4) 2, 0, 2, 2

Sol. (NTA-2)

Motion-4



$[\text{Ni}(\text{CO})_4] \rightarrow$ All ligands are same Zero Geometrical isomers



14. The parameters of the unit cell of a substance are $a = 2.5$, $b = 3.0$, $c = 4.0$, $\alpha = 90^\circ$, $\beta = 120^\circ$, $\gamma = 90^\circ$. The crystal system of the substance is :

- (1) Orthorhombic (2) Triclinic
(3) Hexagonal (4) Monoclinic

Sol. (4)

$a \neq b \neq c$ and $\alpha = \gamma = 90^\circ \neq \beta$
are parameters of monoclinic unit cell.

15. Given below are two statements:

Statement I : Rutherford's gold foil experiment cannot explain the line spectrum of hydrogen atom.

Statement II : Bohr's model of hydrogen atom contradicts Heisenberg's uncertainty principle.

In the light of the above statements, choose the most appropriate answer from the options given below :


- (1) Statement I is false but statement II is true.
(2) Both statement I and statement II are false.
(3) Statement I is true but statement II is false.
(4) Both statement I and statement II are true.

Sol. (4)

Rutherford's gold foil experiment only proved that electrons are held towards nucleus by electrostatic forces of attraction and move in circular orbits with very high speeds.

Bohr's model gave exact formula for simultaneous calculation of speed & distance of electron from the nucleus, something which was deemed impossible according to Heisenberg.

16.



The compound 'A' is a complementary base of _____ in DNA stands.

- (1) Cytosine (2) Adenine
(3) Guanine (4) Uracil

Sol. (2)


Given structure is Thymine and Thymine being paired with adenine.

17. The oxidation states of 'P' in $H_4P_2O_7$, $H_4P_2O_5$ and $H_4P_2O_6$, respectively, are:

- (1) 5, 3 and 4 (2) 6, 4 and 5
(3) 5, 4 and 3 (4) 7, 5 and 6


Sol. (1)

Oxidation state of P in $H_4P_2O_7$, $H_4P_2O_5$ and $H_4P_2O_6$ is 5, 3 & 4 respectively




$$2x + 4(+1) + 7(-2) = 0$$

$$x = +5$$



$$2x + 4(+1) + 5(-2) = 0$$

$$x = +3$$



$$2x + 4(+1) + 6(-2) = 0$$

$$x = +4$$

18. Which one among the following chemical tests is used to distinguish monosaccharide from disaccharide?

- (1) Barfoed test (2) Seliwanoff's test
 (3) Tollen's test (4) Iodine test

Sol. (1)

Barford test is used for distinguish mono-saccharide from disaccharide

19. Match List-I with List-II :

List – I (Drug)	List – II (Class of Drug)
(a)Furacin	(i) Antibiotic
(b)Arsphenamine	(ii) Tranquilizers
(c)Dimetone	(iii) Antiseptic
(d)Valium	(iv) Synthetic antihistamines

Choose the **most appropriate** match :

- (1) (a)-(iii), (b)-(iv), (c)-(ii),(d)-(i)
 (2) (a)-(i), (b)-(iii), (c)-(iv),(d)-(ii)
 (3) (a)-(ii), (b)-(i), (c)-(iii), (d)-(iv)
 (4) (a)-(iii), (b)-(i), (c)-(iv),(d)-(ii)



Sol. (4)

- furacine acts as Antiseptic
 → Arsphenamine also known as salvarsan acts as antibiotic
 → Dimetone is synthetic histamine
 → valium is a Tranqlizer

20. The type of hybridisation and magnetic property of the complex $[\text{MnCl}_6]^{3-}$, respectively, are :

- (1) d^2sp^3 and paramagnetic (2) d^2sp^3 and diamagnetic
 (3) sp^3d^2 and paramagnetic (4) sp^3d^2 and diamagnetic

Sol. (3)



Paramagnetic and having 4 unpaired electrons.

SECTION -B

1. In gaseous triethyl amine the "-C-N-C-" bond angle is _____ degree.

Sol. 180

In gaseous triethyl amine the "-C-N-C-" bond angle is 108 degree.

2. The density of NaOH solution is 1.2 g cm^{-3} . The molality of this solution is _____ m. (Round off to the Nearest Integer)

[Use: Atomic masses: Na: 23.0 u, O: 16.0 u, H: 1.0 u, Density of H_2O : 1.0 g cm^{-3}]

Sol. 5

Consider 1 l solution

$$\begin{aligned} \text{mass of solution} &= (1.2 \times 1000) \text{ g} \\ &= 1200 \text{ gm} \end{aligned}$$

Neglecting volume of NaOH

$$\text{Mass of water} = 1000 \text{ gm}$$


$$\begin{aligned} \Rightarrow \text{Mass of NaOH} &= (1200 - 1000) \text{ gm} \\ &= 200 \text{ gm} \end{aligned}$$

$$\Rightarrow \text{Moles of NaOH} = \frac{200 \text{ g}}{50 \text{ g/mol}} = 5 \text{ mol}$$

$$\Rightarrow \text{molality} = \frac{5 \text{ mol}}{1 \text{ kg}} = 5 \text{ m}$$

3. $\text{PCl}_5 \rightleftharpoons \text{PCl}_3 + \text{Cl}_2$ $K_c = 1.844$
3.0 moles of PCl_5 is introduced in a 1 L closed reaction vessel at 380 K. The number of moles of PCl_5 at equilibrium is _____ $\times 10^{-3}$. (Round off to the Nearest Integer)

Sol. NTA-1400, MOTION-1396



$$t = 0 \quad 3 \text{ moles}$$

$$t = \infty \quad \quad \quad x \quad \quad \quad x$$

$$\Rightarrow \frac{[\text{PCl}_3][\text{Cl}_2]}{[\text{PCl}_5]} = \frac{x^2}{3-x} = 1.844$$

$$\Rightarrow x^2 + 1.844 - 5.532 = 0$$

$$\Rightarrow x = \frac{-1.844 + \sqrt{(1.844)^2 + 4 \times 5.532}}{2}$$

$$\cong 1.604$$


$$\Rightarrow \text{Moles of } \text{PCl}_5 = 3 - 1.604 \cong 1.396$$

4. For water at 100°C and 1 bar,
 $\Delta_{\text{vap}} H - \Delta_{\text{vap}} U = \underline{\hspace{2cm}} \times 10^2 \text{ J mol}^{-1}$. (Round off to the Nearest Integer)

[Use : $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$]

[Assume volume of $\text{H}_2\text{O}(l)$ is much smaller than volume of $\text{H}_2\text{O}(g)$. Assume $\text{H}_2\text{O}(g)$ treated as an ideal gas]

Sol. 31



$$\Delta H = \Delta U + \Delta n_g RT$$

for 1 mole waters;

$$\Delta n_g = 1$$

$$\therefore \Delta n_g RT = 1 \text{ mol} \times 8.31 \text{ J/mol-k} \times 373 \text{ K}$$

$$= 3099.63 \text{ J} \cong 31 \times 10^2 \text{ J}$$

5. The difference between bond orders of CO and NO is NO^\oplus is $\frac{x}{2}$ where $x =$ _____.

(Round off to the Nearest Integer)

Sol. 0

Bond order of CO = 3

Bond order of $\text{NO}^+ = 3$

$$\text{Difference} = 0 = \frac{x}{2}$$


$$x = 0$$

6. The number of geometrical isomers possible in triamminetrinitrocobalt (III) is X and in trioxalatochromate (III) is Y. Then the value of $X + Y$ is _____.

Sol. 2


Triamminetrinitrocobalt(III) $\rightarrow [\text{Co}(\text{NO}_2)_3(\text{NH}_3)_3]$

trioxalatochromate(III) ion $\rightarrow [\text{Cr}(\text{C}_2\text{O}_4)_3]^{3-}$



Two geometrical isomers (X)

$[\text{Cr}(\text{C}_2\text{O}_4)_3]^{3-}$



$$X + Y = 2 + 0 = 2.0$$

7. CO_2 gas adsorbs on charcoal following Freundlich adsorption isotherm. For a given amount of charcoal, the mass of CO_2 adsorbed becomes 64 times when the pressure of CO_2 is doubled. The value of n in the Freundlich isotherm equation is _____ $\times 10^{-2}$. (Round off to the Nearest Integer)

Sol. 17

Freundlich isotherm. ;

$$\frac{x}{m} = k.p^n$$

Substituting values;

$$\left(\frac{64}{1}\right) = (2)^{\frac{1}{n}} \Rightarrow n = \frac{1}{6} = 0.166$$

$$\cong 17 \times 10^{-2}$$

8. 1.46 g of a biopolymer dissolved in a 100 mL water at 300 K exerted an osmotic pressure of 2.42×10^{-3} bar. The molar mass of the biopolymer is _____ $\times 10^4$ g mol⁻¹. (Round off to the Nearest Integer)

[Use : $R = 0.083$ L bar mol⁻¹ K⁻¹]

Sol. 15

$\pi = CRT$; π = osmotic pressure

C = molarity

T = Temperature of solution

let the molar mass be M gm / mol

$$2.42 \times 10^{-3} \text{ bar} = \frac{\left(\frac{1.46\text{g}}{\text{Mgm/mol}}\right)}{0.1\ell} \times \left(\frac{0.083\ell\text{-bar}}{\text{mol-K}}\right) \times (300\text{K})$$

$$\Rightarrow M = 15.02 \times 10^4 \text{ g/mol}$$

9. An organic compound is subjected to chlorination to get compound A using 5.0g of chlorine. When 0.5 g of compound A is reacted with AgNO_3 [Carius Method], the percentage of chlorine in compound A is _____ when it forms 0.3849 g of AgCl . (Round off to the Nearest Integer)

(Atomic masses of Ag and Cl are 107.87 and 35.5 respectively)

Sol. 19

$$n_{\text{Cl}} \text{ in compound} = n_{\text{AgCl}} = \frac{0.3849\text{g}}{(107.87 + 35.5)} \text{ g/mol}$$

$$\Rightarrow \text{mass of chlorine} = n_{\text{Cl}} \times 35.5 = 0.0953 \text{ gm}$$

$$\Rightarrow \% \text{ wt of chlorine} = \frac{0.0953}{0.5} \times 100$$

$$= 19.06 \%$$


- 10.** The conductivity of a weak acid HA of concentration 0.001 mol L^{-1} is $2.0 \times 10^{-5} \text{ S cm}^{-1}$. If $\Lambda_m^\circ(\text{HA}) = 190 \text{ S cm}^2 \text{ mol}^{-1}$, the ionization constant (K_a) of HA is equal to _____ $\times 10^{-6}$. (Round off to the Nearest Integer)

Sol. 12

$$\Lambda_m = 1000 \times \frac{\kappa}{M}$$

$$= 1000 \times \frac{2 \times 10^{-5}}{0.001} = 20 \text{ S cm}^2 \text{ mol}^{-1}$$

$$\Rightarrow \alpha = \frac{\Lambda_m}{\Lambda_m^\circ} = \frac{20}{190} = \left(\frac{2}{19}\right)$$



$$\Rightarrow K_a = 0.001 \left(\frac{\alpha^2}{1 - \alpha} \right) = \frac{0.001 \times \left(\frac{2}{19}\right)^2}{1 - \left(\frac{2}{19}\right)}$$

$$= 12.3 \times 10^{-6}$$

अब मोशन ही है, सर्वश्रेष्ठ विकल्प !

Motion welcomes

Directors of Nucleus Education & Wizard of Mathematics

Now Offline associated with Motion Kota Classroom



Nitin Vijay (NV Sir)
Managing Director
Exp. : 18 yrs



Akhilesh Kanther
(AKK Sir)
Exp. : 17 yrs



Vishal Joshi
(VJ Sir)
Exp. : 18 yrs



Surendra K. Mishra
(SKM Sir)
Exp. : 16 yrs



Gavesh Bhardwaj
(GB Sir)
Exp. : 17 yrs

Academic Pillars of JEE Motion Kota



Ram Ratan Dwivedi
(RRD Sir)
Joint Director
Exp. : 20 yrs



Amit Verma
(AV Sir)
Joint Director
Exp. : 16 yrs



Vijay Pratap Singh
(VPS Sir)
Vice President
Exp. : 20 yrs



Nikhil Srivastava
(NS Sir)
Head JEE Academics
Exp. : 17 yrs



Aatish Agarwal
(AA Sir)
Sr. Faculty
Exp. : 17 yrs



Jayant Chittora
(JC Sir)
Sr. Faculty
Exp. : 16 yrs



Anurag Garg
(AG Sir)
Sr. Faculty
Exp. : 17 yrs



Arjun Gupta
(Arjun Sir)
Sr. Faculty
Exp. : 14 yrs



Devki Nandan Pathak
(DN Sir)
Sr. Faculty
Exp. : 13 yrs



Avinash Kishore
(AVN Sir)
Sr. Faculty
Exp. : 9 yrs



Vipin Sharma
(VS Sir)
Sr. Faculty
Exp. : 12 yrs



Durgesh Pandey
(Pandey Sir)
Sr. Faculty
Exp. : 8 yrs

Join **JEE DROPPER BATCH** Kota Classroom

English & Hindi Medium

Online + Offline Mode

Batch Starting from: **4th August 2021**

Time to use all your skills, Now!

CRASH COURSE

JEE Advanced 2021

Starting from: **9th August '21**

Why should you choose?

- ◆ Live Lectures
- ◆ Best JEE Faculties
- ◆ Doubt Support on learning app
- ◆ Online test with discussion
- ◆ Personalized performance analysis

