



JEE
MAIN
July
2021

MATHS

22nd July 2021 [SHIFT – 2]

QUESTION WITH SOLUTION

JEE | NEET | Foundation

MOTION™

25000+
SELECTIONS SINCE 2007

अब मोशन ही है, सर्वश्रेष्ठ विकल्प!

MOTION welcomes

Directors of Nucleus Education & Wizard of Mathematics

Now Offline associated with Motion Kota Classroom



Nitin Vijay (NV Sir)
Managing Director
Exp. : 18 yrs



Akhilesh Kanther
(AKK Sir)
Exp. : 17 yrs



Vishal Joshi
(VJ Sir)
Exp. : 18 yrs



Surendra K. Mishra
(SKM Sir)
Exp. : 16 yrs



Gavesh Bhardwaj
(GB Sir)
Exp. : 17 yrs

Academic Pillars of JEE Motion Kota



Ram Ratan Dwivedi
(RRD Sir)
Joint Director
Exp. : 20 yrs



Amit Verma
(AV Sir)
Joint Director
Exp. : 16 yrs



Vijay Pratap Singh
(VPS Sir)
Vice President
Exp. : 20 yrs



Nikhil Srivastava
(NS Sir)
Head JEE Academics
Exp. : 17 yrs



Atish Agarwal
(AA Sir)
Sr. Faculty
Exp. : 17 yrs



Jayant Chittora
(JC Sir)
Sr. Faculty
Exp. : 16 yrs



Anurag Garg
(AG Sir)
Sr. Faculty
Exp. : 17 yrs



Arjun Gupta
(Arjun Sir)
Sr. Faculty
Exp. : 14 yrs



Devki Nandan Pathak
(DN Sir)
Sr. Faculty
Exp. : 13 yrs



Avinash Kishore
(AVN Sir)
Sr. Faculty
Exp. : 9 yrs



Vipin Sharma
(VS Sir)
Sr. Faculty
Exp. : 12 yrs



Durgesh Pandey
(Pandey Sir)
Sr. Faculty
Exp. : 8 yrs

Join JEE DROPPER BATCH Kota Classroom

English & Hindi Medium

Online + Offline Mode

Batch Starting from: 4th August 2021

Time to use all your skills, Now!

CRASH COURSE JEE Advanced 2021

Starting from: 9th August'21

Why should you choose?

- ◆ Live Lectures
- ◆ Best JEE Faculties
- ◆ Doubt Support on learning app
- ◆ Online test with discussion
- ◆ Personalized performance analysis



SECTION A

- 1.** Let $f : R \rightarrow R$ be defined as

$$f(x) = \begin{cases} -\frac{4}{3}x^3 + 2x^2 + 3x, & x > 0 \\ 3xe^x, & x \leq 0 \end{cases}$$

Then f is increasing function in the interval.

- (1) $\left(-1, \frac{3}{2}\right)$ (2) $\left(\frac{-1}{2}, 2\right)$ (3) $(0, 2)$ (4) $(-3, -1)$

- Sol.** **(1)**

For $x > 0$ $f'(x) = -4x^2 + 4x + 3$

$F(x)$ is increasing in $\left(-\frac{1}{2}, \frac{3}{2}\right)$

For $x \leq 0$ $f'(x) = 3e^x(1+x)$

$F'(x) > 0 \forall x \in (-1, 0)$

$\Rightarrow f(x)$ is increasing in $(-1, 0)$

So, in complete domain $f(x)$ is increasing in $\left(-1, \frac{3}{2}\right)$

- 2.** Let $[x]$ denote the greatest integer less than or equal to x . Then, the values of $x \in R$ satisfying the equation $[e^x]^2 + [e^x + 1] - 3 = 0$ lie in the interval:

- (1) $[\log_e 2, \log_e 3]$ (2) $[0, 1/e]$ (3) $[0, \log_e 2)$ (4) $[1, e)$

- Sol.** **(3)**

$$[e^x]^2 + [e^x + 1] - 3 = 0$$

$$\Rightarrow [e^x]^2 + [e^x + 1] + 1 - 3 = 0$$

$$\text{Let } [e^x] = t$$

$$\Rightarrow t^2 + t - 2 = 0$$

$$\Rightarrow t = -2, 1$$

$$[e^x] = -2 \text{ (Not possible)}$$

$$\text{or } [e^x] = 1 \quad \therefore 1 \leq e^x < 2$$

$$\Rightarrow \ln(1) \leq x < \ln(2)$$

$$\Rightarrow 0 \leq x < \ln(2)$$

$$\Rightarrow x \in [0, \ln 2)$$

- 3.** Let n denote the number of solutions of the equation $z^2 + \bar{3z} = 0$, where z is a complex

number. Then the value of $\sum_{k=0}^{\infty} \frac{1}{n^k}$ is equal to:

- (1) 1 (2) 2 (3) $\frac{4}{3}$ (4) $\frac{3}{2}$

Sol. (3)

$$z^2 + 3\bar{z} = 0$$

$$\text{Put } z = x + iy$$

$$\Rightarrow x^2 - y^2 + 2ixy + 3(x - iy) = 0$$

$$\Rightarrow (x^2 - y^2 + 3x) + i(2xy - 3y) = 0 + i0$$

$$\therefore x^2 - y^2 + 3x = 0 \quad \dots\dots(1)$$

$$2xy - 3y = 0 \quad \dots\dots$$

$$x = \frac{3}{2}, y = 0$$

$$\text{Put } x = \frac{3}{2} \text{ in equation (1)}$$

$$\frac{9}{4} - y^2 + \frac{9}{2} = 0$$

$$y^2 = \frac{27}{4} \Rightarrow y = \pm \frac{3\sqrt{3}}{2}$$

$$\therefore (x, y) = \left(\frac{3}{2}, \frac{3\sqrt{3}}{2}\right), \left(\frac{3}{2}, -\frac{3\sqrt{3}}{2}\right)$$

$$\text{Put } y = 0 \Rightarrow x^2 - 0 + 3x = 0$$

$$x = 0, -3$$

$$\therefore (x, y) = (0, 0), (-3, 0)$$

∴ No of solutions = n = 4

$$\begin{aligned} \sum_{k=0}^{\infty} \left(\frac{1}{n^k} \right) &= \sum_{k=0}^{\infty} \left(\frac{1}{4^k} \right) \\ &= \frac{1}{1} + \frac{1}{4} + \frac{1}{16} + \frac{1}{64} + \dots\dots \\ &= \frac{1}{1 - \frac{1}{4}} = \frac{4}{3} \end{aligned}$$

4. Let $A = [a_{ij}]$ be a real matrix of order 3×3 , such that $a_{i1} + a_{i2} + a_{i3} = 1$, for $i=1,2,3$. Then, the sum of all the entries of the matrix A^3 is equal to:

(1) 1

(2) 3

(3) 2

(4) 9

Sol. (3)

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

$$\text{Let } X = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$AX = \begin{bmatrix} a_{11} + a_{12} + a_{13} \\ a_{21} + a_{22} + a_{23} \\ a_{31} + a_{32} + a_{33} \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$\Rightarrow AX = X$$

Replace X by AX

$$A^2X = AX = X$$

Replace X by AX

$$A^3X = AX = X$$

$$\text{Let } A^3 = \begin{bmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{bmatrix}$$

$$A^3 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} x_1 + x_2 + x_3 \\ y_1 + y_2 + y_3 \\ z_1 + z_2 + z_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Sum of all the element = 3

5. Let a line $L: 2x+y=k, k>0$ be a tangent to the hyperbola $x^2-y^2=3$. If L is also a tangent to the parabola $y^2=ax$, then a is equal to :

(1) 24 (2) -12 (3) -24 (4) 12

Sol. **(3)**

Tangent to hyperbola of

Slope $m = -2$ (given)

$$y = -2x \pm \sqrt{3(3)}$$

$$(y = mx \pm \sqrt{a^2m^2 - b^2})$$

$$\Rightarrow y + 2x = \pm 3 \Rightarrow 2x + y = 3 (k > 0)$$

For parabola $y^2 = ax$

$$y = mx + \frac{\alpha}{4}$$

$$\Rightarrow y = -2x + \frac{\alpha}{-8}$$

$$\Rightarrow \frac{\alpha}{-8} = 3$$

$$\Rightarrow -24.$$

6. If $\int_0^{100\pi} \frac{\sin^2 x}{e^{\left(\frac{x}{\pi} - \left[\frac{x}{\pi}\right]\right)}} dx = \frac{\alpha\pi^3}{1+4\pi^2}, \alpha \in R,$

where $[x]$ is the greatest integer less than or equal to x , then the value of α is :

(1) 100 (1 - e) (2) 200 (1 - e⁻¹) (3) 150 (e⁻¹ - 1) (4) 50 (e - 1)

Sol. **(2)**

$$I = \int_0^{100\pi} \frac{\sin^2 x}{e^{\left(\frac{x}{\pi}\right)}} dx = 100 \int_0^{\pi} \frac{\sin^2 x}{e^{\frac{x}{\pi}}} dx$$

$$100 \int_0^{\pi} e^{-x/\pi} \frac{(1 - \cos 2x)}{2} dx$$

$$= 50 \left\{ \int_0^{\pi} e^{-x/\pi} dx - \int_0^{\pi} e^{-x/\pi} \cos 2x dx \right\}$$

$$I_1 = \int_0^{\pi} e^{-x/\pi} dx = \left[-\pi e^{-x/\pi} \right]_0^{\pi} = \pi (1 - e^{-1})$$

$$I_2 = \int_0^{\pi} e^{-x/\pi} \cos 2x dx$$

$$\begin{aligned}
 &= -\pi e^{-x/\pi} \cos 2x]_0^\pi - \int_{-\pi}^{\pi} e^{-x/\pi} (-2 \sin 2x) dx \\
 &= \pi(1 - e^{-1}) - 2\pi \int_0^{\pi} e^{-x/\pi} \sin 2x dx \\
 &= \pi(1 - e^{-1}) - 2\pi \left\{ -\pi e^{-x/\pi} \sin 2x]_0^\pi - \int_0^\pi -\pi e^{-x/\pi} 2 \cos 2x dx \right\} \\
 &= \pi(1 - e^{-1}) - 4\pi^2 I_2 \\
 \Rightarrow I_2 &= \frac{\pi(1 - e^{-1})}{1 + 4\pi^2} \\
 \therefore I &= 50 \left\{ \pi(1 - e^{-1}) - \frac{\pi(1 - e^{-1})}{1 + 4\pi^2} \right\} \\
 &= \frac{200(1 - e^{-1})\pi^3}{1 + 4\pi^2}
 \end{aligned}$$

7. Four dice are thrown simultaneously and the numbers shown on these dice are recorded in 2×2 matrices. The probability that such formed matrices have all different entries and are non-singular, is :

(1) $\frac{23}{81}$

(2) $\frac{22}{81}$

(3) $\frac{45}{162}$

(4) $\frac{43}{162}$

Sol. (4)

$$A = \begin{vmatrix} a & b \\ c & d \end{vmatrix} \quad |A| = ad - bc$$

Total case = 6^4

For non-singular matrix $|A| \neq 0 \Rightarrow ad - bc \neq 0$

$\Rightarrow ad \neq bc$

And a, b, c, d are all different numbers in the set $\{1, 2, 3, 4, 5, 6\}$

Now for $ad = bc$

(i) $6 \times 1 = 2 \times 3$

$$\begin{aligned}
 &\Rightarrow a = 6, b = 2, c = 3, d = 1 \\
 &\text{or } a = 1, b = 2, c = 3, d = 6 \\
 &\quad \vdots
 \end{aligned}
 \left. \begin{array}{l} \\ \\ \end{array} \right\} 8 \text{ such cases}$$

(ii) $6 \times 2 = 3 \times 4$

$$\begin{aligned}
 &\Rightarrow a = 6, b = 2, c = 3, d = 2 \\
 &\text{or } a = 1, b = 3, c = 4, d = 6 \\
 &\quad \vdots
 \end{aligned}
 \left. \begin{array}{l} \\ \\ \end{array} \right\} 8 \text{ such cases}$$

favourable cases

$$= {}^6C_4 \underline{4} - 16$$

required probability

$$= \frac{{}^6C_4 \underline{-} 16}{6^4} = \frac{43}{162}$$

- 8.** Let three vectors \vec{a}, \vec{b} and \vec{c} be such that $\vec{a} \times \vec{b} = \vec{c}, \vec{b} \times \vec{c} = \vec{a}$ and $|\vec{a}| = 2$.

Then which one of the following is not true?

(1) Projection of \vec{a} on $(\vec{b} \times \vec{c})$ is 2 (2) $|3\vec{a} + \vec{b} - 2\vec{c}|^2 = 51$

(3) $[\vec{a} \vec{b} \vec{c}] + [\vec{c} \vec{a} \vec{b}] = 8$ (4) $\vec{a} \times ((\vec{b} + \vec{c}) \times (\vec{b} - \vec{c})) = \vec{0}$

Sol. (2)

(1) Projection of \vec{a} on $\vec{b} \times \vec{c}$

$$= \frac{\vec{a} \cdot (\vec{b} \times \vec{c})}{|\vec{b} \times \vec{c}|} = \frac{\vec{a} \cdot \vec{a}}{|\vec{a}|} = |\vec{a}| = 2$$

(2) $\vec{a} \times \vec{b} = \vec{c}$ and $\vec{b} \times \vec{c} = \vec{a}$

$\Rightarrow \vec{a}, \vec{b}, \vec{c}$ are mutually \perp vectors.

$$|\vec{a} \times \vec{b}| = |\vec{c}| \Rightarrow |\vec{a}| |\vec{b}| = |\vec{c}| \Rightarrow |\vec{b}| = |\vec{c}| / 2$$

Also, $|\vec{b} \times \vec{c}| = |\vec{a}| \Rightarrow |\vec{b}| |\vec{c}| = 2 \Rightarrow |\vec{c}| = 2 \& |\vec{b}| = 1$

$$|\vec{a} + \vec{b} - 2\vec{c}|^2 = (3\vec{a} + \vec{b} - 2\vec{c}) \cdot (3\vec{a} + \vec{b} - 2\vec{c})$$

$$= 9|\vec{a}|^2 + |\vec{b}|^2 + 4|\vec{c}|^2$$

$$= (9 \times 4) + 1 + (4 \times 4)$$

$$= 36 + 1 + 16 = 53$$

$$(3) [\vec{a} \vec{b} \vec{c}] + [\vec{c} \vec{a} \vec{b}] = 2[\vec{a} \vec{b} \vec{c}] = 2\vec{a} \cdot (\vec{b} \times \vec{c})$$

$$= 2\vec{a} \cdot \vec{a} = 2|\vec{a}|^2 = 8$$

(4) $\vec{a} \times ((\vec{b} + \vec{c}) \times (\vec{b} - \vec{c}))$

$$= \vec{a}(-\vec{b} \times \vec{c} + \vec{c} \times \vec{b}) = -2(\vec{a} \times (\vec{b} \times \vec{c}))$$

$$= -2(\vec{a} \times \vec{a}) = \vec{0}$$

- 9.** Let S_n denote the sum of first n -terms of an arithmetic progression. If $S_{10} = 530$, $S_5 = 140$, then $S_{20} - S_6$ is equal to :

(1) 1852 (2) 1842 (3) 1872 (4) 1862

Sol. (4)

$$S_{10} = 530 \Rightarrow \frac{10}{2} \{2a + 9d\} = 530$$

$$\Rightarrow 2a + 9d = 106 \dots(1)$$

$$\text{and } S_5 = 140 \Rightarrow \frac{5}{2} \{2a + 4d\} = 140$$

$$\Rightarrow 2a + 4d = 56 \dots(2)$$

$$\Rightarrow 5d = 50 \Rightarrow d = 10 \Rightarrow a = 8$$

$$\text{Now, } S_{20} - S_6 = \frac{20}{2} \{2a + 19d\} - \frac{6}{2} \{2a + 5d\}$$

$$= 14a + 175d$$

$$= (14 \times 8) + (175 \times 10)$$

$$= 1862$$

- 10.** Which of the following Boolean expressions is not a tautology ?

$$(1) (\sim p \Rightarrow q) \vee (\sim q \Rightarrow p)$$

$$(2) (q \Rightarrow p) \vee (\sim q \Rightarrow p)$$

$$(3) (p \Rightarrow q) \vee (\sim q \Rightarrow p)$$

$$(4) (p \Rightarrow \sim q) \vee (\sim q \Rightarrow p)$$

Sol. (1)

$$(1) (\sim p \rightarrow q) \vee (\sim q \rightarrow p)$$

$$= (p \vee q) \vee (q \vee p)$$

$$= (p \vee p) \vee (q \vee q)$$

$$= p \vee q$$

Which is not a tautology.

$$(2) (q \rightarrow p) \vee (\sim q \rightarrow p)$$

$$= (\sim q \vee p) \vee (q \vee p)$$

$$= (\sim q \vee q) \vee p$$

$$= t \vee p = t$$

$$(3) (p \rightarrow q) \vee (\sim q \rightarrow p)$$

$$= (\sim p \vee q) \vee (q \vee p)$$

$$= (\sim p \vee p) \vee q$$

$$= t \vee q = t$$

$$(4) (p \rightarrow \sim q) \vee (\sim q \rightarrow p)$$

$$= (\sim p \vee \sim q) \vee (q \vee p)$$

$$= (\sim p \vee q) \vee (\sim q \vee q)$$

$$= t \vee t = t$$

- 11.** Let the circle $S : 36x^2 + 36y^2 - 108x + 120y + C = 0$ be such that it neither intersects nor touches the co-ordinate axes. If the point of intersection of the lines, $x-2y=4$ and $2x-y=5$ lies inside the circle S , then :

$$(1) 100 < C < 156$$

$$(2) \frac{25}{9} < C < \frac{13}{3}$$

$$(3) 81 < C < 156$$

$$(4) 100 < C < 165$$


Sol. (1)

$$S : 36x^2 + 36y^2 - 108x + 120y + C = 0$$

$$\Rightarrow x^2 + y^2 - 3x + \frac{10}{3}y + \frac{C}{36} = 0$$

$$\text{Centre} = (-g, -f) = \left(\frac{3}{2}, -\frac{10}{6}\right)$$

$$\text{radius} = r = \sqrt{\frac{9}{4} + \frac{100}{36} - \frac{C}{36}}$$



Now,

$$\Rightarrow r < \frac{3}{2}$$

$$\Rightarrow \frac{9}{4} + \frac{100}{36} - \frac{C}{36} < \frac{9}{4}$$

$$\Rightarrow C > 100$$

Now point of intersection of $x - 2y = 4$ and $2x - y = 5$ is $(2, -1)$, which lies inside the circle S.

$$\therefore S(2, -1) < 0$$

$$\Rightarrow (2)^2 + (-1)^2 - 3(2) + \frac{10}{3}(-1) + \frac{C}{36} < 0$$

$$\Rightarrow 4 + 1 - 6 - \frac{10}{3} + \frac{C}{36} < 0$$

$$C < 156$$

From (1) & (2)

$100 < C < 156$ Ans.

- 12.** The number of solutions of $\sin^7 x + \cos^7 x = 1, x \in [0, 4\pi]$ is equal to :

(1) 5

(2) 9

(3) 11

(4) 7

Sol. (1)

$$\sin^7 x \leq \sin^2 x \leq 1 \quad \dots(1)$$

$$\text{and } \cos^7 x \leq \cos^2 x \leq 1 \quad \dots(2)$$

$$\text{also } \sin^2 x + \cos^2 x = 1$$

\Rightarrow equality must hold for (1) & (2)

$$\Rightarrow \sin^7 x = \sin^2 x \text{ & } \cos^7 x = \cos^2 x$$

$$\Rightarrow \sin x = 0 \text{ & } \cos x = 1 \text{ or}$$

$$\cos x = 0 \text{ & } \sin x = 1$$

$$\Rightarrow x = 0, 2\pi, 4\pi, \frac{\pi}{2}, \frac{5\pi}{2}$$

\Rightarrow 5 solutions

- 13.** Let $E_1 : \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, a > b$. Let E_2 be another ellipse such that it touches the end points of major axis of E_1 and the foci E_2 are the end points of minor axis of E_1 . If E_1 and E_2 have same eccentricities, then its value is :


$$(1) \frac{-1 + \sqrt{3}}{2}$$

$$(2) \frac{-1 + \sqrt{6}}{2}$$

$$(3) \frac{-1 + \sqrt{5}}{2}$$

$$(4) \frac{-1 + \sqrt{8}}{2}$$

Sol. (3)



$$e^2 = 1 - \frac{b^2}{a^2} \quad -(1)$$

$$e^2 = 1 - \frac{a^2}{c^2} \quad -(2)$$

$$\Rightarrow \frac{b^2}{a^2} = \frac{a^2}{c^2}$$

$$\Rightarrow c^2 = \frac{a^4}{b^2} \Rightarrow c = \frac{a^2}{b}$$

Also $b = ce$

$$\therefore \frac{b}{e} = \frac{a^2}{b} \Rightarrow \frac{b^2}{a^2} = e$$

$$e^2 = 1 - b^2/a^2 \Rightarrow e^2 + e - 1 = 0$$


14. Let L be the line of intersection of planes $\vec{r} \cdot (\hat{i} - \hat{j} + 2\hat{k}) = 2$ and $\vec{r} \cdot (2\hat{i} + \hat{j} - \hat{k}) = 2$. If $P(\alpha, \beta, \gamma)$ is the foot of perpendicular on L from the point $(1, 2, 0)$, then the value of $35(\alpha + \beta + \gamma)$ is equal to :

(1) 134 (2) 119 (3) 143 (4) 101

Sol. (2)

$$P_1 : x - y + 2z = 2$$

$$P_2 : 2x + y - 3 = 2$$



Let line of Intersection of planes P_1 and P_2 cuts xy plane in point Q .

\Rightarrow z-coordinate of point Q is zero

$$\Rightarrow \begin{cases} x - y = 2 \\ \text{and } 2x + y = 2 \end{cases} \Rightarrow x = \frac{4}{3}, y = \frac{-2}{3}$$

$$\Rightarrow Q\left(\frac{4}{3}, \frac{-2}{3}, 0\right)$$

Vector parallel to the line of intersection

$$\vec{a} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 2 \\ 2 & 1 & -1 \end{vmatrix} = -\hat{i} + 5\hat{j} + 3\hat{k}$$

Equation of Line of intersection

$$\frac{x - \frac{4}{3}}{-1} = \frac{y + \frac{2}{3}}{5} = \frac{z - 0}{3} = \lambda \text{ (say)}$$

Let coordinates of foot of perpendicular be

$$F\left(-\lambda + \frac{4}{3}, 5\lambda - \frac{2}{3}, 3\lambda\right)$$

$$\overline{PF} = \left(-\lambda + \frac{1}{3}\right)\hat{i} + \left(5\lambda - \frac{8}{3}\right)\hat{j} + (3\lambda)\hat{k}$$

$$\overline{PF} \cdot \vec{a} = 0$$

$$\Rightarrow \lambda - \frac{1}{3} + 25\lambda - \frac{40}{3} + 9\lambda = 0$$

$$\Rightarrow 35\lambda = \frac{41}{3} \Rightarrow \lambda = \frac{41}{105}$$

$$\text{Now, } \alpha = -\lambda + \frac{4}{3}, \beta = 5\lambda - \frac{2}{3}, \gamma = 3\lambda$$

$$\Rightarrow \alpha + \beta + \gamma = 7\lambda + \frac{2}{3}$$

$$= 7\left(\frac{41}{105}\right) + \frac{2}{3}$$

$$= \frac{51}{15}$$

$$\Rightarrow 35(\alpha + \beta + \gamma) = \frac{51}{15} \times 35 = 119$$

15. Let $f: R \rightarrow R$ be defined as $f(x) = \begin{cases} \frac{x^3}{(1-\cos 2x)^2} \log_e \left(\frac{1+2xe^{-2x}}{(1-xe^{-x})^2} \right) & , x \neq 0 \\ \alpha & , x = 0 \end{cases}$

If f is continuous at $x=0$, then α is equal to :

- (1) 1 (2) 0 (3) 3 (4) 2

Sol. (1)

For continuity

$$\lim_{x \rightarrow 0} \frac{x^3}{4 \sin^4 x} (\ln(1 + 2e^{-2x}) - 2\ln(1 - xe^{-x})) = \alpha$$

$$\lim_{x \rightarrow 0} \frac{1}{4x} [2xe^{-2x} + 2xe^{-x}] = \alpha \\ = \frac{1}{4}(4) = \alpha = 1$$

- 16.** Let a vector \vec{a} be coplanar with vectors $\vec{b} = 2\hat{i} + \hat{j} + \hat{k}$ and $\vec{c} = \hat{i} - \hat{j} + \hat{k}$. If \vec{a} is perpendicular to $\vec{d} = 3\hat{i} + 2\hat{j} + 6\hat{k}$, and $|\vec{a}| = \sqrt{10}$. Then a possible value of $[\vec{a}\vec{b}\vec{c}] + [\vec{a}\vec{b}\vec{d}] + [\vec{a}\vec{c}\vec{d}]$ is equal to :

- (1) -40 (2) -42 (3) -29 (4) -38

Sol. (2)

$$\vec{a} = \lambda \vec{b} + \mu \vec{c} = \hat{i}(2\lambda + \mu) + \hat{j}(\lambda - \mu) + \hat{k}(\lambda + \mu)$$

$$\vec{a} \cdot \vec{d} = 0 = 3(2\lambda + \mu) + 2(\lambda - \mu) + 6(\lambda + \mu)$$

$$\Rightarrow 14\lambda + 7\mu = 0 \Rightarrow \mu = -2\lambda$$

$$\Rightarrow \vec{a} = (0)\hat{i} - 3\lambda\hat{j} + (-\lambda)\hat{k}$$

$$\Rightarrow |\vec{a}| = \sqrt{10} \quad |\lambda| = \sqrt{10} \Rightarrow |\lambda| = 1$$

$$\Rightarrow \lambda = 1 \text{ or } -1$$

$$[\vec{a}\vec{b}\vec{c}] = 0 \quad (\text{as vectors are coplanar})$$

$$[\vec{a}\vec{b}\vec{c}] + [\vec{a}\vec{b}\vec{d}] + [\vec{a}\vec{c}\vec{d}] = [\vec{a} \quad \vec{b} + \vec{c} \quad \vec{d}]$$

$$= \begin{vmatrix} 0 & -3\lambda & \lambda \\ 3 & 0 & 2 \\ 3 & 2 & 6 \end{vmatrix}$$

$$= 3\lambda(12) + \lambda(6) = 42\lambda = -42$$

- 17.** Let $y=y(x)$ be the solution of the differential equation

$\operatorname{cosec}^2 x \, dy + 2dx = (1+y \cos 2x) \operatorname{cosec}^2 x \, dx$, with $y\left(\frac{\pi}{4}\right) = 0$. Then, the value of $(y(0)+1)^2$ is equal

to :

- (1) $e^{1/2}$ (2) $e^{-1/2}$ (3) e^{-1} (4) e

Sol. (3)

$$\frac{dy}{dx} + 2 \sin^2 x = 1 + y \cos 2x$$

$$\Rightarrow \frac{dy}{dx} + (-\cos 2x)y = \cos 2x$$

$$\text{I.F.} = e^{\int -\cos 2x \, dx} = e^{-\frac{\sin 2x}{2}}$$

Solution of D.E.

$$y \cdot e^{\frac{-\sin 2x}{2}} = \int (\cos 2x) e^{\frac{-\sin 2x}{2}} dx + c$$

$$\Rightarrow y \left(e^{\frac{-\sin 2x}{2}} \right) = -e^{\frac{-\sin 2x}{2}} + c$$

Given

$$y \left(\frac{\pi}{4} \right) = 0$$

$$\Rightarrow 0 = -e^{-1/2} + c \Rightarrow c = e^{-1/2}$$

$$\Rightarrow y \left(e^{\frac{\sin 2x}{2}} \right) = -e^{\frac{\sin 2x}{2}} + e^{-1/2}$$

at $x = 0$

$$y = -1 + e^{-1/2}$$

$$\Rightarrow y(0) = -1 + e^{-1/2}$$

$$\Rightarrow (y(0) + 1)^2 = e^{-1}$$

- 18.** The value of λ and μ such that the system of equations $x+y+z=6$, $3x+5y+5z=26$, $x + 2y + \lambda z=\mu$ has no solution, are :

(1) $\lambda=3, \mu \neq 10$

(2) $\lambda \neq 2, \mu = 10$

(3) $\lambda = 3, \mu = 5$

(4) $\lambda = 2, \mu \neq 10$

Sol. (4)

$$D = \begin{vmatrix} 1 & 1 & 1 \\ 3 & 5 & 5 \\ 1 & 2 & \lambda \end{vmatrix} = 0$$

$$(5\lambda - 10) - 1(3\lambda - 5) + (6 - 5) = 0$$

$$2\lambda - 10 + 5 + 1 = 0$$

$\boxed{\lambda = 2}$ only one option satisfy

Atleast one If $D_1, D_2, D_3 \neq 0$

- 19.** If the domain of the function $f(x) = \frac{\cos^{-1} \sqrt{x^2 - x + 1}}{\sqrt{\sin^{-1} \left(\frac{2x-1}{2} \right)}}$ is the interval $(\alpha, \beta]$, then $\alpha + \beta$ is equal to :

(1) 2

(2) $\frac{3}{2}$

(3) $\frac{1}{2}$

(4) 1

Sol. (2)

$$0 \leq x^2 - x + 1 \leq 1$$

$$\Rightarrow x^2 - x \leq 0$$

$$\Rightarrow x \in [0, 1]$$

$$\text{Also, } 0 < \sin^{-1} \left(\frac{2x-1}{2} \right) \leq \frac{\pi}{2}$$

$$\Rightarrow 0 < \frac{2x-1}{2} \leq 1$$

$$\Rightarrow 0 < 2x - 1 \leq 2$$

$$1 < 2x \leq 3$$

$$\frac{1}{2} < x \leq \frac{3}{2}$$

Taking intersection

$$x \in \left(\frac{1}{2}, 1 \right]$$

$$\Rightarrow \alpha = \frac{1}{2}, \beta = 1$$

$$\Rightarrow \alpha = \frac{1}{2}, \beta = 1$$

- 20.** If the shortest distance between the straight lines $3(x-1)=6(y-2)=2(z-1)$ and $4(x-2)=2(y-\lambda)=(z-3), \lambda \in \mathbb{R}$ is $\frac{1}{\sqrt{38}}$, then the integral value of λ is equal to :

(1) -1

(2) 2

(3) 3


(4) 5

Sol.

(3)

$$L_1 : \frac{(x-1)}{2} = \frac{(y-2)}{1} = \frac{(z-1)}{3} \quad \vec{r}_1 = 2\hat{i} + \hat{j} + 3\hat{k}$$

$$L_2 : \frac{(x-2)}{1} = \frac{y-\lambda}{2} = \frac{z-3}{4} \quad \vec{r}_2 = \hat{i} + 2\hat{j} + 4\hat{k}$$



Shortest distance = Projection of \vec{a} on $\vec{r}_1 \times \vec{r}_2$

$$= \frac{|\vec{a} \cdot (\vec{r}_1 \times \vec{r}_2)|}{|\vec{r}_1 \times \vec{r}_2|}$$

$$|\vec{a} \cdot (\vec{r}_1 \times \vec{r}_2)| = \begin{vmatrix} 1 & \lambda-2 & 2 \\ 2 & 1 & 3 \\ 1 & 2 & 4 \end{vmatrix} = |14 - 5\lambda|$$

$$|\vec{r}_1 \times \vec{r}_2| = \sqrt{38}$$

$$\therefore \frac{1}{\sqrt{38}} = \frac{|14 - 5\lambda|}{\sqrt{38}}$$

$$\Rightarrow |14 - 5\lambda| = 1$$

$$\Rightarrow 14 - 5\lambda = 1 \text{ or } 14 - 5\lambda = -1$$

$$\Rightarrow \lambda = \frac{13}{5} \text{ or } 3$$

\therefore Integral value of $\lambda = 3$

SECTION B

1. The number of elements in the set $\{n \in \{1, 2, 3, \dots, 100\} \mid (11)^n > (10)^n + (9)^n\}$ is _____.

Sol. (96)

$$11^n > 10^n + 9^n$$

$$\Rightarrow 11^n - 9^n > 10^n$$

$$\Rightarrow (10 + 1)^n - (10 - 1)^n > 10^n$$

$$\Rightarrow \left\{ {}^n C_1 \cdot 10^{n-1} + {}^n C_3 10^{n-3} + {}^n C_5 10^{n-5} + \dots \right\} > 10^n$$

$$\Rightarrow 2n \cdot 10^{n-1} + 2 \left\{ {}^n C_3 10^{n-3} + {}^n C_5 10^{n-5} + \dots \right\} > 10^n \quad \dots(1)$$

For $n = 6, 7, 8, \dots, 100$

$$\Rightarrow 2n 10^{n-1} + 2 \left\{ {}^n C_3 10^{n-3} + {}^n C_5 10^{n-5} + \dots \right\} > 10^n$$

$$\Rightarrow 11^n - 9^n > 10^n \text{ For } n = 5, 6, 7, \dots, 100$$

For $n = 4$, Inequality (1) is not satisfied

\Rightarrow Inequality does not hold good for

$N = 1, 2, 3, 4$

So, required number of elements

= 96

2. Let $y=y(x)$ be the solution of the differential equation

$$\left((x+2)e^{\left(\frac{y+1}{x+2}\right)} + (y+1) \right) dx = (x+2)dy, \quad y(1) = 1.$$

If the domain of $y=y(x)$ is an open interval (α, β) , then $|\alpha + \beta|$ is equal to _____.

Sol. (4)

$$y + 1 = Y \Rightarrow dy = dY$$

$$x + 2 = X \Rightarrow dx = dX$$

$$\Rightarrow \left(Xe^{\frac{Y}{X}} + Y \right) dX = XdY$$

$$\Rightarrow XdY - YdX = Xe^{Y/X}dX$$

$$\Rightarrow d\left(\frac{Y}{X}\right) e^{-\frac{Y}{X}} = \frac{dX}{X}$$

$$-e^{-Y/X} = \ln |X| + c$$

$$(3, 2) \rightarrow -e^{-2/3} = \ln |3| + c$$

$$-e^{-\frac{t}{x}} = \ln |X| - e^{-\frac{2}{3}} - \ln 3$$

$$e^{-\frac{Y}{X}} = e^{2/3} + \ln 3 - \ln |X| > 0$$

$$\ln|x| < e^{2/3} + \ell n 3 \rightarrow \lambda$$

$$|x+2| < e^\lambda$$

$$-e^\lambda < x+2 < e^\lambda$$

$$-e^\lambda - 2 < x < e^\lambda - 2$$

$$\alpha + \beta = -4 \Rightarrow |\alpha + \beta| = 4$$

3. Let $A = \{0, 1, 2, 3, 4, 5, 6, 7\}$. Then the number of bijective functions $f : A \rightarrow A$ such that $f(1) + f(2) = 3 - f(3)$ is equal to _____.

Sol. (720)

$$f(1) + f(2) = 3 - f(3)$$

$$\Rightarrow f(1) + f(2) + f(3) = 3$$

$$\text{The only possibility is: } 0 + 1 + 2 = 3$$

\Rightarrow Elements 1, 2, 3 in the domain can be mapped with 0, 1, 2 only.

So number of bijective functions.

$$= |3 \times 5| = 720$$

4. If the constant term, in binomial expansion of $\left(2x^r + \frac{1}{x^2}\right)^{10}$ is 180, than r is equal to _____.

Sol. (8)

$$\left(2x^r + \frac{1}{x^2}\right)^{10}$$

$$\text{General term} = {}^{10}C_R \left(2x^r\right)^{10-R} x^{-2R}$$

$$\Rightarrow 2^{10-R} {}^{10}C_R = 180 \dots\dots\dots (1)$$

$$\& (10-R)r - 2R = 0$$

$$r = \frac{2R}{10-R}$$

$$r = \frac{2(R-10)}{10-R} + \frac{20}{10-R}$$

$$\Rightarrow r = -2 + \frac{20}{10-R} \dots\dots (2)$$

$R = 8$ or 5 reject equation (1) not satisfied

At $R = 8$

$$2^{10-R} {}^{10}C_R = 180 \Rightarrow r = 8$$

5. Let $f : \mathbf{R} \rightarrow \mathbf{R}$ be function defined as $f(x) = \begin{cases} 3\left(1 - \frac{|x|}{2}\right) & \text{if } |x| \leq 2 \\ 0 & \text{if } |x| > 2 \end{cases}$


Let $g : \mathbf{R} \rightarrow \mathbf{R}$ be given by $g(x) = f(x+2) - f(x-2)$. If n and m denote the number of points in \mathbf{R} where g is not continuous and not differentiable, respectively, then $n+m$ is equal to _____.

Sol. (4)

$$f(x-2) = \begin{cases} \frac{3x}{2} & -4 \leq x \leq -2 \\ -\frac{3x}{2} & -2 < x \leq 0 \\ 0 & x \in (-\infty, -4) \cup (0, +\infty) \end{cases}$$

$$f(x+2) = \begin{cases} \frac{3x}{2} + 6 & 0 \leq x \leq 2 \\ -\frac{3x}{2} + 6 & 2 < x \leq 4 \\ 0 & x \in (-\infty, 0) \cup (4, +\infty) \end{cases}$$

$$g(x) = f(x+2) - f(x-2) = \begin{cases} \frac{3x}{2} + 6 & -4 \leq x \leq -2 \\ -\frac{3x}{2} & -2 < x < 2 \\ \frac{3x}{2} - 6 & 2 \leq x \leq 4 \\ 0 & x \in (-\infty, -4) \cup (4, +\infty) \end{cases}$$



$$n = 0$$

$$m = 4 \Rightarrow (n + m = 4)$$

6. Let $A = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$. Then the number of 3×3 matrices B with entries from the set $\{1, 2, 3, 4, 5\}$ and satisfying $AB=BA$ is _____.

Sol. (3125)

$$\text{Let matrix } B = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

$$\therefore AB = BA$$

$$\begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} d & e & f \\ a & b & c \\ g & h & i \end{bmatrix} = \begin{bmatrix} b & a & c \\ e & d & f \\ h & g & i \end{bmatrix}$$

$$\Rightarrow d = b, e = a, f = c, g = h$$

$$\therefore \text{Matrix } B = \begin{bmatrix} a & b & c \\ b & a & c \\ g & g & i \end{bmatrix}$$

No. of ways of selecting a, b, c, g,

$$i = 5 \times 5 \times 5 \times 5 \times 5$$

$$= 5^5 = 3125$$

$$\therefore \text{No. of Matrices } B = 3125$$

7. Consider the following frequency distribution :

Class :	0 – 6	6 – 12	12 – 18	18 – 24	24 – 30
Frequency :	a	b	12	9	5

$$\text{If mean} = \frac{309}{22} \text{ and median} = 14, \text{ then value } (a - b)^2 \text{ is equal to } \underline{\hspace{2cm}}.$$

Sol. (4)

Class	Frequency	X _i	F _i x _i
0-6	a	3	3a
6-12	b	9	9b
12-18	12	15	180
18-24	9	21	189
24-30	5	27	135
	N=(26+a+b)		(504+3a+9b)

$$\text{Mean} = \frac{3a + 9b + 180 + 189 + 135}{a + b + 26} = \frac{309}{22}$$

$$\Rightarrow 66a + 198b + 11088 = 309a + 309b + 8034$$

$$\Rightarrow 243a + 111b = 3054$$

$$\Rightarrow 81a + 37b = 1018 \quad -(1)$$

$$\text{Now, Median} = 12 + \frac{\frac{a+b+26}{2} - (a+b)}{12} \times 6 = 14$$

$$\Rightarrow \frac{13}{2} - \left(\frac{a+b}{4} \right) = 2$$

$$\Rightarrow \frac{a+b}{4} = \frac{9}{2}$$

$$\Rightarrow a + b = 18 \rightarrow (2)$$

From equation (1) & (2)

$$a = 8, b = 10$$

$$\therefore (a - b)^2 = (8 - 10)^2$$

- 8.** If the digits are not allowed to repeat in any number formed by using the digits 0, 2, 4, 6, 8, then the number of all numbers greater than 10,000 is equal to _____.

Sol. (96)

$$\begin{array}{|c|c|c|c|c|} \hline & 0,2,4,6,8 \\ \hline 2,4,6,8 & & & & \\ \hline 4 & 4 & 3 & 2 & 1 \\ \hline \end{array}$$

$$\Rightarrow 4 \times 4 \times 3 \times 2 = 96$$

- 9.** The sum of all the elements in the set $\{n \in \{1, 2, \dots, 100\} \mid H.C.F. \text{ of } n \text{ and } 2040 \text{ is } 1\}$ is equal to _____.

Sol. (1251)

$$2040 = 2^3 \times 3 \times 5 \times 17$$


n should not be multiple of 2, 3, 5 and 17.

$$\begin{aligned}
 \text{Sum of all } n &= (1 + 3 + 5 + \dots + 99) - (3 + 9 + 15 + 21 + \dots + 99) - (5 + 25 + 35 + 55 + 65 \\
 &\quad + 85 + 95) - (17) \\
 &= 2500 - \frac{17}{2}(3 + 99) - 365 - 17 \\
 &= 2500 - 867 - 365 - 17 = 1251
 \end{aligned}$$

- 10.** The area (in sq. units) of the region bounded by the curves $x^2 + 2y - 1 = 0$, $y^2 + 4x - 4 = 0$ and $y^2 - 4x - 4 = 0$, in the upper half plane is _____.

Sol. (2)

Required Area (shaded)



$$\begin{aligned}
 &= 2 \left[\int_0^2 \left(\frac{4 - y^2}{4} \right) dy - \int_0^1 \left(\frac{1 - x^2}{2} \right) dx \right] \\
 &= 2 \left[\frac{4}{3} - \frac{1}{3} \right] = (2)
 \end{aligned}$$

अब मोशन ही है, सर्वश्रेष्ठ विकल्प!

MOTION welcomes

Directors of Nucleus Education & Wizard of Mathematics

Now Offline associated with Motion Kota Classroom



Nitin Vijay (NV Sir)
Managing Director
Exp. : 18 yrs



Akhilesh Kanther
(AKK Sir)
Exp. : 17 yrs



Vishal Joshi
(VJ Sir)
Exp. : 18 yrs



Surendra K. Mishra
(SKM Sir)
Exp. : 16 yrs



Gavesh Bhardwaj
(GB Sir)
Exp. : 17 yrs

Academic Pillars of JEE Motion Kota



Ram Ratan Dwivedi
(RRD Sir)
Joint Director
Exp. : 20 yrs



Amit Verma
(AV Sir)
Joint Director
Exp. : 16 yrs



Vijay Pratap Singh
(VPS Sir)
Vice President
Exp. : 20 yrs



Nikhil Srivastava
(NS Sir)
Head JEE Academics
Exp. : 17 yrs



Aatish Agarwal
(AA Sir)
Sr. Faculty
Exp. : 17 yrs



Jayant Chittora
(JC Sir)
Sr. Faculty
Exp. : 16 yrs



Anurag Garg
(AG Sir)
Sr. Faculty
Exp. : 17 yrs



Arjun Gupta
(Arjun Sir)
Sr. Faculty
Exp. : 14 yrs



Devki Nandan Pathak
(DN Sir)
Sr. Faculty
Exp. : 13 yrs



Avinash Kishore
(AVN Sir)
Sr. Faculty
Exp. : 9 yrs



Vipin Sharma
(VS Sir)
Sr. Faculty
Exp. : 12 yrs



Durgesh Pandey
(Pandey Sir)
Sr. Faculty
Exp. : 8 yrs

Join JEE DROPPER BATCH Kota Classroom

English & Hindi Medium

Online + Offline Mode

Batch Starting from: **4th August 2021**

Time to use all your skills, Now!

CRASH COURSE JEE Advanced 2021

Starting from: **9th August'21**

Why should you choose?

- ◆ Live Lectures
- ◆ Best JEE Faculties
- ◆ Doubt Support on learning app
- ◆ Online test with discussion
- ◆ Personalized performance analysis

