



**JEE  
MAIN  
July  
2021**

**MATHS**  
**27<sup>th</sup> July 2021 [SHIFT – 1]**  
**QUESTION WITH SOLUTION**

**JEE | NEET | Foundation**

**MOTION**<sup>TM</sup>

**25000+**  
SELECTIONS SINCE 2007

अब मोशन ही है, सर्वश्रेष्ठ विकल्प!

MOTION welcomes

Directors of Nucleus Education & Wizard of Mathematics  
Now Offline associated with Motion Kota Classroom



Nitin Vijay (NV Sir)  
Managing Director  
Exp. : 18 yrs



Akhilesh Kanther  
(AKK Sir)  
Exp. : 17 yrs



Vishal Joshi  
(VJ Sir)  
Exp. : 18 yrs



Surendra K. Mishra  
(SKM Sir)  
Exp. : 16 yrs



Gavesh Bhardwaj  
(GB Sir)  
Exp. : 17 yrs

## Academic Pillars of JEE Motion Kota



Ram Ratan Dwivedi  
(RRD Sir)  
Joint Director  
Exp. : 20 yrs



Amit Verma  
(AV Sir)  
Joint Director  
Exp. : 16 yrs



Vijay Pratap Singh  
(VPS Sir)  
Vice President  
Exp. : 20 yrs



Nikhil Srivastava  
(NS Sir)  
Head JEE Academics  
Exp. : 17 yrs



Aatish Agarwal  
(AA Sir)  
Sr. Faculty  
Exp. : 17 yrs



Jayant Chittora  
(JC Sir)  
Sr. Faculty  
Exp. : 16 yrs



Anurag Garg  
(AG Sir)  
Sr. Faculty  
Exp. : 17 yrs



Arjun Gupta  
(Arjun Sir)  
Sr. Faculty  
Exp. : 14 yrs



Devki Nandan Pathak  
(DN Sir)  
Sr. Faculty  
Exp. : 13 yrs



Avinash Kishore  
(AVN Sir)  
Sr. Faculty  
Exp. : 9 yrs



Vipin Sharma  
(VS Sir)  
Sr. Faculty  
Exp. : 12 yrs



Durgesh Pandey  
(Pandey Sir)  
Sr. Faculty  
Exp. : 8 yrs

Join **JEE DROPPER BATCH** Kota Classroom

English & Hindi Medium

Online + Offline Mode

Batch Starting from: **4th August 2021**

Time to use all your skills, Now!

## CRASH COURSE

### JEE Advanced 2021

Starting from: **9th August '21**

### Why should you choose?

- ◆ Live Lectures
- ◆ Best JEE Faculties
- ◆ Doubt Support on learning app
- ◆ Online test with discussion
- ◆ Personalized performance analysis



### SECTION - A

1. If the area of the bounded region


$$R = \left\{ (x, y) : \max\{0, \log_e x\} \leq y \leq 2^x, \frac{1}{2} \leq x \leq 2 \right\}$$

is,  $\alpha (\log_e 2)^{-1} + \beta (\log_e 2) + \gamma$ , then the value of  $(\alpha + \beta - 2\gamma)^2$  is equal to:

- (1) 4                      (2) 1                      (3) 8                      (4) 2

Sol. (4)

$$R \left\{ (x, y) : \max(0, \log_e x) \leq y \leq 2^x, \frac{1}{2} \leq x \leq 2 \right\}$$



$$\int_{\frac{1}{2}}^2 2^x dx - \int_1^2 \ln x dx$$

$$\Rightarrow \left[ \frac{2^x}{\ln 2} \right]_{\frac{1}{2}}^2 - [x \ln x - x]_1^2$$

$$\Rightarrow \frac{(2^2) - 2^{1/2}}{\log_e 2} - (2 \ln 2 - 1)$$

$$\therefore \alpha = 2^2 - \sqrt{2}, \beta = -2, \gamma = 1$$

$$\Rightarrow (\alpha + \beta - 2\gamma)^2$$

$$\Rightarrow (2^2 - \sqrt{2} - 2 - 2)^2$$

$$\Rightarrow (\sqrt{2})^2 = 2$$

2. Let  $\alpha, \beta$  be two roots of the equation  $x^2 + (20)^{\frac{1}{4}}x + (5)^{\frac{1}{2}} = 0$ . Then  $\alpha^8 + \beta^8$  is equal to:

- (1) 10                      (2) 50                      (3) 160                      (4) 100

Sol. (2)

$$(x^2 + \sqrt{5})^2 = \sqrt{20}x^2$$


$$x^4 = -5 \Rightarrow x^8 = 25$$

$$\alpha^8 + \beta^8 = 50$$

3. Let P and Q be two distinct points on a circle which has center at C(2,3) and which passes through origin O, If OC is perpendicular to both the line segments CP and CQ, then the set {P,Q} is equal to:

- (1)  $\{(-1,5), (5,1)\}$   
 (2)  $\{(2+2\sqrt{2}, 3-\sqrt{5}), (2-2\sqrt{2}, 3+\sqrt{5})\}$   
 (3)  $\{(2+2\sqrt{2}, 3+\sqrt{5}), (2-2\sqrt{2}, 3-\sqrt{5})\}$   
 (4)  $\{(4,0), (0,6)\}$

Sol. (1)



$$\tan \theta = -\frac{2}{3}$$

Using symmetric form of line

$$P, Q : (2 \pm \sqrt{13} \cos \theta, 3 \pm \sqrt{13} \sin \theta)$$

$$\left( 2 \pm \sqrt{13} \cdot \left( -\frac{3}{\sqrt{13}} \right), 3 \pm \sqrt{13} \left( \frac{2}{\sqrt{13}} \right) \right)$$

$$(-1,5) \& (5,1)$$

4. Let  $y = y(x)$  be solution of the differential equation  $\log_e \left( \frac{dy}{dx} \right) = 3x + 4y$ , with  $y(0) = 0$ . If

$$y \left( -\frac{2}{3} \log_e 2 \right) = \alpha \log_e 2, \text{ then the value of } \alpha \text{ is equal to:}$$

- (1)  $-\frac{1}{2}$                       (2)  $-\frac{1}{4}$                       (3) 2                      (4)  $\frac{1}{4}$

Sol. (2)

$$\frac{dy}{dx} = e^{3x} \cdot e^{4y} \Rightarrow \int e^{-4y} dy = \int e^{3x} dx$$

$$\frac{e^{-4y}}{-4} = \frac{e^{3x}}{3} + C \Rightarrow -\frac{1}{4} - \frac{1}{3} = C \Rightarrow C = -\frac{7}{12}$$

$$\frac{e^{-4y}}{-4} = \frac{e^{3x}}{3} - \frac{7}{12} \Rightarrow e^{-4y} = \frac{4e^{3x} - 7}{-3}$$

$$e^{4y} = \frac{3}{7 - 4e^{3x}} \Rightarrow 4y = \ln \left( \frac{3}{7 - 4e^{3x}} \right)$$

$$4y = \ln \left( \frac{3}{6} \right) \text{ when } x = -\frac{2}{3} \ln 2$$

$$y = \frac{1}{4} \ln \left( \frac{1}{2} \right) = -\frac{1}{4} \ln 2$$

5. The compound statement  $(P \vee Q) \wedge (\sim P) \Rightarrow Q$  is equivalent to:  
 (1)  $P \vee Q$  (2)  $\sim(P \Rightarrow Q) \Leftrightarrow P \wedge \sim Q$   
 (3)  $P \wedge \sim Q$  (4)  $\sim(P \Rightarrow Q)$

**Sol. (2)**

Using Truth Table

P	Q	$P \vee Q$	$\sim P$	$(P \vee Q) \wedge P$	$(P \vee Q) \wedge \sim P \rightarrow Q$
T	T	T	F	F	T
T	F	T	F	F	T
F	T	T	F	T	T
F	F	F	T	F	T

Check from options

6. Let  
 $A = \{(x,y) \in \mathbb{R} \times \mathbb{R} \mid 2x^2 + 2y^2 - 2x - 2y = 1\}$ ,  
 $B = \{(x,y) \in \mathbb{R} \times \mathbb{R} \mid 4x^2 + 4y^2 - 16y + 7 = 0\}$  and  
 $C = \{(x,y) \in \mathbb{R} \times \mathbb{R} \mid x^2 + y^2 - 4x - 2y + 5 \leq r^2\}$ .  
 Then the minimum value of  $|r|$  such that  $A \cup B \subseteq C$  is equal to:

- (1)  $\frac{3 + \sqrt{10}}{2}$  (2)  $1 + \sqrt{5}$  (3)  $\frac{2 + \sqrt{10}}{2}$  (4)  $\frac{3 + 2\sqrt{5}}{2}$

**Sol. (4)**

$$S_1 : x^2 + y^2 - x - y - \frac{1}{2} = 0; C_1 \left( \frac{1}{2}, \frac{1}{2} \right)$$

$$r_1 = \sqrt{\frac{1}{4} + \frac{1}{4} + \frac{1}{2}} = 1$$


$$S_2 : x^2 + y^2 - 4y + \frac{7}{4} = 0; C_2 : (0, 2)$$

$$r_2 = \sqrt{4 - \frac{7}{4}} = \frac{3}{2}$$

$$S_3 : x^2 + y^2 - 4x - 2y + 5 - r^2 = 0$$

$$C_3 : (2, 1)$$

$$r_3 = \sqrt{4 + 1 - 5 + r^2} = |r|$$



$$C_1 C_3 = \sqrt{\frac{5}{2}}$$

$$\sqrt{\frac{5}{2}} \leq |r-1| \Rightarrow \left. \begin{array}{l} r \leq 1 + \sqrt{\frac{5}{2}} \\ r \geq \frac{3}{2} + \sqrt{5} \end{array} \right\}$$

$$C_2 C_3 = \sqrt{5} \leq \left| r - \frac{3}{2} \right|$$

$$\left. \begin{array}{l} r - \frac{3}{2} \geq \sqrt{5} \\ r - \frac{3}{2} \leq -\sqrt{5} \end{array} \right\}$$

7. The probability that a randomly selected 2 digit number belongs to the set  $(n \in \mathbb{N}; (2^n - 2) \text{ is a multiple of } 3)$  is equal to:

(1)  $\frac{1}{2}$                       (2)  $\frac{1}{3}$                       (3)  $\frac{2}{3}$                       (4)  $\frac{1}{6}$

**Sol. (1)**

Total number of cases =  ${}^{90}C_1 = 90$

Now,  $2^n - 2 = (3 - 1)^n - 2$

$${}^n C_0 3^n - {}^n C_1 \cdot 3^{n-1} + \dots + (-1)^{n-1} \cdot {}^n C_{n-1} 3 + (-1)^n \cdot {}^n C_n - 2$$

$$3(3^{n-1} - n3^{n-2} + \dots + (-1)^{n-1} \cdot n) + (-1)^n - 2$$

$(2^n - 2)$  is multiply of 3 only when n is odd

Req. Probability =  $\frac{45}{90} = \frac{1}{2}$

8. Let  $A = \begin{bmatrix} 1 & 2 \\ -1 & 4 \end{bmatrix}$ . If  $A^{-1} = \alpha I + \beta A$ ,  $\alpha, \beta \in \mathbb{R}$ , I is a  $2 \times 2$  identity matrix, then  $4(\alpha - \beta)$  is equal to:

(1) 5                      (2) 4                      (3) 2                      (4)  $\frac{8}{3}$

**Sol. (2)**

$$A = \begin{bmatrix} 1 & 2 \\ -1 & 4 \end{bmatrix}, |A| = 6$$

$$A^{-1} = \frac{\text{adj}A}{|A|} = \frac{1}{6} \begin{bmatrix} 4 & -2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} \frac{2}{3} & -\frac{1}{3} \\ \frac{1}{6} & \frac{1}{6} \end{bmatrix}$$

$$\begin{bmatrix} \frac{2}{3} & -\frac{1}{3} \\ \frac{1}{6} & \frac{1}{6} \end{bmatrix} = \begin{bmatrix} \alpha & 0 \\ 0 & \alpha \end{bmatrix} + \begin{bmatrix} \beta & 2\beta \\ -\beta & 4\beta \end{bmatrix}$$

$$\left. \begin{array}{l} \alpha + \beta = \frac{2}{3} \\ \beta = -\frac{1}{6} \end{array} \right\} \Rightarrow \alpha = \frac{2}{3} + \frac{1}{6} = \frac{5}{6}$$

$4(\alpha - \beta) = 4(1) = 4$

9. If the mean and variance of the following data:  
6, 10, 7, 13, a, 12, b, 12 are 9 and  $\frac{37}{4}$  respectively, then  $(a-b)^2$  is equal to:
- (1) 12                      (2) 24                      (3) 16                      (4) 32

**Sol. (3)**

$$\text{Mean} = \frac{6+10+7+13+a+12+b+12}{8} = 9$$

$$60 + a + b = 72$$

$$a + b = 12 \quad \dots(1)$$

$$\text{variance} = \frac{\sum x_i^2}{n} - \left(\frac{\sum x_i}{n}\right)^2 = \frac{37}{4}$$

$$\sum x_i^2 = 6^2 + 10^2 + 7^2 + 13^2 + a^2 + b^2 + 12^2 + 12^2$$

$$= a^2 + b^2 + 642$$

$$\frac{a^2 + b^2 + 642}{8} - (9)^2 = \frac{37}{4}$$

$$\frac{a^2 + b^2}{8} + \frac{321}{4} - 81 = \frac{37}{4}$$

$$\frac{a^2 + b^2}{8} = 81 + \frac{37}{4} - \frac{321}{4}$$

$$\frac{a^2 + b^2}{8} = 81 - 71$$

$$\therefore a^2 + b^2 = 80 \quad \dots(2)$$

$$\text{From (1) } a^2 + b^2 + 2ab = 144$$

$$80 + 2ab = 144 \therefore 2ab = 64$$

$$(a - b)^2 = a^2 + b^2 - 2ab = 80 - 64 = 16$$

10. A ray of light through (2,1) is reflected at a point P on the y - axis and then passes through the point (5,3). If this reflected ray is the directrix of an ellipse with eccentricity  $\frac{1}{3}$  and the distance of the nearer focus from this directrix is  $\frac{8}{\sqrt{53}}$ , then the equation of the other directrix can be :


(1)  $2x - 7y - 39 = 0$  or  $2x - 7y - 7 = 0$

(2)  $11x + 7y + 8 = 0$  or  $11x + 7y - 15 = 0$

(3)  $2x - 7y + 29 = 0$  or  $2x - 7y - 7 = 0$

(4)  $11x - 7y - 8 = 0$  or  $11x + 7y + 15 = 0$

**Sol. (3)**



Equation of reflected Ray

$$y - 1 = \frac{2}{7}(x + 2)$$

$$7y - 7 = 2x + 4$$

$$2x - 7y + 11 = 0$$

Let the equation of other directrix is

$$2x - 7y + \lambda$$

Distance of directrix from Focub

$$\frac{a}{e} - ae = \frac{8}{\sqrt{53}}$$

$$3a - \frac{a}{3} = \frac{8}{\sqrt{53}} \text{ or } a = \frac{3}{\sqrt{53}}$$

$$\text{Distance between two directric} = \frac{2a}{e}$$

$$= 2 \times 3 \times \frac{3}{\sqrt{53}} = \frac{18}{\sqrt{53}}$$

$$\left| \frac{\lambda - 11}{\sqrt{53}} \right| = \frac{18}{\sqrt{53}}$$

$$\lambda - 11 = 18 \text{ or } -18$$

$$\lambda = 29 \text{ or } -7$$

$$2x - 7y - 7 = 0 \text{ or } 2x - 7y + 29 = 0$$

11. Let  $f: \left(-\frac{\pi}{4}, \frac{\pi}{4}\right) \rightarrow \mathbb{R}$  be defined as

$$f(x) = \begin{cases} (1 + |\sin x|)^{\frac{3a}{|\sin x|}}, & -\frac{\pi}{4} < x < 0 \\ b, & x = 0 \\ e^{\cot 4x / \cot 2x}, & 0 < x < \frac{\pi}{4} \end{cases}$$

If  $f$  is continuous at  $x = 0$ , then the value of  $6a + b^2$  is equal to:

- (1)  $e$                       (2)  $1+e$                       (3)  $1-e$                       (4)  $e-1$

Sol. (2)

$$\lim_{x \rightarrow 0} f(x) = b$$

$$\lim_{x \rightarrow 0^+} e^{\frac{\cot 4x}{\cot 2x}} = e^{\frac{1}{2}} = b$$

$$\lim_{x \rightarrow 0^-} (1 + |\sin x|)^{\frac{3a}{|\sin x|}} = e^{3a} = e^{\frac{1}{2}}$$

$$a = \frac{1}{6} \Rightarrow 6a = 1$$

$$(6a + b^2) = (1 + e)$$



- 12.** If  $\sin\theta + \cos\theta = \frac{1}{2}$ , then  $16(\sin(2\theta) + \cos(4\theta) + \sin(6\theta))$  is equal to:  
 (1) 27                      (2) -27                      (3) -23                      (4) 23

**Sol. (3)**

$$16[2\sin 4\theta \cos 2\theta + \cos 4\theta]$$

$$16[4\sin 2\theta \cos^2 2\theta + 2\cos^2 2\theta - 1]$$

Now:

$$\sin\theta + \cos\theta = \frac{1}{2}$$

$$1 + \sin 2\theta = \frac{1}{4}$$

$$\sin 2\theta = -\frac{3}{4}$$

$$\cos^2 2\theta = 1 - \frac{9}{16} = \frac{7}{16}$$

$$16\left[-4\left(-\frac{3}{4}\right) \times \frac{7}{16} + 2 \times \frac{7}{16} - 1\right]$$

$$16\left[\frac{-7}{16} - 1\right] \Rightarrow -23$$

- 13.** Let C be the set of all complex numbers. Let  
 $S_1 = \{z \in \mathbb{C} \mid |z - 3 - 2i|^2 = 8\}$ ,  
 $S_2 = \{z \in \mathbb{C} \mid \operatorname{Re}(z) \geq 5\}$  and  
 $S_3 = \{z \in \mathbb{C} \mid |z - \bar{z}| \geq 8\}$ .  
 Then the number of elements in  $S_1 \cap S_2 \cap S_3$  is equal to:  
 (1) 1                      (2) 0                      (3) Infinite                      (4) 2

**Sol. (1)**

$$S_1 : |x - 3 - 2i|^2 = 8$$

$$|x - 3 - 2i| = 2\sqrt{2}$$


$$(x - 3)^2 + (y - 2)^2 = (2\sqrt{2})^2$$

$$S_2 : x \geq 5$$

$$S_3 : |z - \bar{z}| \geq 8$$

$$|2iy| \geq 8$$

$$2|y| \geq 8 \therefore y \geq 4, y \leq -4$$



$$n(S_1 \cap S_2 \cap S_3) = 1$$

- 14.** Let  $\vec{a} = \hat{i} + \hat{j} + 2\hat{k}$  and  $\vec{b} = -\hat{i} + 2\hat{j} + 3\hat{k}$ . Then the vector product  $(\vec{a} + \vec{b}) \times ((\vec{a} \times ((\vec{a} - \vec{b}) \times \vec{b})) \times \vec{b})$  is equal to:
- (1)  $5(30\hat{i} - 5\hat{j} + 7\hat{k})$  (2)  $5(34\hat{i} - 5\hat{j} + 3\hat{k})$   
 (3)  $7(30\hat{i} - 5\hat{j} + 7\hat{k})$  (4)  $7(34\hat{i} - 5\hat{j} + 3\hat{k})$

**Sol. (4)**

$$\vec{a} = \hat{i} + \hat{j} + 2\hat{k}$$

$$\vec{b} = -\hat{i} + 2\hat{j} + 3\hat{k}$$

$$\vec{a} + \vec{b} = 3\hat{j} + 5\hat{k}; \vec{a} \cdot \vec{b} = -1 + 2 + 6 = 7$$

$$((\vec{a} \times ((\vec{a} - \vec{b}) \times \vec{b})) \times \vec{b})$$

$$((\vec{a} \times (\vec{a} \times \vec{b} - \vec{b} \times \vec{b})) \times \vec{b})$$

$$(\vec{a} \times (\vec{a} \times \vec{b} - 0)) \times \vec{b}$$

$$(\vec{a} \times (\vec{a} \times \vec{b})) \times \vec{b}$$

$$((\vec{a} \times \vec{b})\vec{a} - (\vec{a} \cdot \vec{a})\vec{b}) \times \vec{b}$$

$$(\vec{a} \cdot \vec{b})\vec{a} \times \vec{b} - (\vec{a} \cdot \vec{a})(\vec{b} \times \vec{b})$$

$$(\vec{a} \cdot \vec{b})(\vec{a} \times \vec{b})$$

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 2 \\ -1 & 2 & 3 \end{vmatrix} = -\hat{i} - 5\hat{j} + 3\hat{k}$$

$$\therefore 7(-\hat{i} - 5\hat{j} + 3\hat{k})$$

$$(\vec{a} + \vec{b}) \times (7(-\hat{i} - 5\hat{j} + 3\hat{k}))$$

$$7(0\hat{i} + 3\hat{j} + 5\hat{k}) \times (-\hat{i} - 5\hat{j} + 3\hat{k})$$

$$\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & 3 & 5 \\ -1 & -5 & 3 \end{vmatrix}$$

$$\Rightarrow 34\hat{i} - (5)\hat{j} + (3\hat{k})$$

$$\Rightarrow 34\hat{i} - 5\hat{j} + 3\hat{k}$$

$$\therefore 7(34\hat{i} - 5\hat{j} + 3\hat{k})$$

15. The value of the definite integral

$$\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{dx}{(1 + e^{x \cos x})(\sin^4 x + \cos^4 x)}$$

is equal to:

(1)  $\frac{\pi}{\sqrt{2}}$

(2)  $-\frac{\pi}{4}$

(3)  $\frac{\pi}{2\sqrt{2}}$

(4)  $-\frac{\pi}{2}$

Sol. (3)

$$I = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{dx}{(1 + e^{x \cos x})(\sin^4 x + \cos^4 x)} \dots (1)$$

Using  $\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$

$$I = \int_{-\pi/4}^{\pi/4} \frac{dx}{(1 + e^{-x \cos x})(\sin^4 x + \cos^4 x)}$$

Add (1) and (2)

$$2I = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{dx}{\sin^4 x + \cos^4 x}$$

$$2I = 2 \int_0^{\frac{\pi}{4}} \frac{dx}{\sin^4 x + \cos^4 x}$$

$$I = \int_0^{\frac{\pi}{4}} \frac{(1 + \tan^2 x) \sec^2 x}{\tan^4 x + 1} dx$$

$$I = \int_0^{\frac{\pi}{4}} \frac{\left(1 + \frac{1}{\tan^2 x}\right) \sec^2 x}{\left(\tan x - \frac{1}{\tan x}\right)^2 + 2} dx$$

$$\tan x - \frac{1}{\tan x} = t$$

$$\left(1 + \frac{1}{\tan^2 x}\right) \sec^2 x dx = dt$$

$$I = \int_{-\infty}^0 \frac{dt}{t^2 + 2} = \left[ \frac{1}{\sqrt{2}} \tan^{-1} \left( \frac{t}{\sqrt{2}} \right) \right]_{-\infty}^0$$

$$I = 0 - \frac{1}{\sqrt{2}} \left( -\frac{\pi}{2} \right) = \frac{\pi}{2\sqrt{2}}$$

- 16.** Let the plane passing through the point  $(-1, 0, -2)$  and perpendicular to each of the planes  $2x + y - z = 2$  and  $x - y - z = 3$  be  $ax + by + cz + 8 = 0$ . then the value of  $a + b + c$  is equal to:  
 (1) 8                                      (2) 4                                      (3) 3                                      (4) 5

**Sol. (2)**

$$\text{Normal of req. plane } (2\hat{i} + \hat{j} - \hat{k}) \times (\hat{i} - \hat{j} - \hat{k})$$

$$= -2\hat{i} + \hat{j} - 3\hat{k}$$

Equation of plane

$$-2(x + 1) + 1(y - 0) - 3(z + 2) = 0$$

$$-2x + y - 3z - 8 = 0$$

$$2x - y + 3z + 8 = 0$$

$$a + b + c = 4$$

- 17.** The value of  $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \frac{(2j-1) + 8n}{(2j-1) + 4n}$  is equal to:

(1)  $2 - \log_e \left( \frac{2}{3} \right)$                       (2)  $3 + 2 \log_e \left( \frac{2}{3} \right)$                       (3)  $1 + 2 \log_e \left( \frac{3}{2} \right)$                       (4)  $5 + \log_e \left( \frac{3}{2} \right)$

**Sol. (3)**

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \frac{\left( \frac{2j}{n} - \frac{1}{n} + 8 \right)}{\left( \frac{2j}{n} - \frac{1}{n} + 4 \right)}$$

$$\int_0^1 \frac{2x + 8}{2x + 4} dx = \int_0^1 dx + \int_0^1 \frac{4}{2x + 4} dx$$

$$= 1 + 4 \frac{1}{2} (\ln|2x + 4|)_0^1$$

$$1 + 2 \ln \left( \frac{3}{2} \right)$$

**18.** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $f(2)=4$  and  $f'(2) = 1$ . Then, the value of  $\lim_{x \rightarrow 2} \frac{x^2 f(2) - 4f(x)}{x - 2}$

is equal to:

- (1) 4                                      (2) 8                                      (3) 16                                      (4) 12

**Sol. (4)**

Apply L'Hopital Rule

$$\lim_{x \rightarrow 2} \left( \frac{2xf(2) - 4f'(x)}{1} \right)$$

$$= \frac{4(4) - 4}{1} = 12$$

**19.** If the coefficients of  $x^7$  in  $\left(x^2 + \frac{1}{bx}\right)^{11}$  and  $x^{-7}$  in  $\left(x - \frac{1}{bx^2}\right)^{11}$ ,  $b \neq 0$ , are equal, then the value of

$b$  is equal to:

- (1) -1                                      (2) 2                                      (3) -2                                      (4) 1

**Sol. (4)**

Coefficient of  $x^7$  in  $\left(x^2 + \frac{1}{bx}\right)^{11}$

$${}^{11}C_r (x^2)^{11-r} \cdot \left(\frac{1}{bx}\right)^r$$

$${}^{11}C_r x^{22-3r} \cdot \frac{1}{b^r}$$

$$22 - 3r = 7$$

$$r = 5$$

$$\therefore {}^{11}C_5 \cdot \frac{1}{b^5} \cdot x^7$$

Coefficient of  $x^{-7}$  in  $\left(x - \frac{b}{bx^2}\right)^{11}$

$${}^{11}C_r (x)^{11-r} \cdot \left(-\frac{1}{bx^2}\right)^r$$

$${}^{11}C_r x^{11-3r} \cdot \frac{(-1)^r}{b^r}$$

$$11 - 3r = -7 \therefore r = 6$$

$${}^{11}C_6 \cdot \frac{1}{b^6} x^{-7}$$


$${}^{11}C_5 \cdot \frac{1}{b^5} = {}^{11}C_6 \cdot \frac{1}{b^6}$$

Since  $b \neq 0 \therefore b = 1$

20. Two tangents are drawn from the point  $P(-1,1)$  to the circle  $x^2 + y^2 - 2x - 6y + 6 = 0$ . If these tangents touch the circle at points  $A$  and  $B$ , and if  $D$  is a point on the circle such that length of the segments  $AB$  and  $AD$  are equal, then the area of the triangle  $ABD$  is equal to:

- (1) 2                      (2)  $(3\sqrt{2} + 2)$                       (3) 4                      (4)  $3(\sqrt{2} - 1)$

Sol. (3)



$$\Delta ABD = \frac{1}{2} \times 2 \times 4$$

$$= 4$$

### SECTION B

1.  $f(x) = \begin{vmatrix} \sin^2 x & -2 + \cos^2 x & \cos 2x \\ 2 + \sin^2 x & \cos^2 x & \cos 2x \\ \sin^2 x & \cos^2 x & 1 + \cos 2x \end{vmatrix}, x \in [0, \pi]$

Then the maximum value of  $f(x)$  is equal to\_\_\_\_\_.

Sol. (6)

$$\begin{vmatrix} -2 & -2 & 0 \\ 2 & 0 & -1 \\ \sin^2 x & \cos^2 x & 1 + \cos 2x \end{vmatrix} \begin{matrix} (R_1 \rightarrow R_1 - R_2) \\ \& R_2 \rightarrow R_2 - R_3 \end{matrix}$$

$$-2(\cos^2 x) + 2(2 + 2 \cos 2x + \sin^2 x)$$

$$4 + 4 \cos 2x - 2(\cos^2 x - \sin^2 x)$$

$$f(x) = 4 + \frac{2 \cos 2x}{\max=1}$$

$$f(x)_{\max} = 4 + 2 = 6$$

2. Let  $F: [3,5] \rightarrow \mathbb{R}$  be a twice differentiable function on  $(3,5)$  such that  $F(x) = e^{-x}$

$$\int_3^x (3t^2 + 2t + 4F'(t)) dt.$$

If  $F'(4) = \frac{\alpha e^\beta - 224}{(e^\beta - 4)^2}$ , then  $\alpha + \beta$  is equal to\_\_\_\_\_

**Sol. (16)**

$$F(3) = 0$$

$$e^x F(x) = \int_3^x (3t^2 + 2t + 4F'(t)) dt$$

$$e^x F(x) = e^x F'(x) = 3x^2 + 2x + 4F'(x)$$

$$(e^x - 4) \frac{dy}{dx} + e^x y = (3x^2 + 2x)$$

$$\frac{dy}{dx} + \frac{e^x}{(e^x - 4)} y = \frac{(3x^2 + 2x)}{(e^x - 4)}$$

$$y e^{\int \frac{e^x}{(e^x - 4)} dx} = \int \frac{(3x^2 + 2x)}{(e^x - 4)} e^{\int \frac{e^x}{(e^x - 4)} dx} dx$$

$$y(e^x - 4) = \int (3x^2 + 2x) dx + c$$

$$y(e^x - 4) = x^3 + x^2 + c$$

Put  $x = 3 \Rightarrow c = -36$

$$F(x) = \frac{(x^3 + x^2 - 36)}{(e^x - 4)}$$

$$F(x) = \frac{(3x^2 + 2x)(e^x - 4) - (x^3 + x^2 - 36)e^x}{(e^x - 4)^2}$$

$$F'(4) = \frac{56(e^4 - 4) - 44e^4}{(e^4 - 4)^2}$$

$$= \frac{12e^4 - 224}{(e^4 - 4)^2} \Rightarrow \alpha = 12$$

$$\beta = 4$$

$$\alpha + \beta = 16$$

- 3.** Let a plane P pass through the point  $(3, 7, -7)$  and contain the line,  $\frac{x-2}{-3} = \frac{y-3}{2} = \frac{z+2}{1}$ . If distance of the plane P from the origin is d, then  $d^2$  is equal to \_\_\_\_\_.

**Sol. (3)**

$$\vec{BA} = (\hat{i} + 4\hat{j} - 5\hat{k})$$

$$\vec{BA} = (\hat{i} + 4\hat{j} - 5\hat{k})$$

$$\vec{BA} \times \vec{\ell} = \vec{n} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -3 & 2 & 1 \\ 1 & 4 & -5 \end{vmatrix}$$

$$a\hat{i} + b\hat{j} + c\hat{k} = -14\hat{i} - \hat{j} + (-14)\hat{k}$$

$$a = 1, b = 1, c = 1$$

$$\text{Plane is } (x - 2) + (y - 3) + (z + 2) = 0$$

$$x + y + z - 3 = 0$$

$$d = \sqrt{3} \Rightarrow d^2 = 3$$

4. Let  $S = \{1,2,3,4,5,6,7\}$ . Then the number of possible functions  $f: S \rightarrow S$  such that  $f(m.n) = f(m) \cdot f(n)$  for every  $m, n \in S$  and  $m, n \in S$  is equal to\_\_\_\_\_

**Sol. (490)**

$$f(mn) = f(m) \cdot f(n)$$

$$\text{Put } m = 1 \quad \mathbf{f(n) = f(1)} \cdot f(n) \Rightarrow f(1) = 1$$

$$\text{Put } m = n = 2$$

$$f(4) = f(2) \cdot f(2) \begin{cases} f(2) = 1 \Rightarrow f(4) = 1 \\ \text{or} \\ f(2) = 2 \Rightarrow f(4) = 4 \end{cases}$$

$$\text{Put } m = 2, n = 3$$

$$f(6) = f(2) \cdot f(3) \begin{cases} \text{when } f(2) = 1 \\ f(3) = 1 \text{ to } 7 \\ f(2) = 2 \\ f(3) = 1 \text{ or } 2 \text{ or } 3 \end{cases}$$

$f(5), f(7)$  can take any value

$$\text{Total} = (1 \times 1 \times 7 \times 1 \times 7 \times 1 \times 7)$$

$$+ (1 \times 1 \times 3 \times 1 \times 7 \times 1 \times 7)$$


$$= 490$$

5. Let  $f: [0,3] \rightarrow \mathbb{R}$  be defined by


$$f(x) = \min \{x - [x], 1 + [x] - x\}$$

where  $[x]$  is the greatest integer less than or equal to  $x$ . Let  $P$  denote the set containing all  $x \in [0, 3]$  where  $f$  is discontinuous, and  $Q$  denote the set containing all  $x \in (0, 3)$  where  $f$  is not differentiable. Then the sum of number of elements in  $P$  and  $Q$  is equal to \_\_\_\_\_.

**Sol. (5)**



$$1 - \{x\} = 1 - x; 0 \leq x < 1$$



Non differentiable at

$$x = \frac{1}{2}, 1, \frac{3}{2}, 2, \frac{5}{2}$$



6. For real numbers  $\alpha$  and  $\beta$ , consider the following system of linear equations:  
 $x + y - z = 2$ ,  $x + 2y + \alpha z = 1$ ,  $2x - y + z = \beta$ . If the system has infinite solutions, then  $\alpha + \beta$  is equal to \_\_\_\_\_.

**Sol. (5)**

For infinite solutions

$$\Delta = \Delta_1 = \Delta_2 = \Delta_3 = 0$$

$$\Delta = \begin{vmatrix} 1 & 1 & -1 \\ 1 & 2 & \alpha \\ 2 & -1 & 1 \end{vmatrix} = 0$$

$$\Delta = \begin{vmatrix} 3 & 0 & 0 \\ 1 & 2 & \alpha \\ 2 & -1 & 1 \end{vmatrix} = 0$$

$$\Delta = 3(2 + \alpha) = 0$$

$$\Rightarrow \alpha = -2$$

$$\Delta_2 = \begin{vmatrix} 1 & 2 & -1 \\ 1 & 1 & -2 \\ 2 & \beta & 1 \end{vmatrix} = 0$$

$$1(1 + 2\beta) - 2(1 + 4) - (\beta - 2) = 0$$

$$\beta - 7 = 0$$

$$\beta = 7$$

$$\therefore \alpha + \beta = 5 \text{ Ans.}$$

7. Let the domain of the function  $f(x) = \log_4 (\log_5 (\log_3 (18x - x^2 - 77)))$  be  $(a, b)$ . Then the value of the integral  $\int_a^b \frac{\sin^3 x}{(\sin^3 x + \sin^3 (a + b - x))} dx$  is equal to \_\_\_\_\_.

**Sol. (1)**

For domain

$$\log_5 (\log_3 (18x - x^2 - 77)) > 0$$

$$\log_3 (18x - x^2 - 77) > 1$$

$$18x - x^2 - 77 > 3$$

$$x^2 - 18x + 80 < 0$$

$$x \in (8, 10)$$

$$\Rightarrow a = 8 \text{ and } b = 10$$

$$I = \int_a^b \frac{\sin^3 x}{\sin^3 x + \sin^3 (a + b - x)} dx$$

$$I = \int_a^b \frac{\sin^3 (a + b - x)}{\sin^3 x + \sin^3 (a + b - x)} dx$$

$$2I = (b-a) \Rightarrow I = \frac{b-a}{2} (\because a = 8 \text{ and } b = 10)$$

$$I = \frac{10-8}{2} = 1$$

8. If  $\log_3 2, \log_3(2^x-5), \log_3\left(2^x - \frac{7}{2}\right)$  are in an arithmetic progression, then the value of  $x$  is equal to\_\_\_\_\_.

**Sol. (3)**

$$2\log_3(2^x - 5) = \log_3 2 + \log_3\left(2^x - \frac{7}{2}\right)$$

$$\text{Let } 2^x = t$$

$$\log_3(t-5)^2 = \log_3 2\left(t - \frac{7}{2}\right)$$

$$(t-5)^2 = 2t-7$$

$$t^2 - 12t + 32 = 0$$

$$(t-4)(t-8) = 0$$

$$\Rightarrow 2^x = 4 \text{ or } 2^x = 8$$

$$x = 2 \text{ (Rejected)}$$

$$\text{Or } x = 3$$

9. If  $y = y(x), y \in \left[0, \frac{\pi}{2}\right)$  is the solution of the differential equation  $\sec y \frac{dy}{dx} - \sin(x+y) - \sin(x-y) = 0$ , with  $y(0) = 0$ , then  $5y' \left(\frac{\pi}{2}\right)$  is equal to\_\_\_\_\_.

**Sol. (2)**

$$\sec y \frac{dy}{dx} = 2 \sin x \cos y$$

$$\sec^2 y dy = 2 \sin x dx$$

$$\tan y = -2 \cos x + c$$

$$c = 2$$

$$\tan y = -2 \cos x + 2 \Rightarrow \text{at } x = \frac{\pi}{2}$$

$$\tan y = 2$$

$$\sec^2 y \frac{dy}{dx} = 2 \sin x$$

$$5 \frac{dy}{dx} = 2$$

- 10.** Let  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ ,  $\vec{b}$  and  $\vec{c} = \hat{j} - \hat{k}$  be three vectors such that  $\vec{a} \times \vec{b} = \vec{c}$  and  $\vec{a} \cdot \vec{b} = 1$ . If the length of projection vector of the vector  $\vec{b}$  on the vector  $\vec{a} \times \vec{c}$  is  $l$ , then the value of  $3l^2$  is equal to \_\_\_\_\_.

**Sol. (2)**

$$\vec{a} \times \vec{b} = \vec{c}$$

Take Dot with  $\vec{c}$

$$(\vec{a} \times \vec{b}) \cdot \vec{c} = |\vec{c}|^2 = 2$$

Projection of  $\vec{b}$  on  $\vec{a} \times \vec{c} = l$

$$\frac{|\vec{b} \cdot (\vec{a} \times \vec{c})|}{|\vec{a} \times \vec{c}|} = l$$

$$\therefore l = \frac{2}{\sqrt{6}} \Rightarrow l^2 = \frac{4}{6}$$

$$3l^2 = 2$$

अब मोशन ही है, सर्वश्रेष्ठ विकल्प !

MOTION welcomes

Directors of Nucleus Education & Wizard of Mathematics

Now Offline associated with Motion Kota Classroom



Nitin Vijay (NV Sir)  
Managing Director  
Exp. : 18 yrs



Akhilesh Kanther  
(AKK Sir)  
Exp. : 17 yrs



Vishal Joshi  
(VJ Sir)  
Exp. : 18 yrs



Surendra K. Mishra  
(SKM Sir)  
Exp. : 16 yrs



Gavesh Bhardwaj  
(GB Sir)  
Exp. : 17 yrs

## Academic Pillars of JEE Motion Kota



Ram Ratan Dwivedi  
(RRD Sir)  
Joint Director  
Exp. : 20 yrs



Amit Verma  
(AV Sir)  
Joint Director  
Exp. : 16 yrs



Vijay Pratap Singh  
(VPS Sir)  
Vice President  
Exp. : 20 yrs



Nikhil Srivastava  
(NS Sir)  
Head JEE Academics  
Exp. : 17 yrs



Aatish Agarwal  
(AA Sir)  
Sr. Faculty  
Exp. : 17 yrs



Jayant Chittora  
(JC Sir)  
Sr. Faculty  
Exp. : 16 yrs



Anurag Garg  
(AG Sir)  
Sr. Faculty  
Exp. : 17 yrs



Arjun Gupta  
(Arjun Sir)  
Sr. Faculty  
Exp. : 14 yrs



Devki Nandan Pathak  
(DN Sir)  
Sr. Faculty  
Exp. : 13 yrs



Avinash Kishore  
(AVN Sir)  
Sr. Faculty  
Exp. : 9 yrs



Vipin Sharma  
(VS Sir)  
Sr. Faculty  
Exp. : 12 yrs



Durgesh Pandey  
(Pandey Sir)  
Sr. Faculty  
Exp. : 8 yrs

Join **JEE DROPPER BATCH** Kota Classroom

English & Hindi Medium

Online + Offline Mode

Batch Starting from: **4th August 2021**

Time to use all your skills, Now!

# CRASH COURSE

## JEE Advanced 2021

Starting from: **9th August '21**

### Why should you choose?

- ◆ Live Lectures
- ◆ Best JEE Faculties
- ◆ Doubt Support on learning app
- ◆ Online test with discussion
- ◆ Personalized performance analysis

