




**JEE
MAIN
MARCH
2021**

**16th March 2021 | Shift - 2
CHEMISTRY**

JEE | NEET | Foundation

MOTION™

25000+
SELECTIONS SINCE 2007




Identify the reagent(s) 'A' and condition(s) for the reaction

- (1) A = HCl; Anhydrous AlCl_3
- (2) A = HCl, ZnCl_2
- (3) A = Cl_2 , dark, Anhydrous AlCl_3
- (4) A = Cl_2 ; UV light

Ans. (4)

Sol.



2. The INCORRECT statement regarding the structure of C_{60} is:

- (1) It contains 12 six-membered rings and 24 five-membered rings.
- (2) Each carbon atom forms three sigma bonds.
- (3) The five-membered rings are fused only to six-membered rings.
- (4) The six-membered rings are fused to both six and five-membered rings.

Ans. (1)

Sol. it contain 12 five membered ring & 20 six membered ring

3. Match List-I with List-II:

List-I		List-II	
Test/Reagents/Observation(s)		Species detected	
(a)	Lassaigne's Test	(i)	Carbon
(b)	Cu(II) oxide	(ii)	Sulphur
(c)	Silver nitrate	(iii)	N, S, P and halogen
(d)	The sodium fusion extract gives black precipitate with acetic acid & lead acetate	(iv)	Halogen Specifically

The correct match is:


- (1) (a)-(iii), (b)-(i), (c)-(iv), (d)-(ii)
- (2) (a)-(i), (b)-(iv), (c)-(iii), (d)-(ii)
- (3) (a)-(iii), (b)-(i), (c)-(ii), (d)-(iv)
- (4) (a)-(i), (b)-(ii), (c)-(iv), (d)-(iii)

Ans. (1)





Sol. (a)-(iii), (b)-(i), (c)-(iv), (d)-(ii)

Toll Free : 1800-212-1799


www.motion.ac.in | Email : info@motion.ac.in



The structure of X is:



Ans. (1)
Sol.



7. Identify the elements X and Y using the ionisation energy values given below:

Ionization energy (kJ/mol)

	1 st	2 nd
X	495	4563
Y	731	1450

(1) X = F; Y = Mg

(2) X = Mg; Y = F

(3) X = Na; Y = Mg

(4) X = Mg; Y = Na

Ans. (3)

Sol. 2nd I. E of Alkali metals is higher than their respective period.

8. The INCORRECT statements below regarding colloidal solutions is:

(1) A colloidal solution shows colligative properties.

(2) An ordinary filter paper can stop the flow of colloidal particles.

(3) A colloidal solution shows Brownian motion of colloidal particles.

(4) The flocculating power of Al³⁺ is more than that of Na⁺.

Ans. (2)

Sol. Colloidal solutions can pass through ordinary filter paper but cannot pass through special filter colloidal solution coated paper.

9. The characteristics of elements X, Y and Z with atomic numbers, respectively, 33, 53 and 83 are:

(1) X and Z are non-metals and Y is a metalloid.

(2) X and Y are metalloids and Z is a metal

(3) X, Y and Z are metals.

(4) X is a metalloid, Y is a non-metal and Z is a metal.

Ans. (4)

Sol. **Atomic No.** **Element**

(1) 33 —————> As (Metalloid)

(2) 53 —————> I (Non metal)

(3) 83 —————> Bi (Metal)

10. The exact volumes of 1 M NaOH solution required to neutralise 50 mL of 1 M H₃PO₃ solution and 100 mL of 2 M H₃PO₂ solution, respectively, are:

(1) 100 mL and 50 mL

(2) 50 mL and 50 mL

(3) 100 mL and 100 mL

(4) 100 mL and 200 mL

Ans. (4)

Sol. (1) $2\text{NaOH} + \text{H}_3\text{PO}_3 \longrightarrow \text{Na}_2\text{HPO}_3 + 2\text{H}_2\text{O}$

100m mole 50m mole

100m mole = M × V_{ml}

100m mole = 1 × V_{ml}

V_{ml} = 100 ml

(2) $\text{NaOH} + \text{H}_3\text{PO}_2 \longrightarrow \text{NaH}_2\text{PO}_2 + \text{H}_2\text{O}$

200m mole 200m mole

200m mole = M × V_{ml}

V_{ml} = 200 ml

Toll Free : 1800-212-1799

www.motion.ac.in | Email : info@motion.ac.in

15. The correct statements about H_2O_2 are:
(A) used in the treatment of effluents.
(B) used as both oxidising and reducing agents.
(C) the two hydroxyl groups lie in the same plane.
(D) miscible with water.

Choose the correct answer from the options given below:

- (1) (A), (C) and (D) only (2) (A), (B) and (D) only
(3) (A), (B), (C) and (D) (4) (B), (C) and (D) only

Ans. (2)

- Sol. (1) In H_2O_2 oxidation of oxygen is -1 Therefore acts both as O.A and R.A.
(2) H_2O_2 is miscible in water due to inter molecular H-Bonding.
(3) H_2O_2 has open book structure in which both -OH group are not in same plane.

16. The green house gas/es is (are):


- (A) Carbon dioxide (B) Oxygen
(C) Water vapour (D) Methane

Choose the most appropriate answer from the options given below:

- (1) (A) and (B) only (2) (A), (C) and (D) only
(3) (A) and (C) only (4) (A) only

Ans. (2)

- Sol. The green house gases are CO_2 , CH_4 & H_2O vapour.




In the above reaction, the reagent "A" is:

- (1) $NaBH_4$, H_3O^+ (2) HCl , $Zn-Hg$
(3) Alkaline $KMnO_4$, H^+ (4) $LiAlH_4$

Ans. (3)

Sol.



18. Which of the following is least basic?


- (1) $(CH_3CO)_2\ddot{N}H$ (2) $(CH_3CO)\ddot{N}HC_2H_5$
(3) $(C_2H_5)_3\ddot{N}$ (4) $(C_2H_5)_2\ddot{N}H$

Toll Free : 1800-212-1799

www.motion.ac.in | Email : info@motion.ac.in

Ans. (1)

Sol.



Due to higher resonance, ℓp of N is not available for accept H^+

So it is least basic.

19. Fex_2 and Fey_3 are known when x and y are:

- (1) $x=Cl, Br, I$ and $y=F, Cl, Br, I$ (2) $x=F, Cl, Br, I$ and $y=F, Cl, Br$
 (3) $x=F, Cl, Br, I$ and $y=F, Cl, Br, I$ (4) $x=F, Cl, Br$ and $y =F, Cl, Br, I$

Ans. (2)

Sol. FeI_3 , does not react because of I^- being very good reducing agent.

20. The secondary structure of protein is stabilised by:

- (1) van der Waals forces (2) Peptide bond
 (3) Hydrogen bonding (4) glycosidic bond

Ans. (3)

Sol. The secondary structure of protein stabilised by H-bonding.

Section-B

1. At $25^\circ C$, 50 g of iron reacts with HCl to form $FeCl_2$. The evolved hydrogen gas expands against a constant pressure of 1 bar. The work done by the gas during this expansion is _____ J. (Round off to the Nearest Integer).

[Given: $R = 8.14 \text{ J mol}^{-1} \text{ K}^{-1}$. Assume, hydrogen is an ideal gas]

[Atomic mass of Fe is 55.85 u]

Ans. 2218

Sol. $Fe + 2HCl \longrightarrow FeCl_2 + H_2(g)$
50g

$$\text{Moles of Fe} = \frac{50}{55.85} \text{ mol} = \text{moles of } H_2$$

$$\begin{aligned} W_{\text{irrev}} &= -P_{\text{ext}} \cdot \Delta V \\ &= -\text{moles of } H_2 \times RT \\ &= -\frac{50}{55.85} \times 8.314 \times 298 \\ &= -2218.05 \text{ J} \end{aligned}$$

Nearest integer = 2218

2. A 5.0 mol dm^{-3} aqueous solution of KCl has a conductance of 0.55 mS when measured in a cell of cell constant 1.3 cm^{-1} . The molar conductivity of this solution is _____ $\text{mSm}^2 \text{ mol}^{-1}$.

(Round off to the Nearest Integer).

Ans. 14

Sol. $G_{\text{KCl}} = 0.55 \text{ mS} = 55 \times 10^{-5} \text{ S}$
Cell constant = $\ell/A = 1.3 \text{ cm}^{-1}$

Toll Free : 1800-212-1799

www.motion.ac.in | Email : info@motion.ac.in

$$\lambda_M = ??$$

$$R = G(\ell/A) = 55 \times 10^{-5} \times 1.3 \text{ Scm}^{-1}$$

$$\lambda_M = \frac{K \times 1000}{\text{Molarity}} = \frac{55 \times 1.3 \times 10^{-5} \times 1000}{5 \times 10^{-3}}$$

$$\lambda_M = 11 \times 1.3 \times 10 = 11 \times 13 = 143 \text{ S cm}^2 \text{mol}^{-1}$$

$$\lambda_M = \frac{143 \times 1000 \times 10^{-3} \text{ S}}{(10^{-2} \text{ M})^{-2}} \text{ mol}^{-1}$$

$$\lambda_M = 143 \times 1000 \times 10^{-4} (\text{m.S}) \text{m}^2 \cdot \text{mol}^{-1}$$

$$= 14.3$$

Ans. $\lambda_M = 14$ Nearest integer

3. The number of orbitals with $n = 5$, $m_l = +2$ is _____. (Round off to the Nearest Integer).

Ans. 3

Sol. For $n = 5$

$$\ell = 0, 1, 2, 3, 4$$

$$\ell = 2 \rightarrow m = -2, -1, 0, +1, +2$$

$$\ell = 3 \rightarrow m = -3, -2, -1, 0, +1, +2, +3$$

$$\ell = 4 \rightarrow m = -4, -3, -2, -1, 0, +1, +2, +3, +4$$

Total no. of orbitals = 3

4. A and B decompose via first order kinetics with half-lives 54.0 min and 18.0 min respectively. Starting from an equimolar non reactive mixture of A and B, the time taken for the concentration of A to become 16 times that of B is _____ min. (Round off to the Nearest Integer).

Ans. 108

Sol. A $\xrightarrow{1^{\text{st}} \text{ order}}$ $t_{1/2}(\text{A}) = 54$

B $\xrightarrow{1^{\text{st}} \text{ order}}$ $t_{1/2}(\text{B}) = 18$

$$A_0 = B_0 = N_0$$

$$A_t = \frac{A_0}{2^{t/54}}$$

$$B_t = \frac{B_0}{2^{t/18}}$$

$$A_t = 16 \cdot B_t$$

$$\frac{A_0}{2^{t/54}} = 16 \times \frac{B_0}{2^{t/18}}$$

$$2^{t/18 - t/54} = 16$$

$$2^{2t/54} = 16 = 2^4$$

$$\frac{2t}{54} = 4$$

$$t = 108 \text{ min}$$

5. $[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$ absorbs light of wavelength 498 nm during a d-d transition. The octahedral splitting energy for the above complex is _____ $\times 10^{-19}$ J. (Round off to the Nearest Integer).

$$h = 6.626 \times 10^{-34} \text{ Js}; c = 3 \times 10^8 \text{ ms}^{-1}.$$

Ans. (4)

Sol. $\Delta_0 = \frac{hc}{\lambda_{\text{abs}}} = \frac{6.626 \times 10^{-34} \times 3 \times 10^8}{498 \times 10^{-9}}$


$$= \frac{6.626 \times 3}{498} \times 10^{-17} = 0.0399 \times 10^{-17} = 3.99 \times 10^{-19} \simeq 4 \times 10^{-19} \text{ J}$$

Toll Free : 1800-212-1799

www.motion.ac.in | Email : info@motion.ac.in

6. Sulphurous acid (H_2SO_3) has $K_{a1} = 1.7 \times 10^{-2}$ and $K_{a2} = 6.4 \times 10^{-8}$. The pH of 0.588 M H_2SO_3 is _____. (Round off to the Nearest Integer).

Ans. 5



$$\alpha_1 = \sqrt{\frac{1.7 \times 10^{-2}}{0.588}} = \sqrt{\frac{17}{289 \times 2}}$$

Therefore $\frac{\alpha_1 \ll 1}{(1 - \alpha_1) \approx 1}$

Hence $\alpha_2 \ll 1$ & $(1 - \alpha_2) \approx 1$

$\therefore [\text{H}^+] = C\alpha_1$

$$= \sqrt{K_{a1} \times C} = \sqrt{17 \times 10^{-3} \times 0.588}$$

$$= 99.98 \times 10^{-3}$$


$$\text{pH} = 1.99 + 3$$

$$= 4.99 \approx 5$$

7. In Duma's method of estimation of nitrogen, 0.1840 g of an organic compound gave 30 mL of nitrogen collected at 287 K and 758 mm of Hg pressure. The percentage composition of nitrogen in the compound is _____. (Round off to the Nearest Integer).

[Given: Aqueous tension at 287 K = 14 mm of Hg]

Ans. 19



8. Ga (atomic mass 70 u) crystallizes in a hexagonal close packed structure. The total number of voids in 0.581 g of Ga is _____ $\times 10^{21}$. (Round off to the Nearest Integer).

[Given: $N_A = 6.023 \times 10^{23}$]

Ans. 15

Toll Free : 1800-212-1799

www.motion.ac.in | Email : info@motion.ac.in

Sol. No. of moles of Ga = $\frac{0.581}{70}$


No. of atoms of Ga = $\frac{0.581}{70} \times N_A$

∴ Total number of voids = $\frac{0.581}{70} \times N_A \times 3$
 $= 0.0249 \times 6 \times 10^{23}$
 $= 15 \times 10^{21}$

(As there are one octahedral void and two tetrahedral voids per atom)

- 9.** When 35 mL of 0.15 M lead nitrate solution is mixed with 20 mL of 0.12 M chromic sulphate solution, _____ $\times 10^{-5}$ moles of lead sulphate precipitate out. (Round off to the Nearest Integer).

Ans. 525



$3 \text{PbSO}_4 \downarrow + 2\text{Cr}(\text{NO}_3)_3$
 Moles of PbSO_4 = moles of $\text{Pb}(\text{NO}_3)_2$
 $= 5.25 \text{ m mol}$
 $= 525 \times 10^{-5} \text{ mol}$
 Ans. 525

- 10.** At 363 K, the vapour pressure of A is 21 kPa and that of B is 18 kPa. One mole of A and 2 moles of B are mixed. Assuming that this solution is ideal, the vapour pressure of the mixture is _____ kPa. (Round off to the Nearest Integer).

Ans. 19

Sol. $X_A = \frac{1}{1+2} = \frac{1}{3}$ $X_B = \frac{2}{3}$
 $P_A^\circ = 21 \text{ kPa}$ $P_B^\circ = 18 \text{ kPa}$
 $P_{\text{total}} = P_A^\circ X_A + P_B^\circ X_B$
 $= 21 \times \frac{1}{3} + 18 \times \frac{2}{3}$
 $= 7 + 12$
 $= 19 \text{ kPa}$

रिपिटर्स बैच का सर्वश्रेष्ठ परिणाम
सिर्फ मोशन के साथ

MOTION™

Another opportunity to
strengthen your preparation

UNNATI CRASH COURSE

JEE Main May 2021
at Kota Classroom

- ◆ **40 Classes** of each subjects
- ◆ **Doubt Clearing** sessions by **Expert faculties**
- ◆ **Full Syllabus Tests** to improve your question solving skills
- ◆ Thorough learning of concepts with regular classes
- ◆ Get **tips & trick** along with sample papers

Course Fee : ₹ 20,000



Start your **JEE Advanced 2021**
Preparation with

UTTHAN CRASH COURSE

at Kota Classroom

- ◆ Complete course coverage
- ◆ **55 Classes** of each subject
- ◆ **17 Full & 6 Part syllabus** tests will strengthen your exam endurance
- ◆ **Doubt clearing sessions** under the guidance of expert faculties
- ◆ Get **tips & trick** along with sample papers

Course Fee : ₹ 20,000

