



**26<sup>th</sup> Feb. 2021 | Shift - 2**  
**CHEMISTRY**

**JEE | NEET | Foundation**

**MOTION™**

**25000+**  
SELECTIONS SINCE 2007


## Section - A

1. 2,4-DNP test can be used to identify:






- (1) aldehyde
- (2) halogens
- (3) ether
- (4) amine

Ans. (1)

Sol.



2. Identify A in the following chemical reaction.




**Toll Free : 1800-212-1799**

www.motion.ac.in | Email : info@motion.ac.in

Ans. (3)

Sol.



3. The nature of charge on resulting colloidal particles when FeCl<sub>3</sub> is added to excess of hot water is:





- (1) positive
- (2) neutral
- (3) sometimes positive and sometimes negative
- (4) negative

Ans. (1)

Sol. If FeCl<sub>3</sub> is added to excess of hot water, a positively charged sol of hydrated ferric oxide is formed due to adsorption of Fe<sup>3+</sup> ions.

4. Match **List-I** with **List-II**

**List-I**



**List-II**

(i) Wurtz reaction

(ii) Sandmeyer reaction

(iii) Fitting reaction

(iv) Gatterman reaction

Choose the correct answer from the option given below:





(1) (a)-(ii), (b)-(i), (c)-(iv), (d)-(iii)

(2) (a)-(iii), (b)-(iv), (c)-(i), (d)-(ii)

(3) (a)-(ii), (b)-(iv), (c)-(i), (d)-(iii)

(4) (a)-(iii), (b)-(i), (c)-(iv), (d)-(ii)

**Ans. (3)**



**Toll Free : 1800-212-1799**

www.motion.ac.in | Email : info@motion.ac.in

5. In  $\overset{1}{\text{C}}\text{H}_2 = \overset{2}{\text{C}} = \overset{3}{\text{C}}\text{H} - \overset{4}{\text{C}}\text{H}_3$  molecule, the hybridization of carbon 1, 2, 3 and 4 respectively are:

- (1)  $sp^2$ ,  $sp$ ,  $sp^2$ ,  $sp^3$
- (2)  $sp^2$ ,  $sp^2$ ,  $sp^2$ ,  $sp^3$
- (3)  $sp^2$ ,  $sp^3$ ,  $sp^2$ ,  $sp^3$
- (4)  $sp^3$ ,  $sp$ ,  $sp^3$ ,  $sp^3$

**Ans. (1)**

**Sol.**  $\underset{sp^2}{\text{C}}\text{H}_2 = \underset{sp}{\text{C}} = \underset{sp^2}{\text{C}}\text{H} - \underset{sp^3}{\text{C}}\text{H}_3$

6. Match List-I with List-II.

**List-I**

- (a) Sucrose
- (b) Lactose
- (c) Maltose

**List-II**

- (i)  $\beta$ -D-Galactose and  $\beta$ -D-Glucose
- (ii)  $\alpha$ -D-Glucose and  $\beta$ -D-Fructose
- (iii)  $\alpha$ -D-Glucose and  $\alpha$ -D-Glucose

Choose the correct answer from the options given below:

- (1) (a)-(iii), (b)-(ii), (c)-(i)
- (2) (a)-(iii), (b)-(i), (c)-(ii)
- (3) (a)-(i), (b)-(iii), (c)-(ii)
- (4) (a)-(ii), (b)-(i), (c)-(iii)

**Ans. (4)**

**Sol.** Sucrose  $\rightarrow$   $\alpha$ -D- Glucose and  $\beta$ -D- Fructose

Lactose  $\rightarrow$   $\beta$ -D- Galactose and  $\beta$ -D- Glucose

Maltose  $\rightarrow$   $\alpha$ -D- Glucose and  $\alpha$ -D- Glucose

7. Which pair of oxides is acidic in nature?

- (1)  $\text{N}_2\text{O}$ ,  $\text{BaO}$
- (2)  $\text{CaO}$ ,  $\text{SiO}_2$
- (3)  $\text{B}_2\text{O}_3$ ,  $\text{CaO}$
- (4)  $\text{B}_2\text{O}_3$ ,  $\text{SiO}_2$

**Ans. (4)**

**Sol.**  $\text{B}_2\text{O}_3$  and  $\text{SiO}_2$  both are oxides of non-metal and hence are acidic in nature.

**Toll Free : 1800-212-1799**

www.motion.ac.in | Email : info@motion.ac.in

8. Calgon is used for water treatment. Which of the following statement is NOT true about calgon?
- (1) Calgon contains the 2<sup>nd</sup> most abundant element by weight in the earth's crust.
  - (2) It is also known as Graham's salt.
  - (3) It is polymeric compound and is water soluble.
  - (4) It doesnot remove  $\text{Ca}^{2+}$  ion by precipitation.

**Ans. (1)**

**Sol.**  $\text{Na}_6(\text{PO}_3)_6$  or  $\text{Na}_6\text{P}_6\text{O}_{18}$

Order of abundance of element in earth crust is

$\text{O} > \text{Si} > \text{Al} > \text{Fe} > \text{Ca} > \text{Na} > \text{Mg} > \text{K}$

So second most abundant element in earth crust is Si not Ca.

9. Ceric ammonium nitrate and  $\text{CHCl}_3/\text{alc. KOH}$  are used for the identification of functional groups present in \_\_\_\_\_ and \_\_\_\_\_ respectively.
- (1) alcohol, amine
  - (2) amine, alcohol
  - (3) alcohol, phenol
  - (4) amine, phenol

**Ans. (1)**

**Sol.** Alcohol give positive test with ceric ammonium nitrate and primary amines gives carbyl amine test with  $\text{CHCl}_3$ ,  $\text{KOH}$ .

10. Given below are two statements: one is labelled as Assertion A and the other is labelled as Reason R.

Assertion A: In  $\text{TlI}_3$ , isomorphous to  $\text{CsI}_3$ , the metal is present in +1 oxidation state.

Reason R: Tl metals has fourteen *f* electrons in its electronic configuration.

In the light of the above statements, choose the most appropriate answer from the options given below:

- (1) Both A and R are correct and R is the correct explanation of A
- (2) A is not correct but R is correct
- (3) Both A and R are correct R is NOT the correct explanation of A
- (4) A is correct but R is not correct

**Ans. (3)**

**Sol.**  $\text{TlI}_3$  is  $\text{Tl}^+ \text{I}_3^-$

$\text{CsI}_3$  is  $\text{Cs}^+ \text{I}_3^-$

Thallium shows  $\text{Tl}^+$  state due to inert pair effect.

**Toll Free : 1800-212-1799**

**www.motion.ac.in | Email : info@motion.ac.in**


11. The correct order of electron gain enthalpy is:

- (1)  $S > Se > Te > O$
- (2)  $O > S > Se > Te$
- (3)  $S > O > Se > Te$
- (4)  $Te > Se > S > O$

Ans. (1)


Sol. Electron gain enthalpy of O is very low due to small size.

12. Identify A in the given chemical reaction.



- (1)
- (2)
- (3)
- (4)

Ans. (1)



(Internal aldol condensation)

13. Match List-I with List-II

List-I	List-II
(a) Siderite	(i) Cu
(b) Calamine	(ii) Ca
(c) Malachite	(iii) Fe
(d) Cryolite	(iv) Al
	(v) Zn

Choose the correct answer from the options given below:

- (1) (a)-(i), (b)-(ii), (c)-(v), (d)-(iii)
- (2) (a)-(iii), (b)-(v), (c)-(i), (d)-(iv)
- (3) (a)-(i), (b)-(ii), (c)-(iii), (d)-(iv)
- (4) (a)-(iii), (b)-(i), (c)-(v), (d)-(ii)






**Toll Free : 1800-212-1799**

www.motion.ac.in | Email : info@motion.ac.in


**Ans. (2)**

**Sol.** Siderite -  $\text{FeCO}_3$   
 Calamine -  $\text{ZnCO}_3$   
 Malachite -  $\text{CuCO}_3 \cdot \text{Cu(OH)}_2$   
 Cryolite -  $\text{Na}_3\text{AlF}_6$

**14.** Identify A in the given reaction



**Ans. (2)**



**Toll Free : 1800-212-1799**

www.motion.ac.in | Email : info@motion.ac.in

15. Match List-I with List-II.

List-I	List-II
(a) Sodium Carbonate	(i) Deacon
(b) Titanium	(ii) Caster-Kellner
(c) Chlorine	(iii) Van-Arkel
(d) Sodium hydroxide	(iv) Solvay


Choose the correct answer from the option given below:

- (1) (a)-(iii), (b)-(ii), (c)-(i), (d)-(iv)  
 (2) (a)-(iv), (b)-(iii), (c)-(i), (d)-(ii)  
 (3) (a)-(iv), (b)-(i), (c)-(ii), (d)-(iii)  
 (4) (a)-(i), (b)-(iii), (c)-(iv), (d)-(ii)


**Ans. (2)**

**Sol.** Sodium carbonate  $\text{Na}_2\text{CO}_3$  &  $\text{NaHCO}_3$

Titanium : Van arkel method



**Chlorine** : Deacon's process



Sodium hydroxide :- Caster-Kellner cell

16. Match List-I with List-II.

List-I (Molecule)	List-II (Bond order)
(a) $\text{Ne}_2$	(i) 1
(b) $\text{N}_2$	(ii) 2
(c) $\text{F}_2$	(iii) 0
(d) $\text{O}_2$	(iv) 3

Choose the correct answer from the options given below:

- (1) (a)-(iii), (b)-(iv), (c)-(i), (d)-(ii)      (2) (a)-(i), (b)-(ii), (c)-(iii), (d)-(iv)  
 (3) (a)-(ii), (b)-(i), (c)-(iv), (d)-(iii)      (4) (a)-(iv), (b)-(iii), (c)-(ii), (d)-(i)

**Ans. (1)**

**Sol.**

$\text{Ne}_2\text{O}$	BO = 0
$\text{N}_2$	BO = 3
$\text{F}_2$	BO = 1
$\text{O}_2$	BO = 2

As per molecular orbital theory

17. Which of the following forms of hydrogen emits low energy  $\beta^-$  particles?

- (1) Proton  $H^+$
- (2) Deuterium  ${}^2_1H$
- (3) Protium  ${}^1_1H$
- (4) Tritium  ${}^3_1H$

**Ans. (4)**


**Sol.** Tritium isotope of hydrogen is radioactive and emits low energy  $\beta^-$  particles. It is because of high n/p ratio of tritium which makes nucleus unstable.

18. A. Phenyl methanamine  
 B. N, N-Dimethylaniline  
 C. N-Methyl aniline  
 D. Benzenamine


Choose the correct order of basic nature of the above amines.

- (1)  $D > C > B > A$
- (2)  $D > B > C > A$
- (3)  $A > C > B > D$
- (4)  $A > B > C > D$

**Ans. (4)**



19.



Considering the above reaction, the major product among the following is:

- (1)
- (2)
- (3)
- (4)

**Toll Free : 1800-212-1799**


www.motion.ac.in | Email : info@motion.ac.in



2. The number of stereoisomers possible for  $[\text{Co}(\text{ox})_2(\text{Br})(\text{NH}_3)]^{2-}$  is \_\_\_\_\_ [ox = oxalate]

Ans. 3

Sol.  $[\text{Co}(\text{ox})_2\text{Br}(\text{NH}_3)]^{2-}$



Total stereoisomer = 2 (OI) + 1 POE (pair of enantiomers) = 3

3. The average S-F bond energy in  $\text{kJ mol}^{-1}$  of  $\text{SF}_6$  is \_\_\_\_\_. (Rounded off to the nearest integer)

**[Given :** The values of standard enthalpy of formation of  $\text{SF}_6(\text{g})$ ,  $\text{S}(\text{g})$  and  $\text{F}(\text{g})$  are - 1100, 275 and 80  $\text{kJ mol}^{-1}$  respectively.]

Ans. 309

Sol.  $\text{SF}_6(\text{g}) \longrightarrow \text{S}(\text{g}) + 6\text{F}(\text{g})$

$$\Delta H_{\text{reaction}}^{\circ} = 6 \times E_{\text{S-F}} = \Delta H_f^{\circ}[\text{S}(\text{g})] + 6 \times \Delta H_f^{\circ}[\text{F}(\text{g})] - \Delta H_f^{\circ}[\text{SF}_6(\text{g})]$$

$$6 \times E_{\text{S-F}} = 275 + 6 \times 80 - (-1100)$$

$$= 275 + 480 + 1100$$

$$6 \times E_{\text{S-F}} = 1855$$

$$E_{\text{S-F}} = \frac{1855}{6} = 309.1667$$

$\simeq 309 \text{ kJ/mol}$  Ans.

**Toll Free : 1800-212-1799**


www.motion.ac.in | Email : info@motion.ac.in

4. Emf of the following cell at 298 K in V is  $x \times 10^{-2}$ .  
 $\text{Zn}|\text{Zn}^{2+} (0.1 \text{ M})||\text{Ag}^+(0.01 \text{ M})|\text{Ag}$   
 The value of x is \_\_\_\_\_. (Rounded off to the nearest integer)

[Given:  $E_{\text{Zn}^{2+}/\text{Zn}}^0 = -0.76\text{V}$ ;  $E_{\text{Ag}^+/\text{Ag}}^0 = +0.80\text{V}$ ;  $\frac{2.303RT}{F} = 0.059$ ]

Ans. 147

Sol.  $\text{Zn(s)}|\text{Zn}^{2+}(0.1\text{M})||\text{Ag}^+(0.01\text{M})|\text{Ag(s)}$



$$E^0 = 0.80 + 0.76 = 1.56 ; Q = \left\{ \frac{\text{Zn}^{2+}}{(\text{Ag}^+)^2} \right\}$$

$$E = E^0 - \frac{0.059}{n} \log(Q)$$

$$E = 1.56 - \frac{0.059}{2} \log \left[ \frac{0.1}{(0.01)^2} \right]$$

$$E = 1.56 - \frac{0.059}{2} \log \left[ (10)^3 \right]$$

$$E = 1.4715 = 147.15 \times 10^{-2} \text{ volt}$$

$$= x \times 10^{-2}$$

$$X = 147.15 \simeq 147 \text{ Ans.}$$

5. A ball weighing 10g is moving with a velocity of  $90\text{ms}^{-1}$ . If the uncertainty in its velocity is 5%, then the uncertainty in its position is \_\_\_\_\_  $\times 10^{-33}\text{m}$ . (Rounded off to the nearest integer)  
 [Given :  $h = 6.63 \times 10^{-34} \text{ Js}$ ]

Ans. 1

Sol.  $m = 10 \text{ g} = 10^{-2} \text{ Kg}$

$v = 90 \text{ m/sec.}$

$$\Delta v = v \times 5\% = 90 \times \frac{5}{100} = 4.5 \text{ m / sec}$$

$$m \cdot \Delta v \cdot \Delta x \geq \frac{h}{4\pi}$$

$$10^{-2} \times 4.5 \times \Delta x \geq \frac{6.63 \times 3 \times 10^{-34}}{4 \times \frac{22}{7}}$$

$$\Delta x \geq \frac{6.63 \times 7 \times 2 \times 10^{-34}}{9 \times 4 \times 22 \times 10^{-2}}$$

$$\Delta x \geq 1.17 \times 10^{-33} = x \times 10^{-33}$$


$$x = 1.17 \simeq 1$$

Toll Free : 1800-212-1799

www.motion.ac.in | Email : info@motion.ac.in

6. In mildly alkaline medium, thiosulphate ion is oxidized by  $\text{MnO}_4^-$  to "A". The oxidation state of sulphur in "A" is \_\_\_\_\_.

**Ans. 6**



$\therefore$  Oxidation no. of 'S' = +6 Ans.

7. When 12.2 g of benzoic acid is dissolved in 100g of water, the freezing point of solution was found to be  $-0.93^\circ\text{C}$  ( $K_f(\text{H}_2\text{O}) = 1.86 \text{ K kg mol}^{-1}$ ). The number (n) of benzoic acid molecules associated (assuming 100% association) is \_\_\_\_\_.

**Ans. 2**



$$N = \frac{1}{x} = i \{ \alpha = 1 \}$$

$$\Delta T_f = i \times k_f \times m$$

$$0.93 = \frac{1}{n} \times 1.86 \times \frac{12.2 \times 1000}{122 \times 100}$$


$$n = 2$$

8. If the activation energy of a reaction is  $80.9 \text{ kJ mol}^{-1}$ , the fraction of molecules at 700K, having enough energy to react to form products is  $e^{-x}$ . The value of x is \_\_\_\_\_.

(Rounded off to the nearest integer)

[Use  $R = 8.31 \text{ JK}^{-1} \text{ mol}^{-1}$ ]

**Ans. 14**



Fraction of molecules able to cross energy barrier =  $e^{-E_a/RT} = e^{-x}$

$$x = \frac{E_a}{RT} = \frac{80.9 \times 1000}{8.31 \times 700} = 13.91$$


$$x \simeq 14 \text{ Ans}$$

**Toll Free : 1800-212-1799**

**www.motion.ac.in | Email : info@motion.ac.in**

9. The pH of ammonium phosphate solution, if  $pK_a$  of phosphoric acid and  $pK_b$  of ammonium hydroxide are 5.23 and 4.75 respectively, is \_\_\_\_\_.

**Ans. 7**



$$[H^+] = K_a \times \sqrt{\frac{K_w}{K_a \times K_b}}$$

$$pH = pK_a + \frac{1}{2} \{pK_w - pK_a - pK_b\}$$

$$pH = 5.23 + \frac{1}{2} \{14 - 5.23 - 4.75\}$$

$$pH = 5.23 + \frac{1}{2} (4.02) = 7.24 = 7 \text{ (Nearest integer)}$$

10. The number of octahedral voids per lattice site in a lattice is \_\_\_\_\_.  
(Rounded off to the nearest integer)

**Ans. 1**

**Sol.** Assuming FCC

No of lattice sites = 6 face centre + 8 corner = 14

No. of octahedral voids = 13

$$\text{Ratio} = \frac{13}{14} = 0.92857 = 1 \text{ (Nearest integer)}$$

रिपिटर्स बैच का सर्वश्रेष्ठ परिणाम  
सिर्फ मोशन के साथ

**MOTION™**

Another opportunity to  
strengthen your preparation

## **UNNATI CRASH COURSE**

JEE Main May 2021  
at Kota Classroom

- ◆ 40 Classes of each subjects
- ◆ Doubt Clearing sessions by Expert faculties
- ◆ Full Syllabus Tests to improve your question solving skills
- ◆ Thorough learning of concepts with regular classes
- ◆ Get tips & trick along with sample papers

Course Fee : ₹ 20,000



Start your **JEE Advanced 2021**  
Preparation with

## **UTTHAN CRASH COURSE**

at Kota Classroom

- ◆ Complete course coverage
- ◆ 55 Classes of each subject
- ◆ 17 Full & 6 Part syllabus tests will strengthen your exam endurance
- ◆ Doubt clearing sessions under the guidance of expert faculties
- ◆ Get tips & trick along with sample papers

Course Fee : ₹ 20,000



**Toll Free : 1800-212-1799**

www.motion.ac.in | Email : info@motion.ac.in