


हमारा विश्वास... हर एक विद्यार्थी है खास

**JEE  
MAIN  
JAN  
2020**

**PAPER WITH SOLUTION**

**8<sup>th</sup> January 2020 \_ SHIFT - II**

**MATHEMATICS**



**24000+**  
SELECTIONS SINCE 2007

JEE (Advanced)

**5392**

(Under 50000 Rank)

JEE (Main)

**16241**

NEET / AIIMS

**1305**

(since 2016)

NTSE / OLYMPIADS

**1158**

(5<sup>th</sup> to 10<sup>th</sup> class)

**MOTION™**

Nurturing potential through education

H.O. : 394, Rajeev Gandhi Nagar, Kota

[www.motion.ac.in](http://www.motion.ac.in) | [info@motion.ac.in](mailto:info@motion.ac.in)

1. If  $I = \int_1^2 \frac{dx}{\sqrt{2x^3 - 9x^2 + 12x + 4}}$ , then :

यदि  $I = \int_1^2 \frac{dx}{\sqrt{2x^3 - 9x^2 + 12x + 4}}$  है, तो :

(1)  $\frac{1}{9} < I^2 < \frac{1}{8}$       (2)  $\frac{1}{6} < I^2 < \frac{1}{2}$       (3)  $\frac{1}{16} < I^2 < \frac{1}{9}$       (4)  $\frac{1}{8} < I^2 < \frac{1}{4}$

**Sol. 1**

$$I = \int_1^2 \frac{dx}{\sqrt{2x^3 - 9x^2 + 12x + 4}}$$

$$f(x) = \frac{1}{\sqrt{2x^3 - 9x^2 + 12x + 4}}$$

$$f'(x) = \frac{-1}{2} \frac{(6x^2 - 18x + 12)}{(2x^3 - 9x^2 + 12x + 4)^{3/2}}$$


$$f'(x) = \frac{-6(x-1)(x-2)}{2(2x^3 - 9x^2 + 12x + 4)^{3/2}}$$

$$f(1) = \frac{1}{3} \text{ \& } f(2) = \frac{1}{\sqrt{8}}$$

it is increasing function

$$\therefore \frac{1}{3} < I < \frac{1}{\sqrt{8}}$$

$$\frac{1}{9} < I^2 < \frac{1}{8}$$



2. If a line,  $y = mx + c$  is a tangent to the circle,  $(x - 3)^2 + y^2 = 1$  and it is perpendicular to a line  $L_1$ , where  $L_1$  is the tangent to the circle,  $x^2 + y^2 = 1$  at the point  $\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$ ; then :

यदि एक रेखा  $y = mx + c$ , वृत्त  $(x - 3)^2 + y^2 = 1$  की एक स्पर्श रेखा है तथा यह एक रेखा  $L_1$  पर लम्ब है, जहाँ  $L_1$  वृत्त

$x^2 + y^2 = 1$  के बिन्दु  $\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$  पर स्पर्श रेखा है, तो :

(1)  $c^2 - 7c + 6 = 0$     (2)  $c^2 + 7c + 6 = 0$     (3)  $c^2 - 6c + 7 = 0$     (4)  $c^2 + 6c + 7 = 0$

**Sol. 4**

$(x - 3)^2 + y^2 = 1$ , tangent is  $y = mx + c$

for circle  $x^2 + y^2 = 1$  tangent at  $P\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$



**Sol. 1**

$$\alpha = -\frac{1}{2} + i\frac{\sqrt{3}}{2} \Rightarrow \boxed{\alpha = \omega}$$

$$a = (1 + \alpha) \sum_{k=0}^{100} \alpha^{2k}$$

$$a = (1 + \alpha)(1 + \alpha^2 + \alpha^4 + \alpha^6 + \dots + \alpha^{200})$$

$$a = (1 + \alpha) \left\{ \frac{1 \cdot \left[ (\alpha^2)^{101} - 1 \right]}{\alpha^2 - 1} \right\} \Rightarrow a = \frac{(\omega + 1)(\omega^{202} - 1)}{(\omega^2 - 1)}$$

$$\Rightarrow a = \frac{(\omega + 1)(\omega - 1)}{(\omega + 1)(\omega - 1)} \Rightarrow \boxed{a = 1}$$

$$b = \sum_{k=0}^{100} \alpha^{3k} = 1 + \alpha^3 + \alpha^6 + \dots + \alpha^{300}$$

$$b = 1 + \omega^3 + \omega^6 + \dots + \omega^{300} \Rightarrow b = 1 + 1 + \dots 101 \text{ times}$$

$$b = 101$$

$$x^2 - (a + b)x + (ab) = 0$$

$$x^2 - (102)x + 101 = 0$$

**5.** let  $\vec{a} = \hat{i} - 2\hat{j} + \hat{k}$  and  $\vec{b} = \hat{i} - \hat{j} + \hat{k}$  be two vectors. If  $\vec{c}$  is a vector such that  $\vec{b} \times \vec{c} = \vec{b} \times \vec{a}$  and  $\vec{c} \cdot \vec{a} = 0$ , then  $\vec{c} \cdot \vec{b}$  is equal to :

माना दो सदिश  $\vec{a} = \hat{i} - 2\hat{j} + \hat{k}$  तथा  $\vec{b} = \hat{i} - \hat{j} + \hat{k}$  है। यदि एक सदिश  $\vec{c}$  इस प्रकार है कि  $\vec{b} \times \vec{c} = \vec{b} \times \vec{a}$  तथा  $\vec{c} \cdot \vec{a} = 0$  हैं, तो  $\vec{c} \cdot \vec{b}$  बराबर है :

(1)  $\frac{1}{2}$

(2)  $-\frac{1}{2}$

(3)  $-\frac{3}{2}$

(4) -1

**Sol. 2**

$$\vec{a} = \hat{i} - 2\hat{j} + \hat{k}$$

$$\vec{b} = \hat{i} - \hat{j} + \hat{k}$$

$$|\vec{a}| = \sqrt{6} ; |\vec{b}| = \sqrt{3} \quad \& \quad \vec{a} \cdot \vec{b} = 4$$

$$\vec{b} \times \vec{c} = \vec{b} \times \vec{a}$$

$$(\vec{b} \times \vec{c}) - (\vec{b} \times \vec{a}) = 0$$

$$\vec{b} \times (\vec{c} - \vec{a}) = 0$$

$$\vec{b} \parallel (\vec{c} - \vec{a}) \Rightarrow (\vec{c} - \vec{a}) = \lambda \vec{b}$$

$$\vec{c} = \vec{a} + \lambda \vec{b}$$

Now  $\vec{c} \cdot \vec{a} = 0$

$$\Rightarrow \vec{c} \cdot \vec{a} = \vec{a} \cdot \vec{a} + \lambda (\vec{a} \cdot \vec{b})$$

$$0 = |\vec{a}|^2 + \lambda (\vec{a} \cdot \vec{b})$$

$$\lambda = \frac{-|\vec{a}|^2}{\vec{a} \cdot \vec{b}} = \frac{-6}{4} = \frac{-3}{2}$$

$$\therefore \vec{c} = \vec{a} - \frac{3}{2} \vec{b}$$

$$\vec{c} = (\hat{i} - 2\hat{j} + \hat{k}) - \frac{3}{2}(\hat{i} - \hat{j} + \hat{k})$$

$$\vec{c} = -\frac{1}{2}\hat{i} - \frac{1}{2}\hat{j} - \frac{1}{2}\hat{k} \Rightarrow C = -\frac{1}{2}(\hat{i} + \hat{j} + \hat{k})$$

$$\vec{b} \cdot \vec{c} = -\frac{1}{2}(\hat{i} + \hat{j} + \hat{k})(\hat{i} - \hat{j} + \hat{k})$$

$$\vec{b} \cdot \vec{c} = -\frac{1}{2}$$

6. If  $\alpha$  and  $\beta$  be the coefficients of  $x^4$  and  $x^2$  respectively in the expansion of  $(x + \sqrt{x^2 - 1})^6 + (x - \sqrt{x^2 - 1})^6$ , then :

यदि  $(x + \sqrt{x^2 - 1})^6 + (x - \sqrt{x^2 - 1})^6$  के प्रसार में  $x^4$  तथा  $x^2$  के गुणांक क्रमशः  $\alpha$  तथा  $\beta$  हैं, तो :

(1)  $\alpha - \beta = 60$       (2)  $\alpha - \beta = -132$       (3)  $\alpha + \beta = 60$       (4)  $\alpha + \beta = -30$

**Sol.**

$$(x+a)^n + (x-a)^n = 2(T_1 + T_3 + T_5 + \dots)$$

$$(x + \sqrt{x^2 - 1})^6 + (x - \sqrt{x^2 - 1})^6 = 2[T_1 + T_3 + T_5 + T_7]$$

$$= 2[{}^6C_0 x^6 + {}^6C_2 x^4(x^2 - 1) + {}^6C_4 x^2(x^2 - 1)^2 + {}^6C_6 x^0(x^2 - 1)^3]$$

$$= 2[x^6 + 15(x^6 - x^4) + 15x^2(x^4 + 1 - 2x^2) + [x^6 - 3x^4 + 3x^2 - 1]]$$

$$= 2[x^6(2 + 15 + 15 + 1) + x^4(-15 - 30 - 3) + x^2(15 + 3)]$$

coefficient of  $x^4 = \boxed{\alpha = -96}$

$\boxed{\beta = 36}$

$\alpha - \beta = -96 - 36 = -132$

**Increase Your Score  
for JEE Main April'2020**

**उत्कर्ष**  
15 JAN 2020

percentile between 97.0 to 98.99  
in JEE Main (Jan-2020)

Fees - ₹ 22000 Including GST

**उन्नति**  
17 JAN 2020

Below 97 percentile in JEE Main (Jan-2020)  
Tenure: 62 Days | Schedule: 5 Classes Per Day

Fees - ₹ 27500 Including GST

**उत्थान**  
17 JAN 2020

99 percentile and above  
in JEE Main (Jan-2020)

Fees - ₹ 11000 (score 160-200) | Fees - ₹ 5500 (score 200-240) | Fees - ₹ 0 (score above 240)

7. If a hyperbola passes through the point P(10,16) and it has vertices at  $(\pm 6, 0)$ , then the equation of the normal to it at P is :

यदि एक अतिपरवलय बिन्दु P(10, 16) से होकर जाता है तथा इसके शीर्ष  $(\pm 6, 0)$  पर हैं, तो P पर इसके अभिलम्ब का समीकरण है :

(1)  $x + 2y = 42$       (2)  $x + 3y = 58$       (3)  $2x + 5y = 100$       (4)  $3x + 4y = 94$

Sol. **3**

Let hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

vertices  $(\pm a, 0) = (\pm 6, 0) \Rightarrow \boxed{a = 6}$

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \xrightarrow{(10,16)} \frac{(10)^2}{(6)^2} - \frac{(16)^2}{b^2} = 1 \Rightarrow \frac{256}{b^2} = \frac{100}{36} - 1$$

$$\frac{256}{b^2} = \frac{64}{36}$$

$$b^2 = \frac{256 \times 36}{64} \Rightarrow b^2 = 36 \times 4$$

$$b^2 = 9 \times 16$$

$$\boxed{b = 12}$$

$\therefore$  required hyperbola is  $\frac{x^2}{36} - \frac{y^2}{144} = 1$

equation of normal will be

$$\frac{a^2x}{x_1} + \frac{b^2y}{y_1} = a^2 + b^2$$

at P(10,16) normal is

$$\frac{36x}{10} + \frac{144y}{16} = 36 + 144$$

$$\frac{18x}{5} + 9y = 180$$

$$18x + 45y = 900$$

$$\boxed{2x + 5y = 100}$$

8.  $\lim_{x \rightarrow 0} \frac{\int_0^x t \sin(10t) dt}{x}$  is equal to :

$\lim_{x \rightarrow 0} \frac{\int_0^x t \sin(10t) dt}{x}$  बराबर है :

(1)  $\frac{1}{10}$

(2)  $-\frac{1}{5}$

(3)  $\frac{-1}{10}$

(4) 0

Sol. 4

$\lim_{x \rightarrow 0} \frac{\int_0^x t \sin(10t) dt}{x}$

$\frac{0}{0}$  form

∴ apply newton leibnitz's rule

$\lim_{x \rightarrow 0} \frac{x \cdot \sin(10x) - 0}{1} = 0$

9. The system of linear equations

$\lambda x + 2y + 2z = 5$

$2\lambda x + 3y + 5z = 8$

$4x + \lambda y + 6z = 10$  has :

(1) no solution when  $\lambda = 2$

(3) no solution when  $\lambda = 8$

रैखिक समीकरण निकाय

$\lambda x + 2y + 2z = 5$

$2\lambda x + 3y + 5z = 8$

$4x + \lambda y + 6z = 10$

(1) का कोई हल नहीं है जब  $\lambda = 2$

(3) का कोई हल नहीं है जब  $\lambda = 8$

(2) infinitely many solutions when  $\lambda = 2$

(4) a unique solution when  $\lambda = -8$

Sol. 1

$\lambda x + 2y + 2z = 5$

$2\lambda x + 3y + 5z = 8$

$4x + \lambda y + 6z = 10$

$\Delta = \begin{vmatrix} \lambda & 2 & 2 \\ 2\lambda & 3 & 5 \\ 4 & \lambda & 6 \end{vmatrix}$

$\Delta = \lambda (18 - 5\lambda) - 2(12\lambda - 20) + 2(2\lambda^2 - 12)$

$\Delta = 18\lambda - 5\lambda^2 - 24\lambda + 40 + 4\lambda^2 - 24$

$\Delta = -\lambda^2 - 6\lambda + 16$

**Increase Your Score  
for JEE Main April'2020**

**उन्नति**  
17 JAN 2020

Below 97 percentile in JEE Main (Jan-2020)

Tonure: 62 Days | Schedule: 5 Classes Per Day

Fees - ₹ 27500 Including GST

**उत्कर्ष**  
15 JAN 2020

percentile between 97.0 to 98.99  
in JEE Main (Jan-2020)

Fees - ₹ 22000 Including GST

**उत्थान**  
17 JAN 2020

99 percentile and above  
in JEE Main (Jan-2020)

Fees - ₹ 11000  
score 160-200

Fees - ₹ 5500  
score 200-240

Fees - ₹ 0  
score above 240



$$\Delta = -\lambda^2 + 2\lambda - 8\lambda + 16$$

$$\Delta = -\lambda(\lambda - 2) - 8(\lambda - 2)$$

$$\Delta = -(\lambda + 8)(\lambda - 2)$$

for no solutions  $\Delta = 0 \Rightarrow \lambda = -8, \lambda = 2$   
when  $\lambda = 2$

$$\Delta_1 = \begin{vmatrix} 5 & 2 & 2 \\ 8 & 3 & 5 \\ 10 & 2 & 6 \end{vmatrix}$$

$$\Delta_1 = 40 + 4 - 28$$

$$\Delta_1 \neq 0$$

$\therefore$  at  $\lambda = 2$  there is No solution

- 10.** If the 10<sup>th</sup> term of an A.P. is  $\frac{1}{20}$  and its 20<sup>th</sup> term is  $\frac{1}{10}$ , then the sum of its first 200 terms is :

यदि एक समान्तर श्रेणी का 10 वां पद  $\frac{1}{20}$  है तथा इसका 20 वां पद  $\frac{1}{10}$  है, तो इसके प्रथम 200 पदों का योग है :

- (1) 100                      (2)  $50\frac{1}{4}$                       (3) 50                      (4)  $100\frac{1}{2}$

**Sol. 4**

$$T_{10} = a + 9d = \frac{1}{20} \quad \dots(1)$$

$$T_{20} = a + 19d = \frac{1}{10} \quad \dots(2)$$

Equation (2) - (1)

$$-10d = \frac{1}{20} - \frac{1}{10}$$

$$-10d = \frac{1}{20} - \frac{2}{20} = \frac{-1}{20}$$

$$\boxed{d = \frac{1}{200}}$$

$$a + \frac{9}{200} = \frac{1}{20} \Rightarrow a = \frac{1}{20} - \frac{9}{200}$$

$$a = \frac{10}{200} - \frac{9}{200} \Rightarrow \boxed{a = \frac{1}{200}}$$

$$\therefore \boxed{a=d = \frac{1}{200}}$$

$$S_{200} = \frac{200}{2} \left[ \frac{2}{200} + (200-1) \cdot \frac{1}{200} \right] = 100 \left[ \frac{2}{200} + \frac{199}{200} \right]$$

$$S_{200} = \frac{201}{2} = 100\frac{1}{2}$$



11. Let  $f : (1,3) \rightarrow \mathbb{R}$  be a function defined by  $f(x) = \frac{x[x]}{1+x^2}$ , where  $[x]$  denotes the greatest integer  $\leq x$ . Then the range of  $f$  is :

माना  $f : (1,3) \rightarrow \mathbb{R}$  एक फलन है, जो  $f(x) = \frac{x[x]}{1+x^2}$ , द्वारा परिभाषित है जहाँ  $[x]$  महत्तम पूर्णांक  $\leq x$  को दर्शाता है। तो  $f$  का परिसर है :

(1)  $\left(\frac{2}{5}, \frac{1}{2}\right) \cup \left(\frac{3}{5}, \frac{4}{5}\right]$  (2)  $\left(\frac{2}{5}, \frac{4}{5}\right]$  (3)  $\left(\frac{3}{5}, \frac{4}{5}\right]$  (4)  $\left(\frac{2}{5}, \frac{3}{5}\right] \cup \left(\frac{3}{4}, \frac{4}{5}\right]$

Sol. 1

$$f: (1,3) \rightarrow \mathbb{R}, f(x) = \frac{x[x]}{1+x^2}$$

$$f(x) = \begin{cases} \frac{x}{1+x^2}, & x \in (1,2) \\ \frac{2x}{1+x^2}, & x \in [2,3) \end{cases}$$

$$f'(x) = \begin{cases} \frac{(1+x^2)(1) - x(2x)}{(1+x^2)^2}, & x \in (1,2) \\ \frac{(1+x^2)(2) - 2x(2x)}{(1+x^2)^2}, & x \in [2,3) \end{cases}$$

$$f'(x) = \begin{cases} \frac{1-x^2}{1+x^2}, & x \in (1,2) \\ \frac{1-2x^2}{1+x^2}, & x \in [2,3) \end{cases}$$

$\therefore f(x)$  is decreasing function

$$\therefore R_f \in \left(\frac{2}{5}, \frac{1}{2}\right) \cup \left(\frac{6}{10}, \frac{4}{5}\right]$$

12. The differential equation of the family of curves,  $x^2 = 4b(y + b)$ ,  $b \in \mathbb{R}$ , is :

वक्रों  $x^2 = 4b(y + b)$ ,  $b \in \mathbb{R}$  के कुल का अवकल समीकरण है :

(1)  $x(y')^2 = 2yy' - x$  (2)  $xy'' = y'$  (3)  $x(y')^2 = x - 2yy'$  (4)  $x(y')^2 = x + 2yy'$

Sol. 4

$$x^2 = 4b(y + b), b \in \mathbb{R}$$

$$x^2 = 4by + 4b^2$$

$$2x = 4by' + 0$$

**Increase Your Score  
for JEE Main April'2020**

**उत्कर्ष**  
15 JAN 2020

percentile between 97.0 to 98.99  
in JEE Main (Jan-2020)

Fees - ₹ 22000 Including GST

**उन्नति**  
17 JAN 2020

Below 97 percentile in JEE Main (Jan-2020)  
Tenure: 62 Days | Schedule: 5 Classes Per Day

Fees - ₹ 27500 Including GST

**उत्थान**  
17 JAN 2020

99 percentile and above  
in JEE Main (Jan-2020)

Fees - ₹ 11000 score 160-200 | Fees - ₹ 5500 score 200-240 | Fees - ₹ 0 score above 240

$$y' = \frac{x}{2b}$$

$$\Rightarrow \boxed{b = \frac{x}{2y'}}$$

∴ differential equation is

$$x^2 = 4 \cdot y \cdot \frac{x}{2y'} + 4 \left( \frac{x}{2y'} \right)^2$$

$$x^2 = \frac{2xy}{y'} + \frac{x^2}{(y')^2}$$

$$x = \frac{2y}{y'} + \frac{x}{(y')^2}$$

$$\boxed{x(y')^2 = 2yy' + x}$$

- 13.** Let S be the set of all functions  $f: [0,1] \rightarrow \mathbb{R}$ , which are continuous on  $[0,1]$  and differentiable on  $(0,1)$ . Then for every  $f$  in S, there exists a  $c \in (0,1)$ , depending on  $f$ , such that :  
माना सभी फलनों  $f : [0,1] \rightarrow \mathbb{R}$ , जो कि  $[0,1]$  पर संतत हैं तथा  $(0,1)$  पर अवकलनीय हैं, का समुच्चय S है। तो S में प्रत्येक  $f$  के लिए  $f$  पर निर्भर एक  $c \in (0,1)$  का अस्तित्व इस प्रकार है कि :

$$(1) \frac{f(1) - f(c)}{1 - c} = f'(c)$$

$$(2) |f(c) - f(1)| < (1 - c) |f'(c)|$$

$$(3) |f(c) - f(1)| < |f'(c)|$$

$$(4) |f(c) + f(1)| < (1+c)|f'(c)|$$

**Sol. BONUS**

- 14.** The mean and variance of 20 observations are found to be 10 and 4, respectively. On rechecking, it was found that an observation 9 was incorrect and the correct observation was 11. Then the correct variance is :

20 प्रेक्षणों के माध्य तथा प्रसरण क्रमशः 10 तथा 4 पाये गये। पुनः जांच करने पर पाया गया कि एक प्रेक्षण 9 गलत था तथा सही प्रेक्षण 11 था। तो सही प्रसरण है :

$$(1) 4.01$$

$$(2) 4.02$$

$$(3) 3.98$$

$$(4) 3.99$$

**Sol. 4**

Let 20 observation be  $x_1, x_2, \dots, x_{20}$

$$\text{given Mean} = \frac{x_1 + x_2 + \dots + x_{20}}{20} = 10$$

$$x_1 + x_2 + \dots + x_{20} = 200$$

$$\text{Now, } x_1 + x_2 + \dots + x_{20} - 9 + 11 \Rightarrow 202$$

$$V_{ar} = \frac{\sum x_i^2}{n} - (\bar{x})^2$$

$$4 = \frac{x_1^2 + x_2^2 + \dots + x_{20}^2}{20} - 10^2$$

$$2080 = x_1^2 + x_2^2 + \dots + x_{20}^2$$

**24000+**  
SELECTIONS SINCE 2007

JEE (Advanced)

**5392**

JEE (Main)

**16241**

(Under 50000 Rank)

NEET / AIIMS

**1305**

(since 2016)

NTSE / OLYMPIADS

**1158**

(5th to 10th class)

H.O. : 394, Rajeev Gandhi Nagar, Kota

**Toll Free : 1800-212-1799**

www.motion.ac.in | ✉: info@motion.ac.in

Now,  $x_1^2 + x_2^2 + \dots + x_{20}^2 - 81 + 121 \Rightarrow 2080 + 40 = 2120$   
new variance will be

$$\frac{2120}{20} - \left(\frac{202}{20}\right)^2 = 3.99$$

15. The area (in sq. units) of the region  $\{(x,y) \in \mathbb{R}^2 : x^2 \leq y \leq 3 - 2x\}$ , is :  
क्षेत्र  $\{(x,y) \in \mathbb{R}^2 : x^2 \leq y \leq 3 - 2x\}$  का क्षेत्रफल (वर्ग इकाई में) है :

- (1)  $\frac{31}{3}$                       (2)  $\frac{34}{3}$                       (3)  $\frac{29}{3}$                       (4)  $\frac{32}{3}$

Sol. 4

$$\begin{aligned} y &= x^2 \text{ \& } y = 3 - 2x \\ x^2 &= 3 - 2x \\ x^2 + 2x - 3 &= 0 \\ x^2 + 3x - x - 3 &= 0 \\ x(x + 3) - (x - 3) &= 0 \\ (x - 1)(x + 3) &= 0 \\ x &= 1, -3 \end{aligned}$$

$$\text{required area} = \int_{-3}^1 (\text{line} - \text{parabola}) dx$$

$$= \int_{-3}^1 [(3 - 2x) - x^2]$$


$$= \left( 3x - \frac{2x^2}{2} - \frac{x^3}{3} \right)_{-3}^1$$

$$\text{Area} = \left( 3x - x^2 - \frac{x^3}{3} \right)_{-3}^1$$

$$= \left[ 3(1) - (1)^2 - \frac{1}{3} \right] - \left[ 3(-3) - (-3)^2 - \frac{(-3)^3}{3} \right]$$

$$= \left( 2 - \frac{1}{3} \right) - (-18 + 9)$$

$$= \frac{5}{3} + 9 = \frac{32}{3}$$



**Increase Your Score  
for JEE Main April'2020**

**उत्कर्ष**  
15 JAN 2020

percentile between 97.0 to 98.99  
in JEE Main (Jan-2020)

Fees - ₹ 22000 Including GST

**उन्नति**  
17 JAN 2020

Below 97 percentile in JEE Main (Jan-2020)  
Tonure: 62 Days | Schedule: 5 Classes Per Day

Fees - ₹ 27500 Including GST

**उत्थान**  
17 JAN 2020

99 percentile and above  
in JEE Main (Jan-2020)

Fees - ₹ 11000 score 160-200    Fees - ₹ 5500 score 200-240    Fees - ₹ 0 score above 240

16. The length of the perpendicular from the origin, on the normal to the curve,  $x^2 + 2xy - 3y^2 = 0$  at the point (2,2) is

वक्र  $x^2 + 2xy - 3y^2 = 0$  के बिन्दु (2, 2) पर खींचे गये अभिलम्ब पर मूल बिन्दु से डाले गये लम्ब की लम्बाई है :

- (1) 2                                      (2)  $\sqrt{2}$                                       (3)  $2\sqrt{2}$                                       (4)  $4\sqrt{2}$

Sol. 3

$$\begin{aligned} x^2 + 2xy - 3y^2 &= 0 \\ 2x + 2y + 2xy' - 6yy' &= 0 \\ x + y + xy' - 3yy' &= 0 \\ y'(x - 3y) &= -(x + y) \end{aligned}$$

$$\frac{dy}{dx} = \frac{x+y}{3y-x}$$

$$\text{Slope of (N), } -\frac{dx}{dy} = \frac{x-3y}{x+y}$$

$$\therefore \left( -\frac{dx}{dy} \right)_{(2,2)} = \frac{2-6}{2+2} = -1$$

$\therefore$  equation of normal at (2,2) is

$$y - 2 = -(x - 2)$$

$$x + y - 4 = 0$$

$\therefore$  distance from (0,0) will be

$$p = \left| \frac{0+0-4}{\sqrt{2}} \right| = 2\sqrt{2}$$

17. Let S be the set of all real roots of the equation,  $3^x(3^x - 1) + 2 = |3^x - 1| + |3^x - 2|$ . Then S :

- (1) is a singleton                                      (2) contains at least four elements.  
(3) contains exactly two elements                                      (4) is an empty set.

माना समीकरण  $3^x(3^x - 1) + 2 = |3^x - 1| + |3^x - 2|$  के सभी वास्तविक मूलों का समुच्चय S है। तो S :

- (1) एक ही अवयव वाला समुच्चय है।                                      (2) में कम से कम चार अवयव हैं।  
(3) में मात्र दो अवयव हैं                                      (4) एक रिक्त समुच्चय है

Sol. 1

$$3^x(3^x - 1) + 2 = |3^x - 1| + |3^x - 2|$$

$$\text{put } 3^x = t$$

$$t(t - 1) + 2 = |t - 1| + |t - 2|$$

$$t^2 - t + 2 = |t - 1| + |t - 2|$$

from graph


let  $\alpha$  is real solutions

$$\alpha = 3^x$$

$$x = \log_3 \alpha$$

$\therefore$  only one solution

$\therefore$  singleton set



18. If  $A = \begin{pmatrix} 2 & 2 \\ 9 & 4 \end{pmatrix}$  and  $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ , then  $10A^{-1}$  is equal to :

यदि  $A = \begin{pmatrix} 2 & 2 \\ 9 & 4 \end{pmatrix}$  तथा  $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  हैं, तो  $10A^{-1}$  बराबर है :

- (1)  $A - 6I$                       (2)  $4I - A$                       (3)  $6I - A$                       (4)  $A - 4I$

Sol. 1

$$A = \begin{bmatrix} 2 & 2 \\ 9 & 4 \end{bmatrix} \text{ and } I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$10A^{-1} = ?$$

According to Cayley Hamilton equation

$$|A - \lambda I| = 0$$

$$\begin{vmatrix} 2-\lambda & 2 \\ 9 & 4-\lambda \end{vmatrix} = 0$$

$$(2 - \lambda)(4 - \lambda) - 18 = 0$$

$$8 - 2\lambda - 4\lambda + \lambda^2 - 18 = 0$$

$$\lambda^2 - 6\lambda - 10 = 0$$

$$\therefore A^2 - 6A - 10I = 0$$

$$A^{-1}(A^2 - 6A - 10I) = 0$$

$$A - 6I - 10A^{-1} = 0$$

$$10A^{-1} = A - 6I$$

19. The mirror image of the point  $(1,2,3)$  in a plane is  $\left(-\frac{7}{3}, -\frac{4}{3}, -\frac{1}{3}\right)$ . Which of the following point lies on this plane ?

बिन्दु  $(1,2,3)$  का एक समतल में प्रतिबिम्ब (mirror image),  $\left(-\frac{7}{3}, -\frac{4}{3}, -\frac{1}{3}\right)$  है। निम्न में से कौन सा बिन्दु इस समतल

पर स्थित है ?

- (1)  $(1,1,1)$                       (2)  $(1,-1,1)$                       (3)  $(-1,-1,-1)$                       (4)  $(-1,-1,1)$

Sol. 2

for required plane

$$\vec{n} \parallel \overline{AB}$$

$$\vec{n} = -\frac{7}{3} - 1, \frac{-4}{3} - 2, \frac{-1}{3} - 3$$

$$\vec{n} = \frac{-10}{3}, \frac{-10}{3}, \frac{-10}{3}$$

$$\text{D.r of } \vec{n} = 1,1,1$$

**Increase Your Score  
for JEE Main April'2020**

**उत्कर्ष**  
15 JAN 2020

percentile between 97.0 to 98.99  
in JEE Main (Jan-2020)

Fees - ₹ 22000 Including GST

**उन्नति**  
17 JAN 2020

Below 97 percentile in JEE Main (Jan-2020)  
Tenure: 62 Days | Schedule: 5 Classes Per Day

Fees - ₹ 27500 Including GST

**उत्थान**  
17 JAN 2020

99 percentile and above  
in JEE Main (Jan-2020)

Fees - ₹ 11000  
score 160-200

Fees - ₹ 5500  
score 200-240

Fees - ₹ 0  
score above 240

Also mid- point of A & B is  $M \equiv \left( \frac{-7+1}{2}, \frac{-4+2}{2}, \frac{-1+3}{2} \right)$

$$M \equiv \left( \frac{-4}{6}, \frac{2}{6}, \frac{8}{6} \right)$$


$$M \equiv \left( \frac{-2}{3}, \frac{1}{3}, \frac{4}{3} \right)$$

∴ equation of required plane B

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

$$\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = -\frac{2}{3} + \frac{1}{3} + \frac{4}{3}$$

$$\boxed{x + y + z = 1}$$



- 20.** Let A and B be two events such that the probability that exactly one of them occurs is  $\frac{2}{5}$  and the probability that A or B occurs is  $\frac{1}{2}$ , then the probability of both of them occur together is :

माना A तथा B दो घटनाएँ इस प्रकार हैं कि दोनों में से मात्र एक के होने की प्रायिकता  $\frac{2}{5}$  है तथा A या B के होने की प्रायिकता

$\frac{1}{2}$  है, तो दोनों के एक साथ होने की प्रायिकता है :

- (1) 0.02                      (2) 0.01                      (3) 0.20                      (4) 0.10

**Sol.**

$$P(\text{exactly one}) = \frac{2}{5}$$

$$P(A) + P(B) - 2P(A \cap B) = \frac{2}{5} \quad \dots(1)$$

$$P(A \text{ or } B) = P(A \cup B) = \frac{1}{2}$$

$$P(A) + P(B) - P(A \cap B) = \frac{1}{2} \quad \dots(2)$$

$$P(A \cap B) = ?$$

Solve (1) - (2)

$$-P(A \cap B) = \frac{2}{5} - \frac{1}{2}$$

$$P(A \cap B) = \frac{1}{2} - \frac{2}{5}$$

$$= \frac{5}{10} - \frac{4}{10} = \frac{1}{10}$$

$$= P(A \cap B) = 0.10$$

- 21.** The number of 4 letters words (with or without meaning) that can be formed from the eleven letters of the word 'EXAMINATION' is .....
- शब्द 'EXAMINATION' के ग्यारह अक्षरों से बन सकने वाले 4 अक्षरों के शब्दों (अर्थ वाले तथा अर्थविहीन) की संख्या है.....।

**Sol. 2454**

EXAMINATION

(AA)(II)(NN)(EXMOT)

to form four letter words

(1) All same .....not possible

(2) 1 different, 3 same.....not possible

(3) 2 different, 2 same..... ${}^3C_1 \times {}^7C_2 \times \frac{4!}{2!} = 3 \times \frac{7 \times 6 \times 5!}{2!5!} \times \frac{4 \times 3 \times 2!}{2!} = 63 \times 12 = 756$

(4) 2 same, 2 same..... ${}^3C_2 \times \frac{4!}{2!2!} = 3 \times \frac{4 \times 3 \times 2!}{2.2} = 18$

(5) All different ..... ${}^8C_4 \times 4! = \frac{8.7.6.5.4!}{4.3.2.4!} \times 4! = 56 \times 30 = 1680$

**Total = 2454**

- 22.** If  $\frac{\sqrt{2} \sin \alpha}{\sqrt{1 + \cos 2\alpha}} = \frac{1}{7}$  and  $\sqrt{\frac{1 - \cos 2\beta}{2}} = \frac{1}{\sqrt{10}}$ ,  $\alpha, \beta \in \left(0, \frac{\pi}{2}\right)$ , then  $\tan(\alpha + 2\beta)$  is equal to :


यदि  $\frac{\sqrt{2} \sin \alpha}{\sqrt{1 + \cos 2\alpha}} = \frac{1}{7}$  तथा  $\sqrt{\frac{1 - \cos 2\beta}{2}} = \frac{1}{\sqrt{10}}$ ,  $\alpha, \beta \in \left(0, \frac{\pi}{2}\right)$ , हैं, तो  $\tan(\alpha + 2\beta)$  बराबर है.....।

**Sol. 1**

$$\frac{\sqrt{2} \sin \alpha}{\sqrt{1 + \cos 2\alpha}} = \frac{1}{7} \text{ and } \sqrt{\frac{1 - \cos 2\beta}{2}} = \frac{1}{\sqrt{10}}$$

$$\frac{\sqrt{2} \sin \alpha}{\sqrt{2} \cos \alpha} = \frac{1}{7} \text{ \& \ } \frac{\sqrt{2} \sin \beta}{\sqrt{2}} = \frac{1}{\sqrt{10}}$$

$$\tan \alpha = \frac{1}{7} \quad \sin \beta = \frac{1}{\sqrt{10}}$$



**Increase Your Score  
for JEE Main April'2020**

**उत्कर्ष**  
15 JAN 2020

percentile between 97.0 to 98.99  
in JEE Main (Jan-2020)

Fees - ₹ 22000 Including GST

**उन्नति**  
17 JAN 2020

Below 97 percentile in JEE Main (Jan-2020)  
Tenure: 62 Days | Schedule: 5 Classes Per Day

Fees - ₹ 27500 Including GST

**उत्थान**  
17 JAN 2020

99 percentile and above  
in JEE Main (Jan-2020)

Fees - ₹ 11000 score 160-200 | Fees - ₹ 5500 score 200-240 | Fees - ₹ 0 score above 240



$$\tan \beta = \frac{1}{3}$$

$$\therefore \tan 2\beta = \frac{2 \tan \beta}{1 - \tan^2 \beta} = \frac{2\left(\frac{1}{3}\right)}{1 - \frac{1}{9}} = \frac{2/3}{8/9}$$

$$\tan 2\beta = \frac{2}{3} \cdot \frac{9}{8} = \frac{3}{4}$$

$$= \tan(\alpha + 2\beta)$$

$$= \frac{\tan \alpha + \tan 2\beta}{1 - \tan \alpha \cdot \tan 2\beta}$$

$$= \frac{\frac{1}{7} + \frac{3}{4}}{1 - \frac{1}{7} \cdot \frac{3}{4}} = \frac{\frac{4+21}{28}}{\frac{28-3}{28}} = \frac{25}{25} = 1$$

23. The sum  $\sum_{n=1}^7 \frac{n(n+1)(2n+1)}{4}$  is equal to :

योगफल  $\sum_{n=1}^7 \frac{n(n+1)(2n+1)}{4}$  बराबर है .....

Sol. 504

$$\sum_{n=1}^7 \frac{n(n+1)(2n+1)}{4}$$

$$\frac{1}{4} \sum_{n=1}^7 n(n+1)(2n+1)$$

$$\frac{1}{4} \sum_{n=1}^7 ((n^2 + n)(2n+1))$$

$$= \frac{1}{4} \sum_{n=1}^7 (2n^3 + 3n^2 + n)$$

$$\frac{1}{2} \sum_{n=1}^7 n^3 + \frac{3}{4} \sum_{n=1}^7 n^2 + \frac{1}{4} \sum_{n=1}^7 n$$

$$\Rightarrow \frac{1}{2} \left( \frac{7(7+1)}{2} \right)^2 + \frac{3}{4} \left( \frac{7(7+1)(14+1)}{6} \right) + \frac{1}{4} \cdot \frac{7(8)}{2}$$

$$= \frac{1}{2} \frac{49.8.8}{4} + \frac{3.7.8.15}{4.6} + \frac{1}{4} \frac{7.8}{2}$$

$$= (49)(8) + (15 \times 7) + (7)$$

$$= 392 + 105 + 7 = 504$$

24. Let  $f(x)$  be a polynomial of degree 3 such that  $f(-1) = 10$ ,  $f(1) = -6$ ,  $f(x)$  has a critical point at  $x = -1$  and  $f'(x)$  has a critical point at  $x = 1$ . Then  $f(x)$  has a local minima at  $x = \dots\dots\dots$   
 माना घात 3 का एक बहुपद  $f(x)$  इस प्रकार है कि  $f(-1) = 10$ ,  $f(1) = -6$ ,  $f(x)$  का एक क्रांतिक बिन्दु  $x = -1$  है तथा  $f'(x)$  का एक क्रांतिक बिन्दु  $x = 1$  है। तो  $f(x)$  का एक स्थानीय निम्ननिष्ठ है  $x = \dots\dots\dots$  है।

Sol. 3

$$f(x) = ax^3 + bx^2 + cx + d$$

$$f(-1) = 10, f(1) = -6$$

$$-a + b - c + d = 10 \dots(i)$$

$$a + b + c + d = -6 \dots(ii)$$

add (i) + (ii)

$$2(b + d) = 4$$

$$b + d = 2 \dots(iii)$$

$$f'(x) = 3ax^2 + 2bx + c$$

$$f'(-1) = 0$$

$$3a - 2b + c = 0 \dots(iv)$$

$$f''(x) = 6ax + 2b$$

$$f''(1) = 0$$

$$6a + 2b = 0 \dots(v)$$

add (iv) + (v)

$$9a + c = 0 \dots(vi)$$

$$b = -3a$$

$$c + 9 \left( \frac{-b}{3} \right) = 0$$

$$c = 3b$$

$$f(x) = \frac{-b}{3} x^3 + bx^2 + 3bx + (2 - b)$$

$$f'(x) = -bx^2 + 2bx + 3b = 0$$

$$x^2 - 2x - 3 = 0$$

$$(x - 3)(x + 1) = 0$$

$$x = 3, -1 \text{ Minima}$$

at  $x = 3$

25. Let a line  $y = mx(m > 0)$  intersect the parabola,  $y^2 = x$  at a point P, other than the origin. Let the tangent to it at P meet the x - axis at the point Q. If area ( $\Delta OPQ$ ) = 4 sq. units, then m is equal to  $\dots\dots\dots$

माना एक रेखा  $y = mx(m > 0)$ , परवलय  $y^2 = x$  को मूल बिन्दु के अतिरिक्त एक बिन्दु P पर काटती है। माना P पर इसकी स्पर्श रेखा x-अक्ष को बिन्दु Q पर मिलती है। यदि ( $\Delta OPQ$ ) का क्षेत्रफल 4 वर्ग इकाई है, तो m बराबर है  $\dots\dots\dots$ ।

**Increase Your Score  
for JEE Main April'2020**

**उत्कर्ष**  
15 JAN 2020

percentile between 97.0 to 98.99  
in JEE Main (Jan-2020)

Fees - ₹ 22000 Including GST

**उन्नति**  
17 JAN 2020

Below 97 percentile in JEE Main (Jan-2020)  
Tenure: 62 Days | Schedule: 5 Classes Per Day

Fees - ₹ 27500 Including GST

**उत्थान**  
17 JAN 2020

99 percentile and above  
in JEE Main (Jan-2020)

Fees - ₹ 11000 score 160-200 | Fees - ₹ 5500 score 200-240 | Fees - ₹ 0 score above 240

Sol. 0.5

let  $p(t^2, t)$   
Tangent at  $P(t^2, t)$

$$ty = \frac{x + t^2}{2}$$

$2ty = x + t^2 \rightarrow$  equation of tangent

$$Q \equiv (-t^2, 0)$$

$O(0,0)$

$$\Delta(OPQ) = 4$$

$$\frac{1}{2} \begin{vmatrix} 0 & 0 & 1 \\ t^2 & t & 1 \\ -t^2 & 0 & 1 \end{vmatrix} = 4$$


$$|t^3| = 8$$

$$t = 2 \quad (\because t > 0)$$

$\therefore 4y = x + 4$  is tangent

$\therefore P$  is  $(4,2)$

$$y = mx \Rightarrow 2 = 4m \Rightarrow m = \frac{1}{2} \Rightarrow 0.5$$



**24000+**  
SELECTIONS SINCE 2007

JEE (Advanced)

**5392**

JEE (Main)

**16241**

(Under 50000 Rank)

NEET / AIIMS

**1305**

(since 2016)

NTSE / OLYMPIADS

**1158**

(5th to 10th class)

H.O. : 394, Rajeev Gandhi Nagar, Kota

**Toll Free : 1800-212-1799**

www.motion.ac.in | ✉: info@motion.ac.in

कर लो अब पूरी तैयारी

चूक ना जाये इस बारी

## INCREASE YOUR SCORE for JEE Main April 2020

**उत्थान** 17<sup>th</sup> JAN 2020

99 percentile and above  
in JEE Main (Jan-2020)

**उत्कर्ष** 15<sup>th</sup> JAN 2020

percentile between 97.0 to 98.99  
in JEE Main (Jan-2020)

**उन्नति** 17<sup>th</sup> JAN 2020

Below 97 percentile  
in JEE Main (Jan-2020)

**MOTION**<sup>TM</sup>

Nurturing potential through education

Toll Free : 1800-212-1799