

हमारा विश्वास... हर एक विद्यार्थी है खास

**JEE  
MAIN  
Sept.  
2020**

**QUESTION PAPER WITH SOLUTION**

**CHEMISTRY \_ 2 Sep. \_ SHIFT - 1**




**MOTION™**

H.O. : 394, Rajeev Gandhi Nagar, Kota  
www.motion.ac.in | ✉: info@motion.ac.in

हमारा विश्वास... हर एक विद्यार्थी है खास


1. The increasing order of the following compounds towards HCN addition is:



- (1) (iii) < (i) < (iv) < (ii)                      (2) (iii) < (iv) < (i) < (ii)  
 (3) (i) < (iii) < (iv) < (ii)                      (4) (iii) < (iv) < (ii) < (i)

Sol. 1

In HCN, CN<sup>-</sup> is acts as nucleophile, attack first that -CHO group which has maximum positive charge. The magnitude of the (+ve) charge increases by -M and -I group. So reactivity order will be



So, option (1) is correct answer.

2. Which of the following is used for the preparation of colloids?

- (1) Van Arkel Method                                      (2) Ostwald Process  
 (3) Mond Process    (4) Bredig's Arc Method


Sol. 4

Bredig's Arc method  
 Chapter name surface chemistry

3. An open beaker of water in equilibrium with water vapour is in a sealed container. When a few grams of glucose are added to the beaker of water, the rate at which water molecules:

- (1) leaves the vapour increases                      (2) leaves the solution increases  
 (3) leaves the vapour decreases                      (4) leaves the solution decreases

Sol. 1



Hence Rate at which water molecules leaves the vap. increases.

**CRASH COURSE**  
**FOR JEE ADVANCED 2020**

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

- ◆ Doubt Support ◆ Advanced Level Test Access  
 ◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

4. For octahedral Mn(II) and tetrahedral Ni(II) complexes, consider the following statements:

- (I) both the complexes can be high spin.  
 (II) Ni(II) complex can very rarely be low spin.  
 (III) with strong field ligands, Mn(II) complexes can be low spin.  
 (IV) aqueous solution of Mn(II) ions is yellow in colour.

The correct statements are:

- (1) (I), (III) and (IV) only    (2) (I), (II) and (III) only  
 (3) (II), (III) and (IV) only    (4) (I) and (II) only


**Sol. 2**

Mn<sup>2+</sup> [Ar]3d<sup>5</sup> it can form low spin as well as high spin complex depending upon nature of ligand same of Ni<sup>2+</sup> ion with coordination no 4. It can be dsp<sup>2</sup> or sp<sup>3</sup> i:e low spin or high spin depending open nature of ligand.

5. The statement that is not true about ozone is:



- (1) in the stratosphere, it forms a protective shield against UV radiation.  
 (2) in the atmosphere, it is depleted by CFCs.  
 (3) in the stratosphere, CFCs release chlorine free radicals (Cl) which reacts with O<sub>3</sub> to give chlorine dioxide radicals.  
 (4) it is a toxic gas and its reaction with NO gives NO<sub>2</sub>.

**Sol. 3**



Hence option (3)

6. Consider the following reactions:






'x', 'y' and 'z' in these reactions are respectively.

- (1) 4, 5 & 5    (2) 5, 4 & 5  
 (3) 5, 6 & 5    (4) 4, 6 & 5

हमारा विश्वास... हर एक विद्यार्थी है खास

Sol. 4



$(\text{CH}_3\text{CO})_2\text{O}$  reacts with  $-\text{OH}$  group to form acetyl derivative, so as the no. of  $-\text{OH}$  group no. of eq. of  $(\text{CH}_3\text{CO})_2\text{O}$  will be used


So,  $x = 4$

$y = 6$

$z = 5$

So, option (4) will be correct answer.

7. The IUPAC name for the following compound is:



(1) 2,5-dimethyl-5-carboxy-hex-3-enal  
(3) 6-formyl-2-methyl-hex-3-enoic acid

(2) 2,5-dimethyl-6-oxo-hex-3-enoic acid  
(4) 2,5-dimethyl-6-carboxy-hex-3-enal

**CRASH COURSE**  
FOR JEE ADVANCED 2020


FREE Online Lectures Available on 

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access  
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

Sol. 2



2,5-Dimethyl-6-oxohex-3-enoic acid

8. For the following Assertion and Reason, the correct option is

**Assertion (A):** When Cu (II) and sulphide ions are mixed, they react together extremely quickly to give a solid.

**Reason (R):** The equilibrium constant of  $\text{Cu}^{2+}(\text{aq}) + \text{S}^{2-}(\text{aq}) \rightleftharpoons \text{CuS}(\text{s})$  is high because the solubility product is low.

- (1) (A) is false and (R) is true.
- (2) Both (A) and (R) are false.
- (3) Both (A) and (R) are true but (R) is not the explanation for (A).
- (4) Both (A) and (R) are true but (R) is the explanation for (A).





Sol. 4

(A) is (B) true &

(R) is correct explanation of (A)

Ans. 4

9. Which one of the following graphs is not correct for ideal gas?



d = Density, P = Pressure, T = Temperature

- (1) I                      (2) IV                      (3) III                      (4) II

**CRASH COURSE**  
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access  
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**


हमारा विश्वास... हर एक विद्यार्थी है खास

**Sol. 4**


For ideal Gas

$$d = \frac{P \times M}{RT}$$


$d \propto T \rightarrow$  Hyperbolic



$d \propto \frac{1}{T} \rightarrow$  St. line



$d \propto p \rightarrow$  St line



$\therefore$  'II' Graph is incorrect  
Ans (4)

**10.** While titrating dilute HCl solution with aqueous NaOH, which of the following will not be required?



- (1) Bunsen burner and measuring cylinder      (2) Burette and porcelain tile  
(3) Clamp and phenolphthalein                (4) Pipette and distilled water

**Sol. 1**

Bunsen Burner & measuring cylinder are not Required. As titration is already an exothermic process

Ans.(1)

**11.** In Carius method of estimation of halogen, 0.172 g of an organic compound showed presence of 0.08 g of bromine. Which of these is the correct structure of the compound?

- (1)  $H_3C-Br$       (2)       (3)       (4)  $H_3C-CH_2-Br$

**CRASH COURSE**  
FOR JEE ADVANCED 2020

FREE Online Lectures Available on 

Go Premium at ₹ 1100

- ◆ Doubt Support ◆ Advanced Level Test Access
- ◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

Sol. 3

carius method

$$\text{mass \% of 'Br'} = \frac{0.08}{0.172} \times 100 = \frac{8000}{172} = 46.51\%$$

$$\text{option (1) mass \%} = \frac{80}{95} \times 100$$

$$\text{(2) mass \%} = \frac{2 \times 80 \times 100}{252}$$

$$\text{(3) mass \%} = \frac{1 \times 80 \times 100}{80 + 72 + 6 + 14} = \frac{8000}{172} \%$$

$$\text{(4) mass \%} = \frac{1 \times 80 \times 100}{109} \%$$






Option (3) matches with the given mass percentage value

Ans (3)



12. On heating compound (A) gives a gas (B) which is a constituent of air. This gas when treated with  $H_2$  in the presence of a catalyst gives another gas (C) which is basic in nature. (A) should not be:  
 (1)  $(NH_4)_2Cr_2O_7$       (2)  $NaN_3$       (3)  $NH_4NO_2$       (4)  $Pb(NO_3)_2$

Sol. 4

The gas (B) is  $N_2$  which is found in air



13. The major product in the following reaction is:



**CRASH COURSE**  
**FOR JEE ADVANCED 2020**

FREE Online Lectures Available on 


Go Premium at ₹ 1100

◆ Doubt Support    ◆ Advanced Level Test Access  
 ◆ Live Test Paper Discussion    ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

हमारा विश्वास... हर एक विद्यार्थी है खास

Sol. 3



Option (3) is correct answer.





14. In general, the property (magnitudes only) that shows an opposite trend in comparison to other properties across a period is:

- |                         |                            |
|-------------------------|----------------------------|
| (1) Ionization enthalpy | (2) Electronegativity      |
| (3) Atomic radius       | (4) Electron gain enthalpy |

Sol. 3

Ionisation energy, electronegativity & electron gain enthalpy increase across a period but atomic radius decreases

15. The figure that is not a direct manifestation of the quantum nature of atoms is:



**CRASH COURSE**  
FOR JEE ADVANCED 2020

FREE Online Lectures Available on [YouTube](#)

Go Premium at ₹ 1100


◆ Doubt Support ◆ Advanced Level Test Access  
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**








Sol. 4

Internal energy of 'Ar' or any gas, has nothing to do with Quantum nature of atom hence




Ans. option (4)

16. The major aromatic product C in the following reaction sequence will be :



Sol. 3



Option (3) is correct answer.

**CRASH COURSE**  
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100


◆ Doubt Support ◆ Advanced Level Test Access  
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

हमारा विश्वास... हर एक विद्यार्थी है खास

- 17.** Consider that a  $d^6$  metal ion ( $M^{2+}$ ) forms a complex with aqua ligands, and the spin only magnetic moment of the complex is 4.90 BM. The geometry and the crystal field stabilization energy of the complex is:
- (1) tetrahedral and  $-0.6\Delta_t$  (2) tetrahedral and  $-1.6\Delta_t + 1P$   
 (3) octahedral and  $-1.6\Delta_o$  (4) octahedral and  $-2.4\Delta_o + 2P$

**Sol. 1**




$$\mu_{\text{spin}} = 4.9 \text{ BM}$$

$$\text{CFSE} = -0.4 \times 4\Delta_o + 0.6 \times 2\Delta_o$$

$$= [1.6 + 1.2] \Delta_o$$

$$= -0.4 \Delta_o$$



$$\text{CFSE} = -0.6 \times 3\Delta_t + 0.4 \times 3\Delta_t$$

$$= -1.8 \Delta_t + 1.2 \Delta_t$$


$$= -0.6 \Delta_t$$

- 18.** If  $AB_4$  molecule is a polar molecule, a possible geometry of  $AB_4$  is:
- (1) Square planar (2) Tetrahedral  
 (3) Square pyramidal (4) Rectangular planar

**Sol. 1**

Incorrect question Option 1 is more appropriate with respect to given option (Chemical bonding)  
 (Options are incorrect)

- 19.** Which of the following compounds will show retention in configuration on nucleophilic substitution by  $\text{OH}^-$  ion?



**CRASH COURSE**  
**FOR JEE ADVANCED 2020**

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access  
 ◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

**Sol. 1**

In  $\text{CH}_3\text{-CH}(\text{C}_2\text{H}_5)\text{-CH}_2\text{Br}$  attack of  $\text{OH}^-$  is not on chiral carbon, it is adjacent to chiral carbon, so configuration of chiral carbon remains constant.

**20.** The metal mainly used in devising photoelectric cells is:  
 (1) Li (2) Cs (3) Rb (4) Na

**Sol. 2**  
 'Cs' is used in photoelectric cell as its ionisation energy is lowest  
 Hence Ans (2)

**21.** The mass of gas adsorbed,  $x$ , per unit mass of adsorbate,  $m$ , was measured at various pressures,  $p$ .  
 A graph between  $\log \frac{x}{m}$  and  $\log p$  gives a straight line with slope equal to 2 and the intercept equal to 0.4771. The value of  $\frac{x}{m}$  at a pressure of 4 atm is: (Given  $\log 3 = 0.4771$ )

**Sol. 48**

$$\frac{x}{m} = kP^{1/n}$$

$$\log(x/m) = \log_{(k)} + \frac{1}{n} \log(p)$$

$$y = c + mx$$

$$\text{Intercept } C = \log_k = 0.4771$$

$$\text{slop} = \frac{1}{n} = 2, k = 3$$


$$\begin{aligned} \frac{x}{m} &= k(P)^{1/n} \quad \text{at } P = 4 \text{ atm} \\ &= 3(4)^2 \end{aligned}$$

$$\frac{x}{m} = 3 \times 16 = 48 \text{ Ans}$$

**22.** The Gibbs energy change (in J) for the given reaction at  $[\text{Cu}^{2+}] = [\text{Sn}^{2+}] = 1 \text{ M}$  and 298 K is:  
 $\text{Cu(s)} + \text{Sn}^{2+}(\text{aq.}) \rightarrow \text{Cu}^{2+}(\text{aq.}) + \text{Sn(s)}$

$$(E_{\text{Sn}^{2+}|\text{Sn}}^{\circ} = -0.16\text{V}, E_{\text{Cu}^{2+}|\text{Cu}}^{\circ} = 0.34\text{V}, \text{Take } F = 96500 \text{ C mol}^{-1})$$

**Sol. 96500**



$$E_{\text{cell}}^{\circ} = -0.16 - 0.34$$

$$= -0.50$$

$$\begin{aligned} \Delta G^{\circ} &= -nF E_{\text{cell}}^{\circ} \\ &= -2 \times 96500 \times (-0.5) \\ &= +96500 \end{aligned}$$

$$\Delta G = \Delta G^{\circ} + RT \ln Q$$

$$= 96500 + \frac{25}{3} \times 298 \times 2.303 \log(1)$$

$$\Delta G = 96500 \text{ Joules}$$

**CRASH COURSE**  
**FOR JEE ADVANCED 2020**

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access  
 ◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

हमारा विश्वास... हर एक विद्यार्थी है खास


23. The internal energy change (in J) when 90 g of water undergoes complete evaporation at 100° C is \_\_\_\_\_.

(Given:  $\Delta H_{\text{vap}}$  for water at 373 K = 41 kJ/mol,  $R = 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$ )



Sol.  $\text{H}_2\text{O} (\ell) \longrightarrow \text{H}_2\text{O} (\text{g})$

$$\begin{aligned} \Delta E_{\text{vap}} &= \Delta H_{\text{vap}} - \Delta ngRT \\ &= 41000 \times 5 - 5 \times 8.314 \times 373 \\ &= 189494.39 \end{aligned}$$


24. The oxidation states of iron atoms in compounds (A), (B) and (C), respectively, are x, y and z. The sum of x, y and z is \_\_\_\_\_.




Sol. 6



25. The number of chiral carbons present in the molecule given below is \_\_\_\_\_.



Sol. 5



Total chiral carbon = 5

**CRASH COURSE**  
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access  
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

जब इन्होंने पूरा किया अपना सपना  
तो आप भी पा सकते है लक्ष्य अपना

Admission  
**OPEN**

## JEE MAIN RESULT 2019



**Nitin Gupta**

Marks  
**335**  
13th (2019)

Marks  
**149**  
12th (2018)



**Shiv Modi**

Marks  
**318**  
13th (2019)


Marks  
**153**  
12th (2018)



**Ritik Bansal**

Marks  
**308**  
13th (2019)

Marks  
**218**  
12th (2018)



**Shubham Kumar**

Marks  
**300**  
13th (2019)

Marks  
**153**  
12th (2018)

### KOTA'S PIONEER IN DIGITAL EDUCATION

**1,95,00,000+** viewers | **72,67,900+** viewing hours | **2,11,000+** Subscribers

SERVICES	SILVER	GOLD	PLATINUM
Classroom Lectures (VOD)			
Live interaction	NA		
Doubt Support	NA		
Academic & Technical Support	NA		
Complete access to all content	NA		
Classroom Study Material	NA		
Exercise Sheets	NA		
Recorded Video Solutions	NA		
Online Test Series	NA		
Revision Material	NA		
<b>Upgrade to Regular Classroom program</b>	<b>Chargeable</b>	<b>Chargeable</b>	<b>Free</b>
Physical Classroom	NA	NA	
Computer Based Test	NA	NA	
Student Performance Report	NA	NA	
Workshop & Camp	NA	NA	
Motion Solution Lab- Supervised learning and instant doubt clearance	NA	NA	
Personalised guidance and mentoring	NA	NA	

### FEE STRUCTURE

CLASS	SILVER	GOLD	PLATINUM
7th/8th	FREE	₹ 12,000	₹ 35,000
9th/10th	FREE	₹ 15,000	₹ 40,000
11th	FREE	₹ 29,999	₹ 49,999
12th	FREE	₹ 39,999	₹ 54,999
12th Pass	FREE	₹ 39,999	₹ 59,999

+ Student Kit will be provided at extra cost to Platinum Student.

- \* **SILVER (Trial)** Only valid 7 DAYS or First 10 Hour's Lectures.
- \*\* **GOLD (Online)** can be converted to regular classroom (Any MOTION Center) by paying difference amount after lockdown.
- \*\*\* **PLATINUM (Online + Regular)** can be converted to regular classroom (Any MOTION Center) without any cost after lockdown.

New Batch Starting from :  
**16 & 23 September 2020**

**Zero Cost EMI Available**

**MOTION™**

H.O. : 394, Rajeev Gandhi Nagar, Kota  
www.motion.ac.in | ✉ : info@motion.ac.in