

### QUESTION PAPER WITH SOLUTION

CHEMISTRY \_ 3 Sep. \_ SHIFT - 2











H.O.: 394, Rajeev Gandhi Nagar, Kota www.motion.ac.in



# **Motion**

- 1. The five successive ionization enthalpies of an element are 800, 2427, 3658, 25024 and 32824 kJ mol<sup>-1</sup>. The number of valence electrons in the element is:
  - (1) 2
- (2)4
- (3)3
- (4)5

Sol. 3

Fourth & Fifth I.E. are very high (periodic properties) indicating presence of three valence shell electrons

- **2.** The incorrect statement is:
  - (1) Manganate and permanganate ions are tetrahedral
  - (2) In manganate and permanganate ions, the  $\pi$ -bonding takes place by overlap of p-orbitals of oxygen and d-orbitals of manganese
  - (3) Manganate and permanganate ions are paramagnetic
  - (4) Manganate ion is green in colour and permanganate ion is purple in colour
- Sol. 3

<sup>+7</sup> Mn O<sub>4</sub>-

 $d^{o} \rightarrow dimagnetic$ 

Mn O<sub>4</sub>2-

 $d^1 \rightarrow Paramagnetic$ 

- **3.** Match the following drugs with their therapeutic actions:
  - (i) Ranitidine

- (a) Antidepressant
- (ii) Nardil (Phenelzine)
- (b) Antibiotic
- (iii) Chloramphenicol
- (c) Antihistamine
- (iv) Dimetane (Brompheniramine)
- (d) Antacid(e) Analgesic
- (1) (i)-(d); (ii)-(a); (iii)-(b); (iv)-(c)
- (2) (i)-(d); (ii)-(c); (iii)-(a); (iv)-(e)
- (3) (i)-(a); (ii)-(c); (iii)-(b); (iv)-(e)
- (4) (i)-(e); (ii)-(a); (iii)-(c); (iv)-(d)

- Sol. 1
- **4.** An ionic micelle is formed on the addition of:
  - (1) liquid diethyl ether to aqueous NaCl solution
  - (2) sodium stearate to pure toluene

(3) excess water to liquid 
$$H_3C \nearrow N \longrightarrow CH_3$$
  $CH_3$ 

- (4) excess water to liquid  $H_3C$   $N \cap CH_3$
- Sol. 3

ionic micelles formed by addition of water to soap {sodium stearate}

- **5.** Among the statements (I–IV), the correct ones are:
  - (I) Be has smaller atomic radius compared to Mg.
  - (II) Be has higher ionization enthalpy than Al.
  - (III) Charge/radius ratio of Be is greater than that of Al.
  - (IV) Both Be and Al form mainly covalent compounds.
  - (1) (I), (II) and (IV)

(2) (I), (II) and (III)

(3) (II), (III) and (IV)

(4) (I), (III) and (IV)

Sol. 1

Refer S-Block

### **CRASH COURSE**

**FOR JEE ADVANCED 2020** 

FREE Online Lectures Available on You Tube

Go Premium at ₹ 1100

Doubt Support ◆ Advanced Level Test Access

◆ Live Test Paper Discussion ◆ Final Revision Exercises

## **Motion**

Complex A has a composition of  $H_{12}O_6Cl_3Cr$ . If the complex on treatment with conc. $H_2SO_4$  loses 13.5% of its original mass, the correct molecular formula of A is:

[Given: atomic mass of Cr = 52 amu and Cl = 35 amu]

(1) [Cr(H<sub>2</sub>O)<sub>5</sub>Cl]Cl<sub>2</sub>.H<sub>2</sub>O

(2) [Cr(H<sub>2</sub>O)<sub>4</sub>Cl<sub>2</sub>]Cl.2H<sub>2</sub>O

(3)  $[Cr(H_2O)_3Cl_3].3H_2O$ 

 $(4) [Cr(H_2O)_6]CI_3$ 

Sol. 2

Let x molecule of water are lost then

$$13.5 = \left\lceil \frac{x \times 18}{6 \times 18 + 3 \times 35 + 52} \right\rceil \times 100$$

 $x = 1.99 \approx 2$ 

so, complex is [Cr(H<sub>2</sub>O)<sub>4</sub>Cl<sub>2</sub>].2H<sub>2</sub>O

7. The decreasing order of reactivity of the following compounds towards nucleophilic substitution  $(S_N 2)$  is:

- (1) (III) > (II) > (IV) > (I)
- (2) (IV) > (II) > (III) > (I)
- (3)(II) > (III) > (IV) > (I)
- (4) (II) > (III) > (I) > (IV)

Sol. 3

- **8.** The increasing order of the reactivity of the following compounds in nucleophilic addition reaction is: Propanal, Benzaldehyde, Propanone, Butanone
  - (1) Benzaldehyde < Propanal < Propanone < Butanone
  - (2) Propanal < Propanone < Butanone < Benzaldehyde
  - (3) Butanone < Propanone < Benzaldehyde < Propanal
  - (4) Benzaldehyde < Butanone < Propanone < Propanal
- Sol. 3

Rate of Nucleophilic addition ⇒ Aldehyde > Ketone Aliphatic aldehyde > Aromatic aldehyde

#### **CRASH COURSE**

**FOR JEE ADVANCED 2020** 

FREE Online Lectures Available on You Tube

Go Premium at ₹ 1100

Doubt Support ◆ Advanced Level Test Access
 Live Test Paper Discussion ◆ Final Revision Exercises

# MOTION

9. The major product in the following reaction is:

$$\xrightarrow{\text{I -BuOH}} \xrightarrow{\text{Heat}}$$

Sol. 3

$$\frac{\text{t - B}_4\text{OH/}\Delta}{\text{(bulky base)}}$$


- 10. The incorrect statement(s) among (a) – (d) regarding acid rain is (are):
  - (a) It can corrode water pipes.
  - (b) It can damage structures made up of stone.
  - (c) It cannot cause respiratory ailments in animals
  - (d) It is not harmful for trees


  - (1) (a), (b) and (d) (2) (a), (c) and (d) (3) (c) and (d)
- (4) (c) only


Sol. 3


Acid rain can cause respiratory ailments in animals and also harmful for trees and plant.

100 mL of 0.1 M HCl is taken in a beaker and to it 100 mL of 0.1 M NaOH is added in steps of 2 mL 11. and the pH is continuously measured. Which of the following graphs correctly depicts the change in pH?









### **CRASH COURSE**

**FOR JEE ADVANCED 2020** 

FREE Online Lectures Available on You Tube


Go Premium at ₹ 1100


◆ Doubt Support ◆ Advanced Level Test Access

◆ Live Test Paper Discussion ◆ Final Revision Exercises

# **Motion**

Sol. 3





option (3)

- **12.** Consider the hypothetical situation where the azimuthal quantum number, l, takes values 0, 1, 2, ..... n + 1, where n is the principal quantum number. Then, the element with atomic number:
  - (1) 13 has a half-filled valence subshell
- (2) 9 is the first alkali metal
- (3) 8 is the first noble gas
- (4) 6 has a 2p-valence subshell

Sol. 1

1 (1) 
$$X = 1s^2 1p^6 1d^5$$
 - half filled (2)  $X = 1s^2 1p^6 1d^1$  - not alkali metal (3)  $X = 1s^2 1p^6$  - Second nobel gas Option (1)

- **13.** The d-electron configuration of  $[Ru(en)_3]Cl_2$  and  $[Fe(H_2O)_6]Cl_2$ , respectively are:
  - (1)  $t_{2g}^4 e_g^2$  and  $t_{2g}^6 e_g^0$

(2)  $t_{2g}^6 e_g^0$  and  $t_{2g}^6 e_g^0$ 

(3)  $t_{2g}^4 e_g^2$  and  $t_{2g}^4 e_g^2$ 

(4)  $t_{2g}^6 e_g^0$  and  $t_{2g}^4 e_g^2$ 

Sol. 4

$$[Ru(en)_3]Cl_2 \qquad [Fe(H_2O)_6]Cl_2$$

$$Low spin \qquad High spin$$

$$complex \qquad complex$$

$$d^6 - \Box eg^0$$

$$d^6 - \Box t_{2g}^6$$

$$d^6 - \Box t_{2g}^6$$


**FOR JEE ADVANCED 2020** 

FREE Online Lectures Available on You Tube

Doubt Support ◆ Advanced Level Test Access
 Live Test Paper Discussion ◆ Final Revision Exercises

# MOTION

14. Consider the following molecules and statements related to them:



- (a) (B) is more likely to be crystalline than (A)
- (b) (B) has higher boiling point than (A)
- (c) (B) dissolves more readily than (A) in water

Identify the correct option from below:

(1) (a) and (c) are true

(2) only (a) is true

(3) (b) and (c) are true

(4) (a) and (b) are true

Sol. Bonus

All answer are correct

**15**. The strengths of 5.6 volume hydrogen peroxide (of density 1 g/mL) in terms of mass percentage and molarity (M), respectively, are:

(Take molar mass of hydrogen peroxide as 34 g/mol)

(1) 0.85 and 0.5

(2) 0.85 and 0.25

(3) 1.7 and 0.25

(4) 1.7 and 0.5

Sol.

Volume strength = 5.6V

molarity = 
$$\frac{5.6}{11.2}$$
 = 0.5 mol / l

$$mass \% = \left[\frac{0.5 \times 34}{10}\right] \times \frac{1}{1g/ml}$$

Ans. 1.7 & 0.5 option (4)

16. The compound A in the following reactions is:

$$A = \frac{\text{(i) CH}_3\text{MgBr/H}_2\text{O}}{\text{(ii) Conc. H}_2\text{SO}_4/\Delta}$$

$$B \xrightarrow{(i) O_3} C + D$$

$$C \xrightarrow{\text{(i) Conc. KOH}} COO^{\ominus}K^{+} + COO^{\ominus}K^{+}$$

$$D \xrightarrow{Ba(OH)_2} \begin{array}{c} CH_3 & O \\ I & II \\ H_3C-C=CH-C-CH_3 \end{array}$$

$$\begin{array}{c} O \\ II \\ (1) C_6H_5-CH_2-C-CH \end{array}$$

$$\begin{array}{c}
O \\
II \\
(4) C_6H_5-C-CH_2CH
\end{array}$$

#### **CRASH COURSE**

**FOR JEE ADVANCED 2020** 

FREE Online Lectures Available on You Tube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access

◆ Live Test Paper Discussion ◆ Final Revision Exercises

Sol.

- **17.** A mixture of one mole each of H<sub>2</sub>, He and O<sub>2</sub> each are enclosed in a cylinder of volume V at temperature T. If the partial pressure of H<sub>2</sub> is 2 atm, the total pressure of the gases in the cylinder is: (2) 14 atm (3) 38 atm (4) 22 atm (1) 6 atm
- Sol.

$$p_{H_2} = 2 \text{ atm} = xH_2 \times p_{total}$$

$$2 \text{ atm} = \frac{1}{1+1+1} \times P_{\text{total}}$$

$$P_{total} = 6 atm$$

 $P_{total} = 6 atm$ Ans. option (1)

18. Three isomers A, B and C (mol. formula C<sub>8</sub>H<sub>11</sub>N) give the following results:

A and C 
$$\xrightarrow{\text{Diazotization}}$$
 P + Q  $\xrightarrow{\text{(i) Hydrolysis}}$  R (product of A) + S (product of C)  $(KMnO_4+H^+)$ 

R has lower boiling point than S

 $\xrightarrow{C_6H_5SO_2CI}$  alkali-insoluble product

A, B and C, respectively are:

$$(1) \bigcirc \begin{matrix} \mathsf{NH}_2 \\ \mathsf{CH}_2\mathsf{CH}_3 \end{matrix}, \bigcirc \begin{matrix} \mathsf{CH}_2\mathsf{NH}_2 \\ \mathsf{CH}_2\mathsf{CH}_3 \end{matrix}, \bigcirc \begin{matrix} \mathsf{CH}_2\mathsf{CH}_3 \\ \mathsf{CH}_2\mathsf{CH}_3 \end{matrix}, \bigcirc \begin{matrix} \mathsf{NH}_2 \\ \mathsf{CH}_2\mathsf{CH}_3 \end{matrix}, \bigcirc \begin{matrix} \mathsf{NH}_2 \\ \mathsf{CH}_2\mathsf{CH}_3 \end{matrix}, \bigcirc \begin{matrix} \mathsf{CH}_2\mathsf{NH}\mathsf{CH}_3 \\ \mathsf{CH}_2\mathsf{CH}_3 \end{matrix}, \bigcirc \begin{matrix} \mathsf{CH}_2\mathsf{NH}\mathsf{CH}_3 \\ \mathsf{CH}_2\mathsf{CH}_3 \end{matrix}$$

$$(3) \bigcirc \begin{matrix} \mathsf{CH}_2\mathsf{CH}_3 \\ \mathsf{NH}_2 \end{matrix}, \bigcirc \begin{matrix} \mathsf{CH}_2\mathsf{NH}\mathsf{CH}_3 \\ \mathsf{CH}_2\mathsf{CH}_3 \end{matrix}, \bigcirc \begin{matrix} \mathsf{CH}_2\mathsf{CH}_3 \\ \mathsf{CH}_2\mathsf{CH}_3 \end{matrix}$$

### **CRASH COURSE**

**FOR JEE ADVANCED 2020** 

FREE Online Lectures Available on You Tube

#### Go Premium at ₹ 1100

 ◆ Doubt Support ◆ Advanced Level Test Access ◆ Live Test Paper Discussion ◆ Final Revision Exercises

## Motion

#### Sol. 2

$$(A) \qquad (C) \qquad (D) \qquad (D)$$

**19.** For the reaction  $2A + 3B + \frac{3}{2}C \rightarrow 3P$ , which statement is correct?

(1) 
$$\frac{dn_A}{dt} = \frac{dn_B}{dt} = \frac{dn_C}{dt}$$

(2) 
$$\frac{dn_A}{dt} = \frac{3}{2} \frac{dn_B}{dt} = \frac{3}{4} \frac{dn_C}{dt}$$

(3) 
$$\frac{dn_A}{dt} = \frac{2}{3} \frac{dn_B}{dt} = \frac{4}{3} \frac{dn_C}{dt}$$

(4) 
$$\frac{dn_A}{dt} = \frac{2}{3} \frac{dn_B}{dt} = \frac{3}{4} \frac{dn_C}{dt}$$

#### Sol. 3

$$2A + 3B + \frac{3}{2}C \longrightarrow 3P$$

$$ROR = \frac{1}{2} \left\lceil \frac{-d[n_A]}{dt} \right\rceil = \frac{1}{3} \left\lceil \frac{-d[n_B]}{dt} \right\rceil = \frac{2}{3} \left\lceil \frac{-d[n_c]}{dt} \right\rceil = \frac{1}{3} \left\lceil \frac{+d[n_e]}{dt} \right\rceil$$

$$\left[\frac{-dn_{_{A}}}{dt}\right] = \frac{2}{3} \left[\frac{-d[n_{_{B}}]}{dt}\right] = \frac{4}{3} \left[\frac{-d[n_{_{e}}]}{dt}\right]$$

**20.** Consider the following reaction:

$$d \oplus O$$
  $O \oplus a$   $O \oplus$ 

The product 'P' gives positive ceric ammonium nitrate test. This is because of the presence of which of these –OH group(s)?

- (1) (b) only
- (2) (b) and (d)
- (3) (c) and (d)
- (4) (d) only

#### **CRASH COURSE**

**FOR JEE ADVANCED 2020** 

FREE Online Lectures Available on You Tube

Go Premium at ₹ 1100

Doubt Support ◆ Advanced Level Test Access
 Live Test Paper Discussion ◆ Final Revision Exercises

# **Motion**<sup>®</sup>

Sol. 1

$$(\text{OH})_{c} \\ (\text{OH})_{b} \\ \text{OH}_{a} \\ \text{OH}_{b} \\ \text{OH}_{a} \\ \text{CH=O}$$

- **21.** The volume (in mL) of 0.1 N NaOH required to neutralise 10 mL of 0.1 N phosphinic acid is \_\_\_\_\_
- Sol. 10 ml

$$\label{eq:NaOH} \begin{split} \text{NaOH} + \text{H}_3\text{PO}_2 & \longrightarrow \text{NaH}_2\text{PO}_2 + \text{H}_2\text{O} \\ & \text{Phosphinic} \\ \text{Vol.} \times 0.1 = 0.1 \times 10 \\ \text{vol} = 10 \text{ ml Ans.} \end{split}$$

**22.** An acidic solution of dichromate is electrolyzed for 8 minutes using 2A current. As per the following equation

$$Cr_2O_7^{2-} + 14H^+ + 6e^- \longrightarrow 2Cr^{3+} + 7H_2O$$

The amount of  $Cr^{3+}$  obtained was 0.104 g. The efficiency of the process (in %) is (Take: F = 96000 C, At. mass of chromium = 52) \_\_\_\_\_.

Sol. 60 %

[moles of Cr<sup>3+</sup>] × 3 = 
$$\frac{8 \times 60 \times 2}{96000}$$

moles of 
$$Cr^{3+} = \frac{8 \times 4}{9600} = \frac{1}{300}$$
 mol

mass of 
$$Cr^{3+} = \frac{52}{300} g$$

% efficiency = 
$$\frac{Actual \text{ obtained Amt}}{Theo. \text{ obtained Amt}} \times 100$$

$$= \frac{0.104}{\frac{52}{300}} \times 100$$

$$= 30 \times \frac{104}{52} = 60\%$$

- 23. If 250 cm<sup>3</sup> of an aqueous solution containing 0.73 g of a protein A is isotonic with one litre of another aqueous solution containing 1.65 g of a protein B, at 298 K, the ratio of the molecular masses of A and B is  $\_\_\_\_ \times 10^{-2}$  (to the nearest integer).
- Sol. 177

$$\frac{0.73}{M_{A}} \! \times \! \frac{1000}{250} = \frac{1.65}{M_{B}}$$

$$\frac{M_A}{M_B} = \frac{73 \times 4}{165} = 1.769$$
$$= 176.9 \times 10^{-2}$$

 $= 177 \times 10^{-2}$ 

### **CRASH COURSE**

**FOR JEE ADVANCED 2020** 

FREE Online Lectures Available on You Tube

Go Premium at ₹ 1100

Doubt Support ◆ Advanced Level Test Access
 Live Test Paper Discussion ◆ Final Revision Exercises



- **24.**  $6.023 \times 10^{22}$  molecules are present in 10 g of a substance 'x'. The molarity of a solution containing 5 g of substance 'x' in 2 L solution is \_\_\_\_\_ ×  $10^{-3}$ .
- Sol. 25

Mol. wt of 'x' = 
$$\frac{10}{6.023 \times 10^{22}} \times 6.023 \times 10^{23}$$
  
= 100 g/mol

$$M = \frac{5/100}{2} = \left(\frac{5}{200} \times 1000\right) \times 10^{-3}$$

$$M = 25 \times 10^{-3} \text{ mol/lit}$$

- **25.** The number of C=0 groups present in a tripeptide Asp–Glu–Lys is \_\_\_\_\_.
- Sol. 5

$$\begin{array}{c|c} H_2N-CH & O \\ \hline \\ CH_2 & CH_2 \\ \hline \\ C=0 & CH_2 \\ \hline \\ OH & C=0 \\ \hline \\ OH & OH \\ \end{array}$$

### Admission **OPEN**

## जब इन्होंने पूरा किया अपना सपना तो आप भी पा सकते है लक्ष्य अपना

### **JEE MAIN RESULT 2019**









### **KOTA'S PIONEER IN DIGITAL EDUCATION** 1,95,00,000+ viewers | 72,67,900+ viewing hours | 2,11,000+ Subscribers

SERVICES	SILVER	GOLD	PLATINUM
Classroom Lectures (VOD)			
Live interaction	NA		
Doubt Support	NA		
Academic & Technical Support	NA		
Complete access to all content	NA		
Classroom Study Material	NA		
Exercise Sheets	NA		
Recorded Video Solutions	NA		
Online Test Series	NA		
Revision Material	NA		
Upgrade to Regular Classroom program	Chargeable	Chargeable	Free
Physical Classroom	NA	NA	
Computer Based Test	NA	NA	
Student Performance Report	NA	NA	
Workshop & Camp	NA	NA	
Motion Solution Lab- Supervised learning and instant doubt clearance	NA	NA	
Personalised guidance and mentoring	NA	NA	

FEE STRUCTURE					
CLASS	SILVER	GOLD	PLATINUM		
7th/8th	FREE	₹ 12,000	₹ 35,000		
9th/10th	FREE	₹ 15,000	₹ 40,000		
11th	FREE	₹ 29,999	₹ 49,999		
12th	FREE	₹ 39,999	₹ 54,999		
12th Pass	FREE	₹ 39,999	₹ 59,999		
+ Student Kit will be provided at extra cost to Platinum Student.					

- SILVER (Trial) Only valid 7 DAYS or First 10 Hour's Lectures.
- GOLD (Online) can be converted to regular classroom (Any  $MOTION \ Center) \ by paying \ difference \ amount \ after \ lockdown.$
- \*\*\* PLATINUM (Online + Regular) can be converted to regular classroom (Any MOTION Center) without any cost after

New Batch Starting from:

16 & 23 September 2020

**Zero Cost EMI Available** 

