

हमारा विश्वास... हर एक विद्यार्थी है खास

**JEE  
MAIN  
Sept.  
2020**

**QUESTION PAPER WITH SOLUTION**






**CHEMISTRY \_ 5 Sep. \_ SHIFT - 2**




**MOTION™**

H.O. : 394, Rajeev Gandhi Nagar, Kota  
www.motion.ac.in | ✉: info@motion.ac.in

1. The major product formed in the following reaction is :



Sol. 1



2. Hydrogen peroxide, in the pure state, is :

(1) Linear and blue in color

(2) Linear and almost colorless

(3) Non-planar and almost colorless

(4) Planar and blue in color

Sol. 3

$\text{H}_2\text{O}_2$  has openbook structure it is non planar

3. Boron and silicon of very high purity can be obtained through :

(1) Liquefaction

(2) Electrolytic refining


(3) Zone refining

(4) Vapour phase refining

Sol. 3

Fact

4. The following molecule acts as an :



(1) Anti-histamine

(2) Antiseptic





(3) Anti-depressant

(4) Anti-bacterial

Sol. 1

Anti-histamine

5. Among the following compounds, geometrical isomerism is exhibited by :



**CRASH COURSE**  
FOR JEE ADVANCED 2020



FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100





◆ Doubt Support ◆ Advanced Level Test Access  
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: 07 Sept. 2020

Sol. 1 & 2



6. Adsorption of a gas follows Freundlich adsorption isotherm. If  $x$  is the mass of the gas adsorbed on mass  $m$  of the adsorbent, the correct plot of  $\frac{x}{m}$  versus  $p$  is :



Sol. 2

As temp. increases extent of Adsorption decreases  
Therefore correct option (2)

$$\frac{x}{m} = KP^{1/n}$$

$$\frac{x}{m} \text{ v/s } P \rightarrow \text{non linear curve}$$

**CRASH COURSE**  
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube


Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access  
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: 07 Sept. 2020

7. The compound that has the largest H-M-H bond angle (M=N, O, S, C) is :  
 (1) CH<sub>4</sub>                      (2) H<sub>2</sub>S                      (3) NH<sub>3</sub>                      (4) H<sub>2</sub>O

Sol. 1



8. The correct statement about probability density (except at infinite distance from nucleus) is :  
 (1) It can be zero for 3p orbital                      (2) It can be zero for 1s orbital  
 (3) It can never be zero for 2s orbital                      (4) It can negative for 2p orbital

Sol. 1

$$\Psi_{R/S}^2 > 0 \text{ always}$$


$$\Psi_{R/S}^2 \text{ can be } = 0; \text{ As '2s' has 1 Radial Node}$$

$$\Psi_R^2 \text{ can never be negative}$$

$$\Psi_R^2 (3P) \text{ can be } = 0 \text{ as 3P has Radial Nodes}$$

Ans. Option (1)

9. The rate constant (k) of a reaction is measured at different temperatures (T), and the data are plotted in the given figure. The activation energy of the reaction in kJ mol<sup>-1</sup> is : (R is gas constant)



- Sol. 4                      (1) R                      (2) 2/R                      (3) 1/R                      (4) 2R

$$\ln(k) = \ln(A) - \frac{E_a}{R} \left( \frac{1}{T} \right)$$

$$\ln(A) = 10$$

$$\text{Slope} = \frac{-E_a}{R} \times 10^{-3} = -10/5$$

$$E_a = 2000R \text{ J/mol}$$

$$E_a = 2R \text{ KJ/mol}$$

**CRASH COURSE**  
**FOR JEE ADVANCED 2020**


FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

◆ Doubt Support    ◆ Advanced Level Test Access  
 ◆ Live Test Paper Discussion    ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**


10. The variation of molar conductivity with concentration of an electrolyte (X) in aqueous solution is shown in the given figure.



The electrolyte X is :






- (1) HCl                      (2) CH<sub>3</sub>COOH                      (3) NaCl                      (4) KNO<sub>3</sub>

Sol. 2



Such type of variation is always for weak electrolyte  
Hence Ans (2) CH<sub>3</sub>COOH

11. The final major product of the following reaction is :



**CRASH COURSE**  
FOR JEE ADVANCED 2020


FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100






◆ Doubt Support ◆ Advanced Level Test Access  
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: 07 Sept. 2020


Sol. 3



12. The major product of the following reaction is :



Sol. 3



13. Lattice enthalpy and enthalpy of solution of NaCl are  $788 \text{ kJ mol}^{-1}$ , and  $4 \text{ kJ mol}^{-1}$ , respectively. The hydration enthalpy of NaCl is :

(1)  $-780 \text{ kJ mol}^{-1}$

(2)  $784 \text{ kJ mol}^{-1}$

(3)  $-784 \text{ kJ mol}^{-1}$

(4)  $780 \text{ kJ mol}^{-1}$

Sol. 3

$$\Delta H_{\text{sol}} = \text{L.E.} + \Delta H_{\text{hyd}}$$

$$4 = 788 + \Delta H_{\text{hyd}}$$

$$\Delta H_{\text{hyd}} = -784 \text{ KJ/mol Ans}$$

14. Reaction of ammonia with excess  $\text{Cl}_2$  gives :

(1)  $\text{NH}_4\text{Cl}$  and  $\text{N}_2$

(2)  $\text{NH}_4\text{Cl}$  and  $\text{HCl}$

(3)  $\text{NCl}_3$  and  $\text{HCl}$

(4)  $\text{NCl}_3$  and  $\text{NH}_4\text{Cl}$

**CRASH COURSE**  
FOR JEE ADVANCED 2020



FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access  
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: 07 Sept. 2020


**Sol. 3**



**15.** Which one of the following polymers is not obtained by condensation polymerisation ?

- (1) Bakelite (2) Nylon 6  
(3) Buna-N (4) Nylon 6, 6

**Sol. 2**



**16.** Consider the complex ions, trans-[Co(en)<sub>2</sub>Cl<sub>2</sub>]<sup>+</sup> (A) and cis-[Co(en)<sub>2</sub>Cl<sub>2</sub>]<sup>+</sup> (B)

The correct statement regarding them is :

- (1) Both (A) and (B) can be optically active.  
(2) (A) can be optically active, but (B) cannot be optically active.  
(3) Both (A) and (B) cannot be optically active.  
(4) (A) cannot be optically active, but (B) can be optically active.

**Sol. 4**

Due to presence of Pos (A) cannot be optically active, but (B) can be optically active

**17.** An element crystallises in a face-centred cubic (fcc) unit cell with cell edge a. The distance between the centres of two nearest octahedral voids in the crystal lattice is :

- (1) a                      (2)  $\frac{a}{2}$                       (3)  $\sqrt{2}a$                       (4)  $\frac{a}{\sqrt{2}}$

**Sol. 4**

Nearest octahedral voids

One along edge center & other at Body centre

$$\text{Distance} = \sqrt{\left(\frac{a}{2}\right)^2 + \left(\frac{a}{2}\right)^2} = \sqrt{2} \frac{a}{2}$$

$$= \frac{a}{\sqrt{2}} \text{ Ans.}$$

**18.** The correct order of the ionic radii of O<sup>2-</sup>, N<sup>3-</sup>, F<sup>-</sup>, Mg<sup>2+</sup>, Na<sup>+</sup> and Al<sup>3+</sup> is :

- (1) N<sup>3-</sup> < O<sup>2-</sup> < F<sup>-</sup> < Na<sup>+</sup> < Mg<sup>2+</sup> < Al<sup>3+</sup>                      (2) N<sup>3-</sup> < F<sup>-</sup> < O<sup>2-</sup> < Mg<sup>2+</sup> < Na<sup>+</sup> < Al<sup>3+</sup>  
(3) Al<sup>3+</sup> < Na<sup>+</sup> < Mg<sup>2+</sup> < O<sup>2-</sup> < F<sup>-</sup> < N<sup>3-</sup>                      (4) Al<sup>3+</sup> < Mg<sup>2+</sup> < Na<sup>+</sup> < F<sup>-</sup> < O<sup>2-</sup> < N<sup>3-</sup>

**Sol. 4**

all are Isoelectronic

$$(1) \frac{\text{N}^{3-} \text{O}^{2-} \text{F}^- \text{Na}^+ \text{Mg}^{2+} \text{Al}^{3+}}{\text{Z} \uparrow, \text{Z}_{\text{eff}} \uparrow, \text{Ionic Radii} \downarrow}$$

$$(2) \text{Al}^{3+} < \text{Mg}^{2+} < \text{Na}^+ < \text{F}^- < \text{O}^{2-} < \text{N}^{3-}$$

**CRASH COURSE**  
**FOR JEE ADVANCED 2020**


FREE Online Lectures Available on **YouTube**

Go Premium at ₹ 1100

- ◆ Doubt Support ◆ Advanced Level Test Access  
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

19. The increasing order of boiling points of the following compounds is :




(1) I < III < IV < II

(2) IV < I < II < III

(3) I < IV < III < II

(4) III < I < II < IV

Sol. 3



20. The one that is NOT suitable for the removal of permanent hardness of water is :

(1) Ion-exchange method



(2) Calgon's method

(3) Treatment with sodium carbonate

(4) Clark's method

Sol. 4





Clark's method is used for Removal of Temporary hardness



21. For a reaction  $\text{X} + \text{Y} \rightleftharpoons 2\text{Z}$ , 1.0 mol of X, 1.5 mol of Y and 0.5 mol of Z were taken in a 1 L vessel and allowed to react. At equilibrium, the concentration of Z was  $1.0 \text{ mol L}^{-1}$ . The equilibrium

constant of reaction is  $\frac{x}{15}$ . The value of x is \_\_\_\_\_.

Sol. 16



$$K_{\text{eq}} = \frac{(1)^2}{\frac{5}{4} \times \frac{3}{4}} = \frac{16}{15}$$

x = 16 Ans.

**CRASH COURSE**  
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100


◆ Doubt Support ◆ Advanced Level Test Access  
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: 07 Sept. 2020



**22.** The volume, in mL, of 0.02 M  $K_2Cr_2O_7$  solution required to react with 0.288 g of ferrous oxalate in acidic medium is \_\_\_\_\_.  
(Molar mass of Fe = 56 g mol<sup>-1</sup>)

**Sol.** **50 ml**




$$\frac{0.02 \times \text{vol} \times 6}{1000} = 3 \times \frac{0.288}{144} \times 100$$

$$\text{Vol.} = \frac{200}{4} = 50 \text{ ml Ans.}$$


**23.** Considering that  $\Delta_0 > P$ , the magnetic moment (in BM) of  $[Ru(H_2O)_6]^{2+}$  would be \_\_\_\_\_.

**Sol.** **0**



**24.** For a dimerization reaction,  $2A(g) \rightarrow A_2(g)$  at 298 K,  $\Delta U^\ominus = -20 \text{ kJ mol}^{-1}$ ,  $\Delta S^\ominus = -30 \text{ kJ mol}^{-1}$ , then the  $\Delta G^\ominus$  will be \_\_\_\_\_ J.

**Sol.** **-13538 J**



$$\Delta U^\ominus = -20 \text{ kJ}$$

$$\Delta H^\ominus = -20000 + (-1)R \times 298$$

$$\Delta G^\ominus = -20000 - 298R + 30 \times 298$$


$$\Delta G^\ominus = -20,000 + 298 \left( \frac{90 - 25}{3} \right)$$

$$\Delta G^\ominus = 20,000 + \frac{298 \times 65}{3}$$

$$\Delta G^\ominus = -13538 \text{ J}$$

**25.** The number of chiral carbons present in sucrose is \_\_\_\_\_.

**Sol.** **9**



**CRASH COURSE**  
**FOR JEE ADVANCED 2020**

FREE Online Lectures Available on **YouTube**

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access  
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

जब इन्होंने पूरा किया अपना सपना  
तो आप भी पा सकते है लक्ष्य अपना

Admission  
**OPEN**

## JEE MAIN RESULT 2019



**Nitin Gupta**

Marks  
**335**  
13th (2019)

Marks  
**149**  
12th (2018)



**Shiv Modi**

Marks  
**318**  
13th (2019)

Marks  
**153**  
12th (2018)



**Ritik Bansal**

Marks  
**308**  
13th (2019)

Marks  
**218**  
12th (2018)



**Shubham Kumar**

Marks  
**300**  
13th (2019)

Marks  
**153**  
12th (2018)

### KOTA'S PIONEER IN DIGITAL EDUCATION

**1,95,00,000+** viewers | **72,67,900+** viewing hours | **2,11,000+** Subscribers

SERVICES	SILVER	GOLD	PLATINUM
Classroom Lectures (VOD)			
Live interaction	NA		
Doubt Support	NA		
Academic & Technical Support	NA		
Complete access to all content	NA		
Classroom Study Material	NA		
Exercise Sheets	NA		
Recorded Video Solutions	NA		
Online Test Series	NA		
Revision Material	NA		
<b>Upgrade to Regular Classroom program</b>	<b>Chargeable</b>	<b>Chargeable</b>	<b>Free</b>
Physical Classroom	NA	NA	
Computer Based Test	NA	NA	
Student Performance Report	NA	NA	
Workshop & Camp	NA	NA	
Motion Solution Lab- Supervised learning and instant doubt clearance	NA	NA	
Personalised guidance and mentoring	NA	NA	

### FEE STRUCTURE

CLASS	SILVER	GOLD	PLATINUM
7th/8th	FREE	₹ 12,000	₹ 35,000
9th/10th	FREE	₹ 15,000	₹ 40,000
11th	FREE	₹ 29,999	₹ 49,999
12th	FREE	₹ 39,999	₹ 54,999
12th Pass	FREE	₹ 39,999	₹ 59,999

+ Student Kit will be provided at extra cost to Platinum Student.

- \* **SILVER (Trial)** Only valid 7 DAYS or First 10 Hour's Lectures.
- \*\* **GOLD (Online)** can be converted to regular classroom (Any MOTION Center) by paying difference amount after lockdown.
- \*\*\* **PLATINUM (Online + Regular)** can be converted to regular classroom (Any MOTION Center) without any cost after lockdown.

New Batch Starting from :  
**16 & 23 September 2020**

**Zero Cost EMI Available**

**MOTION™**

H.O. : 394, Rajeev Gandhi Nagar, Kota  
www.motion.ac.in | ✉ : info@motion.ac.in