

हमारा विश्वास... हर एक विद्यार्थी है खास

**JEE
MAIN
Sept.
2020**

QUESTION PAPER WITH SOLUTION

CHEMISTRY _ 6 Sep. _ SHIFT - 2



MOTION™

H.O. : 394, Rajeev Gandhi Nagar, Kota
www.motion.ac.in | ✉: info@motion.ac.in

1. Match the following :

Test/Method

- (i) Lucas Test
(ii) Dumas method
(iii) kjeldjhl's method
(iv) Hinsberg Test

Reagent


- (a) $C_6H_5SO_2Cl/aq. KOH$
(b) $HNO_3/AgNO_3$
(c) CuO/CO_2
(d) Conc. HCl and $ZnCl_2$
(e) H_2SO_4

- (1) (i)-(d), (ii)-(c), (iii)-(e), (iv)-(a) (2) (i)-(b), (ii)-(a), (iii)-(c), (iv)-(d)
(3) (i)-(b), (ii)-(d), (iii)-(e), (iv)-(a) (4) (i)-(d), (ii)-(c), (iii)-(b)-(iv)-(e)

Sol. 1

By Theory


2. The IUPAC name of the following compound is:



- (1) 2-nitro-4-hydroxymethyl-5-amino benzaldehyde
(2) 3-amino-4-hydroxymethyl-5-nitro benzaldehyde
(3) 4-amino-2-formyl-5-hydroxymehtyl nitrobenzene
(4) 5-amino-4-hydroxymethyl-2-nitro benzaldehyde

Sol. 4

3. For the Given Cell;



Change in Gibbs energy (ΔG) is negative, if :

- (1) $C_2 = \sqrt{2}C_1$ (2) $C_2 = \frac{C_1}{\sqrt{2}}$
(3) $C_1 = 2C_2$ (4) $C_1 = C_2$

Sol. 1

$$E = 0 - \frac{0.059}{2} \log \left[\frac{C_1}{C_2} \right]$$

$$\Delta G = -nFE$$

$$= +nF \times \frac{RT}{nF} \times 2.303 \log \left[\frac{C_1}{C_2} \right]$$

$$\log \left[\frac{C_1}{C_2} \right] < 0$$

$$C_1 < C_2$$

$$C_2 = \sqrt{2}C_1$$

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100


- ◆ Doubt Support
- ◆ Advanced Level Test Access
- ◆ Live Test Paper Discussion
- ◆ Final Revision Exercises

Start Date: 07 Sept. 2020

4. Reaction of an inorganic sulphite X with dilute H_2SO_4 generates compound Y. Reaction of Y with NaOH gives X. Further, the reaction of X with Y and water affords compound Z. Y and Z, respectively, are :


- (1) SO_2 and $NaHSO_3$ (2) S and Na_2SO_3
 (3) SO_2 and Na_2SO_3 (4) SO_3 and $NaHSO_3$

Sol. 1



Ans. (1) $Y = SO_2$
 $Z = NaHSO_3$

5. The value of K_c is 64 at 800 K for the reaction $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$
 The value of K_c for the following reaction is :



- (1) $1/4$ (2) 8 (3) $1/8$ (4) $1/64$

Sol. 3

$$K_c = (64)^{-1/2}$$

$$K_c = \frac{1}{8}$$

6. The correct match between Item - I (Starting material) and Item - II (reagent) for the preparation of benzaldehyde is :

Item - I	Item - II
(I) Benzene	(P) HCl and $SnCl_2 \cdot H_3O^+$
(II) Benzonitrile	(Q) H_2 , Pd- $BaSO_4$, and quinoline
(III) Benzoyl Chloride	(R) CO, HCl and $AlCl_3$

- (1) (I) - (R), (II) - (P) and (III) - (Q)
 (2) (I) - (P), (II) - (Q) and (III) - (R)
 (3) (I) - (Q), (II) - (R) and (III) - (P)
 (4) (I) - (R), (II) - (Q) and (III) - (P)

CRASH COURSE
FOR JEE ADVANCED 2020




FREE Online Lectures Available on  YouTube

Go Premium at ₹ 1100

- ◆ Doubt Support ◆ Advanced Level Test Access
- ◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**


Sol. 1



7. For a d^4 metal ion in an octahedral field, the correct electronic configuration is :

- (1) $e_g^2 t_{2g}^2$ when $\Delta_0 < P$ (2) $t_{2g}^4 e_g^0$ when $\Delta_0 < P$
 (3) $t_{2g}^3 e_g^1$ when $\Delta_0 > P$ (4) $t_{2g}^3 e_g^1$ when $\Delta_0 < P$

Sol. 4



8. The correct match between Item - I and Item - II is :

- | Item - I | Item - II |
|--|------------------------------------|
| (a) Natural rubber | (I) 1,3-butadiene + styrene |
| (b) Neoprene | (II) 1,3-butadiene + acrylonitrile |
| (c) Buna-N | (III) Chloroprene |
| (d) Buna-S | (IV) Isoprene |
| (1) (a) - (III), (b) - (IV), (c) - (I), (d) - (II) | |
| (2) (a) - (IV), (b) - (III), (c) - (II), (d) - (I) | |
| (3) (a) - (IV), (b) - (III), (c) - (I), (d) - (II) | |
| (4) (a) - (III), (b) - (IV), (c) - (II), (d) - (I) | |


Sol. 2

By Theory

9. Which one of the following statement is not true ?
 (1) Lactose contains α -glycoside linkage between C_1 of galactose and C_4 of glucose.
 (2) Lactose is a reducing sugar and it gives Fehling's test.
 (3) On acid hydrolysis, lactose gives one molecule of D(+)-glucose and one molecule of D(+)-galactose.
 (4) Lactose($C_{11}H_{22}O_{11}$) is a disaccharide and it contains 8 hydroxyl groups.

Sol. 1

Lactose contains β -glycosidic linkage between C_1 of galactose and C_4 of glucose.



10. The element that can be refined by distillation is :
 (1) tin (2) gallium
 (3) zinc (4) nickel

Sol. 3

Zinc \rightarrow Refined by distillation

11. Match the following compounds (Column-I) with their uses (Column-II) :

S. No.	Column - I	S. No.	Column - II
(I)	$Ca(OH)_2$	(A)	Casts of statues
(II)	NaCl	(B)	White wash
(III)	$CaSO_4 \cdot \frac{1}{2} H_2O$	(C)	Antacid
(IV)	$CaCO_3$	(D)	Washing soda preparation
(1)	(I)-(B), (II)-(C), (III)-(D), (IV)-(A)	(2)	(I)-(C), (II)-(D), (III)-(B), (IV)-(A)
(3)	(I)-(B), (II)-(D), (III)-(A), (IV)-(C)	(4)	(I)-(D), (II)-(A), (III)-(C), (IV)-(B)

Sol. 3

- (i) $Ca(OH)_2 \rightarrow$ used in white wash due to its disinfectant nature
 (ii) $CaCO_3 \rightarrow$ it used as an Antacid

(iii) $CaSO_4 \cdot \frac{1}{2} H_2O \rightarrow$ Formaking casts of statues and busts

(iii) Preparation of wasing soda (sodium carbonate)

- (1) $2NH_3 + H_2O + CO_2 \longrightarrow (NH_4)_2CO_3$
 (2) $(NH_4)_2CO_3 + H_2O + CO_2 \longrightarrow 2NH_4HCO_3$
 (3) $NH_4HCO_3 + NaCl \longrightarrow NH_4Cl + NaHCO_3$
 (4) $2NaHCO_3 \longrightarrow Na_2CO_3 + CO_2 + H_2O$

Ans. (3)


12. Mischmetal is an alloy consisting mainly of :
 (1) lanthanoid and actinoid metals (2) lanthanoid metals
 (3) actinoid metals (4) actinoid and transition metals

Sol. 2
 Misch metal - well known alloy is mischmetal which consists of Lanthanoid metal (~ 95%) and iron (~ 5%) and Traces of S, C, Ca and Al

13. For a reaction,
 $4M(s) + nO_2(g) \rightarrow 2M_2O_n(s)$
 the free energy change is plotted as a function of temperature. The temperature below which the oxide is stable could be inferred from the plot as the point at which :
 (1) the free energy change shows a change from negative to positive value.
 (2) the slope changes from positive to zero
 (3) the slope changes from positive to negative.
 (4) the slope changes from negative to positive.




Sol. 1
 $\Delta G = \Delta H - T\Delta S$
 $\Delta G = -ve$ (stable oxide)
 $\Delta G = +ve$ (unstable oxide)

14. The increasing order of the boiling points of the major products A, B and C of the following reaction will be :



- (1) $A < B < C$ (2) $C < A < B$
 (3) $A < C < B$ (4) $B < C < A$

Sol. 4



$$B.P \propto \frac{1}{\text{Branching}}$$

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access
 ◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

15. The average molar mass of chlorine is 35.5 g mol^{-1} . The ratio of ^{35}Cl to ^{37}Cl in naturally occurring chlorine is close to :

- (1) 1 : 1 (2) 3 : 1
(3) 2 : 1 (4) 4 : 1





Sol. 2

$$\frac{35x + 37y}{x + y} = 35.5$$


$$1.5y = -0.5x$$

$$x/y = 3/1$$

16. which of the following compound can be prepared in good yield by Gabriel phthalimide synthesis ?



Sol. 1



In this reaction, the alkyl halide should be $\text{CH}_2 - \text{Cl}$, which can give $\text{S}_{\text{N}}2$ reaction easily.

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube


Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

17. The reaction of NO with N_2O_4 at 250 K gives:
 (1) N_2O (2) NO_2
 (3) N_2O_5 (4) N_2O_3

Sol. 4



18. A set of solution is prepared using 180g of water as a solvent and 10g of different non-volatile solutes A, B and C. The relative lowering of vapour pressure in the presence of these solutes are in the order [Given, molar mass of A = $100g\ mol^{-1}$ B = $200g\ mol^{-1}$; C = $10,000g\ mol^{-1}$]
 (1) $A > C > B$ (2) $B > C > A$
 (3) $C > B > A$ (4) $A > B > C$

Sol. 4

$$RLVP_A = \frac{0.1}{10.1} = \frac{1}{101}$$

$$RLVP_B = \frac{0.05}{10.05} = \frac{1}{201}$$

$$RLVP_C = \frac{10^{-3}}{10} = 10^{-4}$$

$$A > B > C$$



19. Dihydrogen of high purity (> 99.95%) is obtained through :
 (1) the electrolysis of acidified water using Pt electrodes.
 (2) the reaction of Zn with dilute HCl
 (3) the electrolysis of brine solution.
 (4) the electrolysis of warm $Ba(OH)_2$ solution using Ni electrodes.

Sol. 4

High purity (>99.95%) dihydrogen is obtained by electrolysing warm aqueous barium hydroxide solution between nickel electrodes

20. A crystal is made up of metal iron ' M_1 ' and ' M_2 ' and oxide ions. Oxide ions form a ccp lattice structure. The cation ' M_1 ' occupies 50% of octahedral voids and the cation ' M_2 ' occupies 12.5% of tetrahedral voids of oxide lattice. The oxidation number of ' M_1 ' and ' M_2 ' are, respectively :
 (1) +2, +4 (2) +3, +1
 (3) +4, +2 (4) +1, +3

Sol. 1



$$+2x + y = 8$$

$$x = +2$$

$$y = +4$$

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access
 ◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

21. For Freundlich adsorption isotherm, a plot of $\log(x/m)$ (y-axis) and $\log p$ (x-axis) gives a straight line. The intercept and slope for the line is 0.4771 and 2, respectively. The mass of gas, adsorbed per gram of adsorbent if the initial pressure is 0.04 atm, is $\times 10^{-4}g$. ($\log 3=0.4771$)

Sol. 48

$$x/m = KP^{1/n}$$

$$\log(x/m) = \log(k) + \frac{1}{n} \log(P)$$

$$K = 3, \frac{1}{n} = 2 \Rightarrow n = \frac{1}{2}$$

$$x/m = 3(P)^{1/n}$$

$$\begin{aligned} x/m &= 3 \times (4 \times 10^{-2})^2 \\ &= 48 \times 10^{-4}g \end{aligned}$$


22. The atomic number of Unnilunium is

Sol. 101

Fact

23. A solution of phenol in chloroform when treated with aqueous NaOH gives compound P as a major product. The mass percentage of carbon in P is (to the nearest integer)
(Atomic mass : C = 12; H = 1; O = 16)

Sol. 68.85%



$$\text{mass of Salicylaldehyde} = 12 \times 7 + 6 \times 1 + 16 \times 2 = 122$$

$$\text{mass of carbon} = 12 \times 7 = 84$$

$$\text{The mass \% of carbon} = \frac{84}{122} \times 100 = 68.85\%$$

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

24. The rate of a reaction decreased by 3.555 times when the temperature was changed from 40°C to 30°C. The activation energy (in KJ mol⁻¹) of the reaction is..... [Take; R = 8.314 J mol⁻¹ K⁻¹ In 3.555 = 1.268]

Sol. $K_{40^{\circ}\text{C}} = K$; $K_{30^{\circ}\text{C}} = \frac{K}{3.555}$

$$\ln \{3.555\} = \frac{E_a}{R} \left\{ \frac{1}{303} - \frac{1}{313} \right\}$$

$$E_a = \frac{1.268 \times 8.314 \times 313 \times 303}{10}$$


$$E_a = 99980.7 \text{ J/mol.}$$

$$E_a = 99.98 \text{ kJ/mol.}$$

$$E_a = 100 \text{ kJ/mol.}$$

25. If the solubility product of AB₂ is 3.20 × 10⁻¹¹ M³, then the solubility of AB₂ in pure water is × 10⁻⁴ mol L⁻¹ [Assuming that neither kind of ion reacts with water].

Sol. 2



$$K_{\text{sp}} = 4s^3 = 32 \times 10^{-12}$$

$$S = 2 \times 10^{-4} \text{ mol/lit}$$

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

जब इन्होंने पूरा किया अपना सपना
तो आप भी पा सकते है लक्ष्य अपना

Admission
OPEN

JEE MAIN RESULT 2019



Nitin Gupta

Marks
335
13th (2019)

Marks
149
12th (2018)



Shiv Modi

Marks
318
13th (2019)

Marks
153
12th (2018)



Ritik Bansal

Marks
308
13th (2019)

Marks
218
12th (2018)



Shubham Kumar

Marks
300
13th (2019)

Marks
153
12th (2018)

KOTA'S PIONEER IN DIGITAL EDUCATION

1,95,00,000+ viewers | **72,67,900+** viewing hours | **2,11,000+** Subscribers

SERVICES	SILVER	GOLD	PLATINUM
Classroom Lectures (VOD)			
Live interaction	NA		
Doubt Support	NA		
Academic & Technical Support	NA		
Complete access to all content	NA		
Classroom Study Material	NA		
Exercise Sheets	NA		
Recorded Video Solutions	NA		
Online Test Series	NA		
Revision Material	NA		
Upgrade to Regular Classroom program	Chargeable	Chargeable	Free
Physical Classroom	NA	NA	
Computer Based Test	NA	NA	
Student Performance Report	NA	NA	
Workshop & Camp	NA	NA	
Motion Solution Lab- Supervised learning and instant doubt clearance	NA	NA	
Personalised guidance and mentoring	NA	NA	

FEE STRUCTURE

CLASS	SILVER	GOLD	PLATINUM
7th/8th	FREE	₹ 12,000	₹ 35,000
9th/10th	FREE	₹ 15,000	₹ 40,000
11th	FREE	₹ 29,999	₹ 49,999
12th	FREE	₹ 39,999	₹ 54,999
12th Pass	FREE	₹ 39,999	₹ 59,999

+ Student Kit will be provided at extra cost to Platinum Student.

- * **SILVER (Trial)** Only valid 7 DAYS or First 10 Hour's Lectures.
- ** **GOLD (Online)** can be converted to regular classroom (Any MOTION Center) by paying difference amount after lockdown.
- *** **PLATINUM (Online + Regular)** can be converted to regular classroom (Any MOTION Center) without any cost after lockdown.

New Batch Starting from :
16 & 23 September 2020

Zero Cost EMI Available

MOTION™

H.O. : 394, Rajeev Gandhi Nagar, Kota
www.motion.ac.in | ✉ : info@motion.ac.in