

CHEMISTRY _ 6 Sep. _ SHIFT - 1











H.O.: 394, Rajeev Gandhi Nagar, Kota www.motion.ac.in | \simes: info@motion.ac.in



Motion[®]

- **1.** The INCORRECT statement is :
 - (1) Cast iron is used to manufacture wrought iron.
 - (2) Brass is an alloy of copper and nickel.
 - (3) German silver is an alloy of zinc, copper and nickel.
 - (4) Bronze is an alloy of copper and tin
- Sol. 2

Brass - (copper Zinc)

Bronze - (copper tin)

- **2.** The species that has a spin-only magnetic moment of 5.9 BM, is: $(T_d = tetrahedral)$
 - (1) [Ni(CN)₄]²⁻ (square planar)
- (2) $Ni(CO)_4(T_d)$

(3) $[MnBr_4]^{2-}(T_d)$

(4) $[NiCl_{4}]^{2}$ - (T_{d})

Sol. 3

 $[MnBr_4]^{2-}$

$$\mu = \sqrt{5(5+2)} = 5.9 \text{ BM}$$

3. For the reaction

$$Fe_2N(s) + \frac{3}{2}H_2(g) \rightleftharpoons 2Fe(s) + NH_3(g)$$

(1)
$$K_c = K_n (RT)^{1/2}$$

(2)
$$K_c = K_p (RT)^{-1/2}$$

(3)
$$K_c = K_p(RT)^{\frac{3}{2}}$$

$$(4) K_c = K_p(RT)$$

Sol. 1

$$Fe_2N(s) + \frac{3}{2}H_2(g) \rightleftharpoons 2Fe(s) + NH_3(g)$$

$$\Delta n_g = 1 - \frac{3}{2} = \frac{-1}{2}$$

$$\frac{K_p}{K_c} = (RT)^{\Delta n_g} = (RT)^{-\frac{1}{2}}$$

$$K_c = \frac{K_p}{(RT)^{-\frac{1}{2}}} = K_p.(RT)^{\frac{1}{2}}$$

CRASH COURSE

FOR JEE ADVANCED 2020

FREE Online Lectures Available on You Tube

Go Premium at ₹ 1100

Doubt Support ◆ Advanced Level Test Access
 Live Test Paper Discussion ◆ Final Revision Exercises

MOTION

4. Consider the following reactions:

$$(C_{7}H_{14}) \xrightarrow{\text{ozonolysis}} 'B' + 'C'$$

$$(B' \xrightarrow{\Delta} Yellow ppt \xrightarrow{\Delta} Silver mirror$$

'A' is:

gives white

Sol.

Ozonolysis
$$CH_3-CHO + (C)$$

(A)

(B)

$$CH_3-CHO + (C)$$

$$CI$$

$$ZnCl_2/HCI$$

$$CI$$

$$Cl$$

$$Cturbidity in 5 min)$$

CRASH COURSE

FOR JEE ADVANCED 2020

FREE Online Lectures Available on You Tube

Go Premium at ₹ 1100

 ◆ Doubt Support ◆ Advanced Level Test Access ◆ Live Test Paper Discussion ◆ Final Revision Exercises

Motion

5. Arrange the following solutions in the decreasing order of pOH:

(A) 0.01 M HCI

(B) 0.01 M NaOH

(C) 0.01 M CH₃COONa

(D) 0.01 M NaCl

(1)(A) > (C) > (D) > (B)

(2)(B) > (D) > (C) > (A)

Sol.

- (i) 10^{-2} M HCl \Rightarrow [H⁺] = 10^{-2} M \rightarrow pH = 2
- (ii) 10^{-2} M NaOH \Rightarrow [OH $^{-}$] = 10^{-2} M \rightarrow pOH = 2
- (iii) $10^{-2} \text{ M CH}_3\text{COO-Na}^+ \Rightarrow [\text{OH}^+] > 10^{-7} \Rightarrow \text{pOH} < 7$
- (iv) 10^{-2} M NaČl \Rightarrow Neutral pOH = 7
- (i) > (iv) > (iii) > (ii)

6. The variation of equilibrium constant with temperature is given below:

Temperature

Equilibrium Constant

$$T_1 = 25^{\circ}C$$

 $T_2 = 100^{\circ}C$

$$K_1 = 10$$

 $K_2 = 100$

The value of ΔH^0 , ΔG^0 at T_1 and ΔG^0 at T_2 (in Kj mol⁻¹) respectively, are close to [use $R = 8.314JK^{-1} \text{ mol}^{-1}$]

- (1) 28.4, -7.14 and -5.71
- (2) 0.64, 7.14 and -5.71
- (3) 28.4, 5.71 and -14.29
- (4) 0.64, 5.71 and -14.29

Sol.

$$\text{In} \left\lceil \frac{k_2}{k_1} \right\rceil = \frac{\Delta H^{\circ}}{R} \left\{ \frac{1}{T_1} - \frac{1}{T_2} \right\}$$

$$In(10) = \frac{\Delta H^{\circ}}{R} \left\{ \frac{1}{298} - \frac{1}{373} \right\}$$

$$\frac{373 \times 298 \times 8.314 \times 2.303}{75} = \Delta H^{\circ} = 28.37 \text{ kJ mol}^{-1}$$

$$\Delta G^{\circ}_{T_1} = -RT_1 ln(K_1) = -298R ln(10) = -5.71 kJ mol^{-1}$$


$$\Delta G^{\circ}_{T_2} = -RT_2 \ln(K_2) = -373R \ln(100)$$

= -14.283 kJ/mol

7. Consider the following reactions

$$A \rightarrow P1$$
; $B \rightarrow P2$; $C \rightarrow P3$; $D \rightarrow P4$,

The order of the above reactions are a,b,c and d, respectively. The following graph is obtained when log[rate] vs. log[conc.] are plotted:



CRASH COURSE

FOR JEE ADVANCED 2020

FREE Online Lectures Available on You Tube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access

◆ Live Test Paper Discussion ◆ Final Revision Exercises

Motion

Among the following the correct sequence for the order of the reactions is :

(1) c > a > b > d

(2) d > a > b > c

(3) d > b > a > c

(4) a > b > c > d

Sol. 3

 $A \rightarrow P1$

 $B \rightarrow P2$

 $\mathsf{C}\to\mathsf{P3}$

 $D \rightarrow P4$

Rate = K (conc.) order

log(rate) = log(K) + order log (case)y c + m.x

Straight line

Slope = order

According graph

d > b > a > c order of slope

8. The major product obtained from the following reactions is :

$$O_2N - \left\langle \begin{array}{c} O \end{array} \right\rangle - C \equiv C - \left\langle \begin{array}{c} O \end{array} \right\rangle - OCH_3 \xrightarrow{\quad Hg^{2^+}/H^+}$$

Sol. 3

$$O_2N$$
 $C=C$ OCH_3 Hg^{2+}/H^+

$$O_2N \longrightarrow CH_2 - C \longrightarrow OCH_3$$

CRASH COURSE

FOR JEE ADVANCED 2020

FREE Online Lectures Available on You Tube

Go Premium at ₹ 1100

Doubt Support ◆ Advanced Level Test Access
 Live Test Paper Discussion ◆ Final Revision Exercises

Motion^m

- **9.** Which of the following compounds shows geometrical isomerism?
 - (1) 2-methylpent-1-ene

(2) 4-methylpent-2-ene

(3) 2-methylpent-2-ene

(4) 4-methylpent-1-ene

Sol. 2

10. The lanthanoid that does NOT shows +4 oxidation state is :

(1) Dy

(2) Ce

(3) Tb

(4) Eu

Sol. 4

Fact

11. The major products of the following reactions are :

$$\begin{array}{c} \text{CH}_3 \\ \text{I} \\ \text{CH}_3\text{- CH - CH - CH}_3 \\ \text{OSO}_2\text{CH}_3 \end{array} \xrightarrow{\begin{array}{c} \text{(i) KOt}_{\text{Bu}} / \Delta \\ \text{(ii) O}_3 / \text{H}_2\text{O}_2 \end{array}} \rightarrow \end{array}$$

Sol. 1

$$CH_{3} \xrightarrow{CH} CH - CH - CH_{3} \xrightarrow{KOt_{Bu}} CH_{3} - CH - CH = CH_{2}$$

$$OSO_{2}CH_{3} \xrightarrow{CH_{3}} CH_{3}$$

$$CH_{3} \xrightarrow{CH_{3}-CH_{2}COOH} + HCOOH$$

$$CH_{3} \xrightarrow{CH_{3}-CH_{3}-CH_{3}}$$

CRASH COURSE

FOR JEE ADVANCED 2020

FREE Online Lectures Available on You Tube

Go Premium at ₹ 1100

Doubt Support ◆ Advanced Level Test Access
 Live Test Paper Discussion ◆ Final Revision Exercises

12. The major product of the following reaction is :

$$CH_3$$
 $2HBr \rightarrow$
 NO_2

(1)
$$Br \longrightarrow NO_2$$
 Br

$$(4) \qquad \begin{array}{c} H_3C & Br \\ \\ NO_2 & Br \end{array}$$

Sol. 2

$$CH_3$$
 $2HBr$
 Br
 NO_2
 Br
 NO_2

13. The increasing order of pK_h values of the following compounds is :

$$N(CH_3)_2$$
 $N(CH_3)_2$ $NHCH_3$ $NHCH_3$ OCH_3 OC

- (1) I < II < III < IV
- (3) I < II < IV < III

- (2) II < IV < III < I
- (4) II < I < III < IV

FOR JEE ADVANCED 2020

FREE Online Lectures Available on You Tube

Go Premium at ₹ 1100

Doubt Support ◆ Advanced Level Test Access
 Live Test Paper Discussion ◆ Final Revision Exercises

Motion[®]

Sol. 3

Order of pK_b

$$N(CH_3)_2$$
 $N(CH_3)_2$ $NHCH_3$ $NHCH_3$ OCH_3 OCH_3

- **14.** kraft temperature is the temperature :
 - (1) Above which the aqueous solution of detergents starts boiling
 - (2) Below which the formation of micelles takes place.
 - (3) Above which the formation of micelles takes place.
 - (4) Below which the aqueous solution of detergents starts freezing.

Sol. 3

 T_{κ} + temp. above which formation of micelles takes place.

15. The set that contains atomic numbers of only transition elements, is ?

(1) 9, 17, 34, 38

(2) 21, 25, 42, 72

(3) 37, 42, 50, 64

(4) 21, 32, 53, 64

Sol. 2

Tranition elements = 21 to 30 37 to 48

57 & 72 to 80

Ans. 21, 25, 42 & 72

16. Consider the Assertion and Reason given below.

Assertion (A): Ethene polymerized in the presence of Ziegler Natta Catalyst at high temperature and pressure is used to make buckets and dustbins.

Reason (R): High density polymers are closely packed and are chemically inert.

Choose the correct answer from the following:

- (1) (A) and (R) both are wrong.
- (2) Both (A) and (R) are correct and (R) is the correct explanation of (A)
- (3) (A) is correct but (R) is wrong
- (4) Both (A) and (R) are correct but (R) is not the correct explanation of (A).

Sol. 2

From ziegler - Natta catalyst HDPE is produced, HDPE is closely packed and are chemically inert, so used to make backet and dustbin.

CRASH COURSE

FOR JEE ADVANCED 2020

FREE Online Lectures Available on You Tube

Go Premium at ₹ 1100

Doubt Support ◆ Advanced Level Test Access

◆ Live Test Paper Discussion ◆ Final Revision Exercises

MOTION

17. A solution of two components containing n_1 moles of the 1^{st} component and n_2 moles of the 2^{nd} component is prepared. M₁ and M₂ are the molecular weights of component 1 and 2 respectively. If d is the density of the solution in g mL⁻¹, C_2 is the molarity and x_2 is the mole fraction of the 2^{nd} component, then C2 can be expressed as:

(1)
$$C_2 = \frac{dx_1}{M_2 + x_2(M_2 - M_1)}$$

(2)
$$C_2 = \frac{1000x_2}{M_1 + x_2(M_2 - M_1)}$$

(3)
$$C_2 = \frac{dx_2}{M_2 + x_2(M_2 - M_1)}$$

(4)
$$C_2 = \frac{1000 dx_2}{M_1 + x_2 (M_2 - M_1)}$$

Sol.

$$C_2 = \frac{X_2}{[X_2M_1 + (1 - X_2)M_2]/d} \times 1000$$

$$C_2 = \frac{1000 \, dx_2}{M_1 + (M_2 - M_1)x_2}$$

- 18. The correct statement with respect to dinitrogen is?
 - (1) Liquid dinitrogen is not used in cryosurgery.
 - (2) N₂ is paramagnetic in nature
 - (3) It can combine with dioxygen at 25°C
 - (4) It can be used as an inert diluent for reactive chemicals.

Sol.

- (1) Liquid nitrogen is used as a refrigerant to preserve biological material food items and in cryosurgery.
- (2) N_2 is diamagnetic, with no unpaired electrons.
- (3) N₂ does not combine with oxygen, hydrogen or most other elements. Nitrogen will combine with oxygen, however; in the presence of lightining or a spark.
- (4) In iron and chemical Industry inert diluent for reactive chemicals.
- Among the sulphates of alkaline earth metals, the solubilities of BeSO₄ and MgSO₄ in water, 19. respectively, are:

(1) Poor and high

(2) High and high

(3) Poor and poor

(4) High and poor

Sol.

Order of solubility of sulphate of Alkaline earth metals $BeSO_4 > MgSO_4 > CaSO_4 > SrSO_4 > BaSO_4$

FOR JEE ADVANCED 2020

FREE Online Lectures Available on You Tube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access

◆ Live Test Paper Discussion ◆ Final Revision Exercises

Motion^m

- **20.** The presence of soluble fluoride ion upto 1ppm concentration in drinking water, is :
 - (1) Harmful to skin

(2) Harmful to bones

(3) Safe for teeth

(4) Harmful for teeth

Sol. 3

Environmental chemistry - safe for teeth

- 21. A spherical balloon of radius 3cm containing helium gas has a pressure of 48×10^{-3} bar. At the same temperature, the pressure, of a spherical balloon of radius 12cm containing the same amount of gas will be...... $\times 10^{-6}$ bar.
- Sol. 750

moles =
$$\frac{48 \times 10^{-3} \times \frac{4}{3\pi} (3 \text{cm})^{3}}{R \times T}$$

$$moles = \frac{P \times \frac{4}{3\pi} (12cm)^3}{R T}$$

$$P \times 144 \times 12 = 48 \times 9 \times 3 \times 10^{-3}$$

$$P = \frac{27}{36} \times 10^{-3}$$

$$P = \frac{27000}{36} \times 10^{-6}$$

$$P = \frac{3000}{4} \times 10^{-6}$$

$$P = 750 \times 10^{-6} \, bar$$

[Assume 100% ionisation of the complex and $CaCl_2$, coordination number of Cr as 6, and that all NH_3 molecules are present inside the coordination sphere]

$$\Delta T_b = i \times K_b \times m$$

$$i \times 0.1 \times K_b = 3 \times 0.05 \times K_b \times 2$$

$$i = 3$$

$$[Cr(NH_3)_5. Cl] Cl_2 \rightarrow [Cr(NH_3)_5Cl]^{+2} + 2Cl^{-1}$$

$$x = 5$$

23. Potassium chlorate is prepared by the electrolysis of KCl in basic solution

 $60H^{-} + Cl^{-} \longrightarrow ClO_{3}^{-} + 3H_{2}O + 6e^{-}$

If only 60% of the current is utilized in the reaction, the time (rounded to the nearest hour) required to produce 10g of KClO₃ using a current of 2A is

(Given: $F = 96,500 \text{ C mol}^{-1}$; molar mass of $KClO_3 = 122g \text{ mol}^{-1}$)

CRASH COURSE

FOR JEE ADVANCED 2020

FREE Online Lectures Available on You Tube

Go Premium at ₹ 1100

Doubt Support ◆ Advanced Level Test Access
 Live Test Paper Discussion ◆ Final Revision Exercises



Sol.

$$\frac{10}{122} \times 6 = \frac{2 \times t(hr) \times 3600 \times 60\%}{96500}$$

$$t(hr) = \frac{96500}{122 \times 72} = 10.98 hr$$

= 11 hours

- 24. In an estimation of bromine by Carius method, 1.6 g of an organic compound gave 1.88 g of AgBr. The mass percentage of bromine in the compound is (Atomic mass, Ag=108, Br=80 g mol⁻¹)
- Sol. 50 %

Carius method

% of Br =
$$\frac{\text{wt of AgBr}}{\text{wt. of organic compound}} \times 100 \times \frac{\text{molar mass of Br}}{\text{AgBr}}$$

$$= \frac{1.88}{1.6} \times \frac{80}{188} \times 100 = \frac{15040}{300.8} = 50\%$$

- The number of CI = O bonds in perchloric acid is, "....." 25.
- Sol.

Admission **OPEN**

जब इन्होंने पूरा किया अपना सपना तो आप भी पा सकते है लक्ष्य अपना

JEE MAIN RESULT 2019









KOTA'S PIONEER IN DIGITAL EDUCATION 1,95,00,000+ viewers | 72,67,900+ viewing hours | 2,11,000+ Subscribers

SERVICES	SILVER	GOLD	PLATINUM
Classroom Lectures (VOD)			
Live interaction	NA		
Doubt Support	NA		
Academic & Technical Support	NA		
Complete access to all content	NA		
Classroom Study Material	NA		
Exercise Sheets	NA	S.	
Recorded Video Solutions	NA	8	
Online Test Series	NA		
Revision Material	NA		
Upgrade to Regular Classroom program	Chargeable	Chargeable	Free
Physical Classroom	NA	NA	
Computer Based Test	NA	NA	
Student Performance Report	NA	NA	
Workshop & Camp	NA	NA	
Motion Solution Lab- Supervised learning and instant doubt clearance	NA	NA	
Personalised guidance and mentoring	NA	NA	

•				
CLASS	SILVER	GOLD	PLATINUM	
7th/8th	FREE	₹ 12,000	₹ 35,000	
9th/10th	FREE	₹ 15,000	₹ 40,000	
11th	FREE	₹ 29,999	₹ 49,999	
12th	FREE	₹ 39,999	₹ 54,999	
12th Pass	FREE	₹ 39,999	₹ 59,999	

- SILVER (Trial) Only valid 7 DAYS or First 10 Hour's Lectures.
- GOLD (Online) can be converted to regular classroom (Any
- MOTION Center) by paying difference amount after lockdown. *** PLATINUM (Online + Regular) can be converted to regular
- classroom (Any MOTION Center) without any cost after

New Batch Starting from:

16 & 23 September 2020

Zero Cost EMI Available

