

हमारा विश्वास... हर एक विद्यार्थी है खास

**JEE
MAIN
Sept.
2020**

QUESTION PAPER WITH SOLUTION

MATHEMATICS _ 2 Sep. _ SHIFT - 2



MOTION™

H.O. : 394, Rajeev Gandhi Nagar, Kota
www.motion.ac.in | ✉: info@motion.ac.in

हमारा विश्वास... हर एक विद्यार्थी है खास

Q.1 Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function which satisfies $f(x+y) = f(x) + f(y) \forall x, y \in \mathbb{R}$. If $f(1) = 2$ and

$g(n) = \sum_{k=1}^{(n-1)} f(k), n \in \mathbb{N}$ then the value of n , for which $g(n) = 20$, is:

- (1) 9 (2) 5 (3) 4 (4) 20

Sol.

(2)
 $f(1) = 2 ; f(x+y) = f(x) + f(y)$
 $x = y = 1 \Rightarrow f(2) = 2 + 2 = 4$
 $x = 2, y = 1 \Rightarrow f(3) = 4 + 2 = 6$
 $g(n) = f(1) + f(2) + \dots + f(n-1)$
 $= 2 + 4 + 6 + \dots + 2(n-1)$
 $= 2 \sum (n-1)$
 $= 2 \frac{(n-1) \cdot n}{2}$
 $= n^2 - n$
 Given $g(n) = 20 \Rightarrow n^2 - n = 20$
 $n^2 - n - 20 = 0$
 $n = 5$

Q.2 If the sum of first 11 terms of an A.P., a_1, a_2, a_3, \dots is $0 (a_1 \neq 0)$ then the sum of the A.P., $a_1, a_3, a_5, \dots, a_{23}$ is ka_1 , where k is equal to:

- (1) $-\frac{121}{10}$ (2) $-\frac{72}{5}$ (3) $\frac{72}{5}$ (4) $\frac{121}{10}$

Sol.

(2)
 $\sum_{k=1}^{11} a_k = 0 \Rightarrow 11a + 55d = 0$
 $a + 5d = 0$
 Now $a_1 + a_3 + \dots + a_{23} = ka_1$
 $12a + d(2+4+6+\dots+22) = ka$
 $12a + 2d \cdot 66 = ka$
 $12(a+11d) = ka$
 $12 \left(a + 11 \left(-\frac{a}{5} \right) \right) = ka$
 $12 \left(1 - \frac{11}{5} \right) = k$
 $k = -\frac{72}{5}$

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on  YouTube

Go Premium at ₹ 1100

- ◆ Doubt Support ◆ Advanced Level Test Access
- ◆ Live Test Paper Discussion ◆ Final Revision Exercises


Start Date: **07 Sept. 2020**

हमारा विश्वास... हर एक विद्यार्थी है खास

Q.5 The area (in sq. units) of an equilateral triangle inscribed in the parabola $y^2 = 8x$, with one of its vertices on the vertex of this parabola, is:

- (1) $128\sqrt{3}$ (2) $192\sqrt{3}$ (3) $64\sqrt{3}$ (4) $256\sqrt{3}$

Sol. (2)



A : $(a \cos 30^\circ, a \sin 30^\circ)$
lies on parabola

$$\frac{a^2}{4} = 8 \cdot \frac{a \cdot \sqrt{3}}{2}$$

$$\boxed{a = 16\sqrt{3}}$$

$$\text{Area of equilateral } \Delta = \frac{\sqrt{3}}{4} a^2$$

$$\Delta = \frac{\sqrt{3}}{4} \cdot 16 \cdot 16 \cdot 3$$

$$\Delta = 192\sqrt{3}$$

Q.6 The imaginary part of $(3 + 2\sqrt{-54})^{1/2} - (3 - 2\sqrt{-54})^{1/2}$ can be :

- (1) $\sqrt{6}$ (2) $-2\sqrt{6}$ (3) 6 (4) $-\sqrt{6}$

Sol. (2)

$$(3 + 2i\sqrt{54})^{1/2} - (3 - 2i\sqrt{54})^{1/2}$$

$$= (9 + 6i^2 + 2 \cdot 3i\sqrt{6})^{1/2} - (9 + 6i^2 - 2 \cdot 3i\sqrt{6})^{1/2}$$

$$= \left((3 + \sqrt{6}i)^2 \right)^{1/2} - \left((3 - \sqrt{6}i)^2 \right)^{1/2}$$

$$= \pm(3 + \sqrt{6}i) \mp (3 - \sqrt{6}i) = -2\sqrt{6} i$$

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on **YouTube**


Go Premium at ₹ 1100

- ◆ Doubt Support ◆ Advanced Level Test Access
- ◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

Q.7 A plane passing through the point (3,1,1) contains two lines whose direction ratios are 1, -2, 2 and 2, 3, -1 respectively. If this plane also passes through the point (α , -3, 5), then α is equal to:
 (1) -5 (2) 10 (3) 5 (4) -10

Sol. (3)



$$\vec{n}_p = (-4, 5, 7)$$

Equation of plane :

$$P : -4(x-3) + 5(y-1) + 7(z-1) = 0$$

$$P : -4x + 5y + 7z + 12 - 5 - 7 = 0$$

$$P : 4x - 5y - 7z = 0$$

Pass (α , -3, 5)

$$4\alpha + 15 - 35 = 0$$

$$4\alpha = 20$$

$$\alpha = 5$$

Q.8 Let $A = \{X = (x, y, z)^T : PX = 0 \text{ and } x^2 + y^2 + z^2 = 1\}$, where $P = \begin{bmatrix} 1 & 2 & 1 \\ -2 & 3 & -4 \\ 1 & 9 & -1 \end{bmatrix}$, then the set A:

- (1) contains more than two elements
 (3) contains exactly two elements

- (2) is a singleton.
 (4) is an empty set.

Sol. (3)

$$X = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \text{ \& } x^2 + y^2 + z^2 = 1$$

$$PX = 0$$

$$\begin{bmatrix} 1 & 2 & 1 \\ -2 & 3 & -4 \\ 1 & 9 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$x + 2y + z = 0 \dots\dots(1)$$

$$-2x + 3y - 4z = 0 \dots(2)$$

$$x + 9y - z = 0 \dots\dots(3)$$

from (1) & (3)

$$\Rightarrow 2x + 11y = 0$$

from (1) & (2)

$$\Rightarrow 2x + 11y = 0$$

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

- ◆ Doubt Support ◆ Advanced Level Test Access
- ◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

हमारा विश्वास... हर एक विद्यार्थी है खास

$$\begin{aligned} &\text{from (2) \& (3)} \\ &-6x - 33y = 0 \\ &\Rightarrow 2x + 11y = 0 \\ &\text{put in (1)} \\ &-7y + 2z = 0 \end{aligned}$$

$$\text{Now } \left(\frac{11y}{2}\right)^2 + y^2 + \left(\frac{7y}{2}\right)^2 = 1$$

$$\begin{aligned} y^2(121 + 1 + 49) &= 4 \\ y^2(171) &= 4 \end{aligned}$$

$$y = \pm \frac{2}{\sqrt{171}} \Rightarrow x = \pm \frac{7}{\sqrt{171}} \Rightarrow z = \mp \frac{11}{\sqrt{171}} \Rightarrow \text{Only two pair possible}$$

- Q.9** The equation of the normal to the curve $y = (1+x)^{2y} + \cos^2(\sin^{-1}x)$ at $x=0$ is:
 (1) $y+4x=2$ (2) $2y+x=4$ (3) $x+4y=8$ (4) $y=4x+2$

Sol. (3)
 at $x = 0 \Rightarrow y = 1 + \cos^2(0) = 2$
 $p : (0, 2)$

$$\text{Now } y^1 = (1+x)^{2y} \left\{ \frac{2y}{1+x} + \ln(1+x) \cdot 2y \right\} - \sin 2(\sin^{-1}x) \cdot \frac{1}{\sqrt{1-x^2}}$$

$$y \Big|_{(0,2)} = 4 - 0$$

$$N_o : y - 2 = -\frac{1}{4}(x-0)$$


$$N_o : 4y - 8 = -x$$

$$N_o : \boxed{x + 4y = 8}$$

- Q.10** Consider a region $R = \{(x, y) \in \mathbb{R}^2 : x^2 \leq y \leq 2x\}$. If a line $y = \alpha$ divides the area of region R into two equal parts, then which of the following is true.?

- (1) $\alpha^3 - 6\alpha^2 + 16 = 0$ (2) $3\alpha^2 - 8\alpha^{3/2} + 8 = 0$
 (3) $\alpha^3 - 6\alpha^{3/2} - 16$ (4) $3\alpha^2 - 8\alpha + 8 = 0$

Sol. (2)



CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on **YouTube**

Go Premium at ₹ 1100

- ◆ Doubt Support ◆ Advanced Level Test Access
- ◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

$$A = \text{Area} = \int_0^2 (2x - x^2) dx = 4 - \frac{8}{3} = \frac{4}{3}$$

$$\text{Now } \int_0^{\alpha/2} (2x - x^2) dx + \int_{\alpha/2}^{\sqrt{\alpha}} (\alpha - x^2) dx = \frac{1}{2}A$$

$$\frac{\alpha^2}{4} - \frac{\alpha^3}{24} + \alpha(\sqrt{\alpha} - \alpha/2) - \left(\frac{\alpha\sqrt{\alpha}}{3} - \frac{\alpha^3}{24}\right) = \frac{4}{6}$$

$$\frac{\alpha^2}{4} + \alpha\sqrt{\alpha} - \frac{\alpha^2}{2} - \frac{\alpha\sqrt{\alpha}}{3} = \frac{4}{6}$$

$$-3\alpha^2 + 8\alpha\sqrt{\alpha} = 8$$

$$3\alpha^2 - 8\alpha\sqrt{\alpha} + 8 = 0$$

Q.11 Let $f : (-1, \infty) \rightarrow \mathbb{R}$ be defined by $f(0)=1$ and $f(x) = \frac{1}{x} \log_e(1+x), x \neq 0$. Then the function f :

- (1) increases in $(-1, \infty)$
- (2) decreases in $(-1, 0)$ and increases in $(0, \infty)$
- (3) increases in $(-1, 0)$ and decreases in $(0, \infty)$
- (4) decreases in $(-1, \infty)$.

Sol. (4)

$$f(x) = \frac{1}{x} \ln(1+x)$$

$$f' = \frac{x - \frac{1}{1+x} - \ln(1+x)}{x^2}$$

$$f' = \frac{1 - \frac{1}{1+x} - \ln(1+x)}{x^2}$$

$$f' < 0 \quad \forall x \in (-1, \infty)$$

Q.12 Which of the following is a tautology?

- (1) $(p \rightarrow q) \wedge (q \rightarrow p)$
- (2) $(\sim p) \wedge (p \vee q) \rightarrow q$
- (3) $(q \rightarrow p) \vee \sim(p \rightarrow q)$
- (4) $(\sim q) \vee (p \wedge q) \rightarrow q$

Sol. (2)

| $\sim p$ | $p \vee q$ | $\sim p \wedge (p \vee q)$ | $\sim p \wedge (p \vee q) \rightarrow q$ |
|----------|------------|----------------------------|--|
| F | T | F | T |
| F | T | F | T |
| T | T | T | T |
| T | F | F | T |

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

- ◆ Doubt Support ◆ Advanced Level Test Access
- ◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

हमारा विश्वास... हर एक विद्यार्थी है खास

Q.13 Let $f(x)$ be a quadratic polynomial such that $f(-1)+f(2)=0$. If one of the roots of $f(x)=0$ is 3, then its other roots lies in:

- (1) (0,1) (2) (1,3) (3) (-1,0) (4) (-3,-1)

Sol. (3)

$$\begin{aligned} \text{Let } f(x) &= a(x-3)(x-\alpha) \\ f(-1)+f(2) &= 0 \\ a[(-1-3)(-1-\alpha)+(2-3)(2-\alpha)] &= 0 \\ a[4+4\alpha-2+\alpha] &= 0 \\ 5\alpha+2 &= 0 \end{aligned}$$

$$\alpha = -\frac{2}{5}$$

Q.14 Let S be the sum of the first 9 terms of the series :

$$\{x+ka\} + \{x^2+(k+2)a\} + \{x^3+(k+4)a\} + \{x^4+(k+6)a\} + \dots \quad \text{where } a \neq 0 \quad \text{and } a \neq 1.$$

If $S = \frac{x^{10} - x + 45a(x-1)}{x-1}$, then k is equal to:

- (1) 3 (2) -3 (3) 1 (4) -5

Sol. (2)

$$\begin{aligned} S &= \{x+ka\} + \{x^2+(k+2)a\} + \{x^3+(k+4)a\} \text{ up to 9 term} \\ S &= (x+x^2+\dots+x^9) + a\{k+(k+2)+(k+4)+\dots\text{up to 9 term}\} \end{aligned}$$

$$S = \frac{x(1-x^9)}{1-x} + a\{9k+2.36\}$$

$$S = \frac{x^{10} - x}{x-1} + 9ak + 72a$$

$$S = \frac{x^{10} - x + 45a(x-1)}{x-1} = \frac{x^{10} - x + (9k+72)a(x-1)}{x-1}$$

$$= 45 = 9k + 72$$

$$9k = -27$$

$$k = -3$$

Q.15 The set of all possible values of θ in the interval $(0, \pi)$ for which the points $(1,2)$ and $(\sin \theta, \cos \theta)$ lie on the same side of the line $x+y=1$ is:

- (1) $\left(0, \frac{\pi}{4}\right)$ (2) $\left(0, \frac{\pi}{2}\right)$ (3) $\left(0, \frac{3\pi}{4}\right)$ (4) $\left(\frac{\pi}{4}, \frac{3\pi}{4}\right)$

Sol. (2)


CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**



$(\sin\theta, \cos\theta)$ lie on $x^2 + y^2 = 1$


Shaded points satisfy

$$\Rightarrow \theta \in (0, \pi / 2)$$

Q.16 Let $n > 2$ be an integer. Suppose that there are n Metro stations in a city located along a circular path. Each pair of stations is connected by a straight track only. Further, each pair of nearest stations is connected by blue line, whereas all remaining pairs of stations are connected by red line. If the number of red lines is 99 times the number of blue lines, then the value of n is:

- (1) 201 (2) 199 (3) 101 (4) 200

Sol. (1)



Red line = 99 blue line

$${}^n C_2 - n = 99n$$

$$\frac{n(n-1)}{2} = 100n$$

$$n-1 = 200$$

$$\boxed{n = 201}$$

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

- ◆ Doubt Support ◆ Advanced Level Test Access
- ◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

हमारा विश्वास... हर एक विद्यार्थी है खास

Q.17 If a curve $y=f(x)$, passing through the point $(1,2)$ is the solution of the differential equation, $2x^2dy=(2xy+y^2)dx$, then $f\left(\frac{1}{2}\right)$ is equal to:

(1) $\frac{-1}{1+\log_e 2}$

(2) $1+\log_e 2$

(3) $\frac{1}{1+\log_e 2}$

(4) $\frac{1}{1-\log_e 2}$

Sol. (3)

$$2 \frac{dy}{dx} = 2 \frac{y}{x} + \left(\frac{y}{x}\right)^2 \rightarrow \text{HDE}$$

$$\therefore y = vx$$

$$2\left(v + x \frac{dv}{dx}\right) = 2v + v^2$$

$$2 \frac{dv}{v^2} = \frac{dx}{x}$$

$$-\frac{2}{v} = \ln x + c$$

$$-\frac{2x}{y} = \ln x + c$$

$$\downarrow (1,2)$$

$$c = -1$$

$$c : \ln x + \frac{2x}{y} = 1$$

$$\text{For } f(1/2) \Rightarrow \ln\left(\frac{1}{2}\right) + \frac{2}{2y} = 1$$

$$y = \frac{1}{1+\ln 2}$$

Q.18 For some $\theta \in \left(0, \frac{\pi}{2}\right)$, if the eccentricity of the hyperbola, $x^2 - y^2 \sec^2 \theta = 10$ is $\sqrt{5}$ times the eccentricity of the ellipse, $x^2 \sec^2 \theta + y^2 = 5$, then the length of the latus rectum of the ellipse, is:

(1) $\frac{4\sqrt{5}}{3}$

(2) $\frac{2\sqrt{5}}{3}$

(3) $2\sqrt{6}$

(4) $\sqrt{30}$

Sol. (1)

$$H : x^2 - y^2 \sec^2 \theta = 10$$

$$E : x^2 \sec^2 \theta + y^2 = 5$$

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

$$\sqrt{1 + \frac{10 \cos^2 \theta}{10}} = \sqrt{5} \sqrt{1 - \frac{5 \cos^2 \theta}{5}}$$

$$1 + \cos^2 \theta = 5 - 5 \cos^2 \theta$$

$$6 \cos^2 \theta = 4$$

$$\cos \theta = \pm \sqrt{\frac{2}{3}}$$

$$l(\text{LR}) \text{ of ellipse} = \frac{2.5 \cos^2 \theta}{\sqrt{5}}$$

$$= 2\sqrt{5} \cdot \frac{2}{3} = \boxed{\frac{4\sqrt{5}}{3}}$$

Q.19 $\lim_{x \rightarrow 0} \left(\tan \left(\frac{\pi}{4} + x \right) \right)^{1/x}$ is equal to:

(1) e

(2) e²

(3) 2

(4) 1

Sol.

$$\lim_{x \rightarrow 0} \left(\tan \left(\frac{\pi}{4} + x \right) \right)^{1/x} (1^\infty) = e^L$$

$$L = \lim_{x \rightarrow 0} \frac{\tan \left(\frac{\pi}{4} + x \right) - 1}{x}$$

$$L = \lim_{x \rightarrow 0} \frac{1 + \tan x}{1 - \tan x} - 1$$

$$L = \lim_{x \rightarrow 0} 2 \left(\frac{\tan x}{x} \right) \cdot \left(\frac{1}{1 - \tan x} \right)$$

$$L = +2$$

Ans. e²

Q.20 Let a, b, c ∈ R be all non-zero and satisfy a³+b³+c³ = 2. If the matrix

$$A = \begin{pmatrix} a & b & c \\ b & c & a \\ c & a & b \end{pmatrix}$$

satisfies A^TA=I, then a value of abc can be:

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

हमारा विश्वास... हर एक विद्यार्थी है खास

(1) $\frac{2}{3}$

(2) 3

(3) $-\frac{1}{3}$

(4) $\frac{1}{3}$

Sol.

(4)

$$a^3 + b^3 + c^3 = 2$$

$$A^T A = I$$

$$\begin{bmatrix} a & b & c \\ b & c & a \\ c & a & b \end{bmatrix} \begin{bmatrix} a & b & c \\ b & c & a \\ c & a & b \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= a^2 + b^2 + c^2 = 1$$

$$\& ab + bc + ca = 0$$

$$\text{Now } (a+b+c)^2 = \sum a^2 + 2\sum ab$$

$$(\sum a)^2 = 1 + 0 \Rightarrow (\sum a)^2 = 1 \Rightarrow \sum a = \pm 1$$

$$\text{Now } \sum a^3 - 3abc = (\sum a)(\sum a^2 - \sum ab)$$

$$2 - 3abc = \pm 1(1 - 0)$$

$$2 - 3abc = \pm 1$$

$$\begin{array}{l} \begin{array}{c} (+) \quad \quad \quad (-) \\ \diagdown \quad \diagup \\ 3abc = 1 \quad 3abc = 3 \\ \boxed{abc = \frac{1}{3}} \quad \boxed{abc = 1} \end{array} \end{array}$$

Q.21 Let the position vectors of points 'A' and 'B' be $\hat{i} + \hat{j} + \hat{k}$ and $2\hat{i} + \hat{j} + 3\hat{k}$, respectively. A point 'P' divides the line segment AB internally in the ratio $\lambda : 1 (\lambda > 0)$. If O is the origin and

$$\vec{OB} \cdot \vec{OP} - 3 \left| \vec{OA} \times \vec{OP} \right|^2 = 6, \text{ then } \lambda \text{ is equal to } \underline{\hspace{2cm}}$$

Sol. 0.8

$$\vec{OA} = \langle 1, 1, 1 \rangle, \vec{OB} = \langle 2, 1, 3 \rangle$$

$$\begin{array}{c} \lambda \quad \quad \quad 1 \\ \text{A} \quad \quad \quad \text{P} \quad \quad \quad \text{B} \end{array}$$

$$\vec{OP} = \left(\frac{2\lambda + 1}{\lambda + 1}, 1, \frac{3\lambda + 1}{\lambda + 1} \right)$$

$$\vec{OB} \cdot \vec{OP} = \frac{2(2\lambda + 1)}{\lambda + 1} + 1 + \frac{3(3\lambda + 1)}{\lambda + 1}$$

$$= \frac{14\lambda + 6}{\lambda + 1}$$

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

$$\begin{aligned}
 |\overline{OA} \times \overline{OP}|^2 &= |\overline{OA}|^2 |\overline{OP}|^2 - (\overline{OA} \cdot \overline{OP})^2 \\
 3 \cdot \left(\frac{(2\lambda + 1)^2 + (\lambda + 1)^2 + (3\lambda + 1)^2}{(\lambda + 1)^2} \right) &- \left(\frac{2\lambda + 1 + \lambda + 1 + 3\lambda + 1}{\lambda + 1} \right)^2 \\
 &= \frac{1}{(\lambda + 1)^2} \{ 3(14\lambda^2 + 12\lambda + 3) - (6\lambda + 3)^2 \} \\
 &= \frac{1}{(\lambda + 1)^2} \{ 6\lambda^2 \}
 \end{aligned}$$

$$\text{Now } \frac{14\lambda + 6}{\lambda + 1} - 3 \left(\frac{6\lambda^2}{(\lambda + 1)^2} \right) = 6$$

$$\begin{aligned}
 (14\lambda + 6)(\lambda + 1) - 18\lambda^2 &= 6(\lambda + 1)^2 \\
 -4\lambda^2 + 20\lambda + 6 &= 6\lambda^2 + 12\lambda + 6 \\
 10\lambda^2 - 8\lambda &= 0 \\
 \lambda(10\lambda - 8) &= 0
 \end{aligned}$$

$$\therefore \lambda > 0$$

$$\boxed{\lambda = .8}$$

Q.22 Let $[t]$ denote the greatest integer less than or equal to t . Then the value of

$$\int_1^2 |2x - [3x]| dx \text{ is } \underline{\hspace{2cm}}$$

Sol. 1

$$\begin{aligned}
 &\int_1^2 |2x - [3x]| dx \\
 3x &= t \\
 &= \frac{1}{3} \int_3^6 \left| \frac{2t}{3} - [t] \right| dt \\
 &= \frac{1}{9} \left[\int_3^4 |2t - 3[t]| dt \right] \\
 &= \frac{1}{9} \left[\int_3^4 |2t - 9| dt + \int_4^5 |2t - 12| dt + \int_5^6 |2t - 15| dt \right] \\
 &= \frac{1}{9} \left[\int_3^4 (9 - 2t) dt + \int_4^5 (12 - 2t) dt + \int_5^6 (15 - 2t) dt \right]
 \end{aligned}$$

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

हमारा विश्वास... हर एक विद्यार्थी है खास

$$\begin{aligned}
 &= \frac{1}{9} [9 \cdot 1 + 12 \cdot 1 + 15 \cdot 1 - [4^2 - 3^2] - [5^2 - 4^2] - [6^2 - 5^2]] \\
 &= \frac{1}{9} [36 - [4^2 - 3^2 + 5^2 - 4^2 + 6^2 - 5^2]] \\
 &= \frac{1}{9} [36 - 36 + 9] = 1
 \end{aligned}$$

Q.23 If $y = \sum_{k=1}^6 k \cos^{-1} \left\{ \frac{3}{5} \cos kx - \frac{4}{5} \sin kx \right\}$, then $\frac{dy}{dx}$ at $x=0$ is _____

Sol. 91

$$y = \sum_{k=1}^6 k \cos^{-1} \{ \cos(kx + \theta) \}$$

$$\text{where } \tan \theta = \frac{4}{3}$$

$$y = \cos^{-1}(\cos(x+\theta)) + 2\cos^{-1}(\cos(2x+\theta)) + \dots + 6\cos^{-1}(\cos(6x+\theta))$$

$$\begin{aligned}
 \left. \frac{dy}{dx} \right|_{x=0} &= \frac{\sin(\theta)}{\sqrt{1 - \cos^2 \theta}} + \dots \\
 &= 1 \cdot 1 + 2 \cdot 2 + 3 \cdot 3 + \dots + 6 \cdot 6 \\
 &= \sum 6^2 = \frac{6 \cdot 7 \cdot 13}{6} = 91
 \end{aligned}$$

Q.24 If the variance of the terms in an increasing A.P., $b_1, b_2, b_3, \dots, b_{11}$ is 90, then the common difference of this A.P. is _____

Sol. 3

$$\text{Var}(x) = \frac{\sum bi^2}{11} - \left(\frac{\sum bi}{11} \right)^2$$

$$90 = \frac{a^2 + (a+d)^2 + (a+2d)^2 + \dots + (a+10d)^2}{11}$$

$$\left(\frac{a + a + d + a + 2d + \dots + (a + 10d)}{11} \right)^2$$

$$10890 = 11 \{ 11a^2 + 385d^2 + 110ad \} - \{ 11a + 55d \}^2$$

$$10890 = 1210d^2$$

$$d = 3$$

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on YouTube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

Q.25 For a positive integer n , $\left(1 + \frac{1}{x}\right)^n$ is expanded in increasing powers of x . If three consecutive coefficients in this expansion are in the ratio, 2:5:12, then n is equal to _____

Sol. 118

Let 3 consecutive coH are

$${}^n C_{r-1} : {}^n C_r : {}^n C_{r+1} :: 2:5:12$$

$$\frac{{}^n C_{r-1}}{{}^n C_r} = \frac{2}{5} \quad \& \quad \frac{{}^n C_r}{{}^n C_{r+1}} = \frac{5}{12}$$

$$\frac{r}{n-r+1} = \frac{2}{5} \quad \& \quad \frac{r+1}{(n-r)} = \frac{5}{12}$$

$$7r = 2n + 2 \quad \& \quad 17r = 5n - 12$$

$$\Rightarrow \frac{2n+2}{7} = \frac{5n-12}{17}$$

$$= 34n + 34 = 35n - 84$$

$$\Rightarrow n = 118$$

CRASH COURSE
FOR JEE ADVANCED 2020

FREE Online Lectures Available on  YouTube

Go Premium at ₹ 1100

◆ Doubt Support ◆ Advanced Level Test Access
◆ Live Test Paper Discussion ◆ Final Revision Exercises

Start Date: **07 Sept. 2020**

जब इन्होंने पूरा किया अपना सपना
तो आप भी पा सकते है लक्ष्य अपना

Admission
OPEN

JEE MAIN RESULT 2019



Nitin Gupta

Marks
335
13th (2019)

Marks
149
12th (2018)



Shiv Modi

Marks
318
13th (2019)


Marks
153
12th (2018)



Ritik Bansal

Marks
308
13th (2019)

Marks
218
12th (2018)



Shubham Kumar

Marks
300
13th (2019)

Marks
153
12th (2018)

KOTA'S PIONEER IN DIGITAL EDUCATION

1,95,00,000+ viewers | **72,67,900+** viewing hours | **2,11,000+** Subscribers

| SERVICES | SILVER | GOLD | PLATINUM |
|--|-------------------|-------------------|-------------|
| Classroom Lectures (VOD) | | | |
| Live interaction | NA | | |
| Doubt Support | NA | | |
| Academic & Technical Support | NA | | |
| Complete access to all content | NA | | |
| Classroom Study Material | NA | | |
| Exercise Sheets | NA | | |
| Recorded Video Solutions | NA | | |
| Online Test Series | NA | | |
| Revision Material | NA | | |
| Upgrade to Regular Classroom program | Chargeable | Chargeable | Free |
| Physical Classroom | NA | NA | |
| Computer Based Test | NA | NA | |
| Student Performance Report | NA | NA | |
| Workshop & Camp | NA | NA | |
| Motion Solution Lab- Supervised learning and instant doubt clearance | NA | NA | |
| Personalised guidance and mentoring | NA | NA | |

FEE STRUCTURE

| CLASS | SILVER | GOLD | PLATINUM |
|-----------|--------|----------|----------|
| 7th/8th | FREE | ₹ 12,000 | ₹ 35,000 |
| 9th/10th | FREE | ₹ 15,000 | ₹ 40,000 |
| 11th | FREE | ₹ 29,999 | ₹ 49,999 |
| 12th | FREE | ₹ 39,999 | ₹ 54,999 |
| 12th Pass | FREE | ₹ 39,999 | ₹ 59,999 |

+ Student Kit will be provided at extra cost to Platinum Student.

* **SILVER (Trial)** Only valid 7 DAYS or First 10 Hour's Lectures.

** **GOLD (Online)** can be converted to regular classroom (Any MOTION Center) by paying difference amount after lockdown.

*** **PLATINUM (Online + Regular)** can be converted to regular classroom (Any MOTION Center) without any cost after lockdown.

New Batch Starting from :
16 & 23 September 2020

Zero Cost EMI Available

MOTION™

H.O. : 394, Rajeev Gandhi Nagar, Kota
www.motion.ac.in | ✉ : info@motion.ac.in